

# APPLICATION FOR PATENT

5 Inventors: Iris Pecker, Israel Vlodavsky and Elena Feinstein

10 Title: POLYNUCLEOTIDE ENCODING A POLYPEPTIDE  
HAVING HEPARANASE ACTIVITY AND EXPRESSION  
OF SAME IN GENETICALLY MODIFIED CELLS

15 This is a continuation of U.S. Patent Application No. 09/776,874,  
filed February 6, 2001, which is a continuation of U.S. Patent Application  
No. 09/258,892, filed March 1, 1999, which is a continuation-in-part of  
20 PCT/US98/17954, filed August 31, 1998, which claims priority from U.S.  
Patent Application 09/109,386, filed July 2, 1998, now abandoned, which is  
a continuation-in-part of U.S. Patent Application 08/922,170, filed  
September 2, 1997, now, U.S. Patent No. 5,968,822.

## 25 FIELD AND BACKGROUND OF THE INVENTION

The present invention relates to a polynucleotide, referred to  
hereinbelow as *hpa*, encoding a polypeptide having heparanase activity,  
vectors (nucleic acid constructs) including same and genetically modified  
30 cells expressing heparanase. The invention further relates to a recombinant  
protein having heparanase activity and to antisense oligonucleotides,  
constructs and ribozymes for down regulating heparanase activity. In  
addition, the invention relates to heparanase promoter sequences and their  
uses.

*Heparan sulfate proteoglycans:* Heparan sulfate proteoglycans (HSPG) are ubiquitous macromolecules associated with the cell surface and extra cellular matrix (ECM) of a wide range of cells of vertebrate and invertebrate tissues (1-4). The basic HSPG structure includes a protein core to which several linear heparan sulfate chains are covalently attached. These polysaccharide chains are typically composed of repeating hexuronic and D-glucosamine disaccharide units that are substituted to a varying extent with N- and O-linked sulfate moieties and N-linked acetyl groups (1-4). Studies on the involvement of ECM molecules in cell attachment, growth and differentiation revealed a central role of HSPG in embryonic morphogenesis, angiogenesis, neurite outgrowth and tissue repair (1-5). HSPG are prominent components of blood vessels (3). In large blood vessels they are concentrated mostly in the intima and inner media, whereas in capillaries they are found mainly in the subendothelial basement membrane where they support proliferating and migrating endothelial cells and stabilize the structure of the capillary wall. The ability of HSPG to interact with ECM macromolecules such as collagen, laminin and fibronectin, and with different attachment sites on plasma membranes suggests a key role for this proteoglycan in the self-assembly and insolubility of ECM components, as well as in cell adhesion and locomotion. Cleavage of the heparan sulfate (HS) chains may therefore result in degradation of the subendothelial ECM and hence may play a

decisive role in extravasation of blood-borne cells. HS catabolism is observed in inflammation, wound repair, diabetes, and cancer metastasis, suggesting that enzymes which degrade HS play important roles in pathologic processes. Heparanase activity has been described in activated immune system cells and highly metastatic cancer cells (6-8), but research has been handicapped by the lack of biologic tools to explore potential causative roles of heparanase in disease conditions.

*Involvement of Heparanase in Tumor Cell Invasion and Metastasis:* Circulating tumor cells arrested in the capillary beds of different organs must invade the endothelial cell lining and degrade its underlying basement membrane (BM) in order to invade into the extravascular tissue(s) where they establish metastasis (9, 10). Metastatic tumor cells often attach at or near the intercellular junctions between adjacent endothelial cells. Such attachment of the metastatic cells is followed by rupture of the junctions, retraction of the endothelial cell borders and migration through the breach in the endothelium toward the exposed underlying BM (9). Once located between endothelial cells and the BM, the invading cells must degrade the subendothelial glycoproteins and proteoglycans of the BM in order to migrate out of the vascular compartment. Several cellular enzymes (e.g., collagenase IV, plasminogen activator, cathepsin B, elastase, etc.) are thought to be involved in degradation of BM (10). Among these enzymes is an endo- $\beta$

-D-glucuronidase (heparanase) that cleaves HS at specific intrachain sites (6, 8, 11). Expression of a HS degrading heparanase was found to correlate with the metastatic potential of mouse lymphoma (11), fibrosarcoma and melanoma (8) cells. Moreover, elevated levels of heparanase were detected in sera from metastatic tumor bearing animals and melanoma patients (8) and in tumor biopsies of cancer patients (12).

The control of cell proliferation and tumor progression by the local microenvironment, focusing on the interaction of cells with the extracellular matrix (ECM) produced by cultured corneal and vascular endothelial cells, was investigated previously by the present inventors. This cultured ECM closely resembles the subendothelium *in vivo* in its morphological appearance and molecular composition. It contains collagens (mostly type III and IV, with smaller amounts of types I and V), proteoglycans (mostly heparan sulfate- and dermatan sulfate- proteoglycans, with smaller amounts of chondroitin sulfate proteoglycans), laminin, fibronectin, entactin and elastin (13, 14). The ability of cells to degrade HS in the cultured ECM was studied by allowing cells to interact with a metabolically sulfate labeled ECM, followed by gel filtration (Sepharose 6B) analysis of degradation products released into the culture medium (11). While intact HSPG are eluted next to the void volume of the column ( $K_{av} < 0.2$ ,  $M_r \sim 0.5 \times 10^6$ ), labeled degradation fragments of HS side chains



are eluted more toward the  $V_t$  of the column ( $0.5 < k_{av} < 0.8$ ,  $M_r = 5-7 \times 10^3$ ) (11).

The heparanase inhibitory effect of various non-anticoagulant species of heparin that might be of potential use in preventing extravasation of blood-borne cells was also investigated by the present inventors. Inhibition of heparanase was best achieved by heparin species containing 16 sugar units or more and having sulfate groups at both the N and O positions. While O-desulfation abolished the heparanase inhibiting effect of heparin, O-sulfated, N-acetylated heparin retained a high inhibitory activity, provided that the N-substituted molecules had a molecular size of about 4,000 daltons or more (7). Treatment of experimental animals with heparanase inhibitors (e.g., non-anticoagulant species of heparin) markedly reduced (>90%) the incidence of lung metastases induced by B16 melanoma, Lewis lung carcinoma and mammary adenocarcinoma cells (7, 8, 16). Heparin fractions with high and low affinity to anti-thrombin III exhibited a comparable high anti-metastatic activity, indicating that the heparanase inhibiting activity of heparin, rather than its anticoagulant activity, plays a role in the anti-metastatic properties of the polysaccharide (7).

*Heparanase activity in the urine of cancer patients:* In an attempt to further elucidate the involvement of heparanase in tumor progression and its relevance to human cancer, urine samples for heparanase activity

were screened (16a). Heparanase activity was detected in the urine of some, but not all, cancer patients. High levels of heparanase activity were determined in the urine of patients with an aggressive metastatic disease and there was no detectable activity in the urine of healthy donors.

5       Heparanase activity was also found in the urine of 20% of normal and microalbuminuric insulin dependent diabetes mellitus (IDDM) patients, most likely due to diabetic nephropathy, the most important single disorder leading to renal failure in adults.

*Possible involvement of heparanase in tumor angiogenesis:*

10       Fibroblast growth factors are a family of structurally related polypeptides characterized by high affinity to heparin (17). They are highly mitogenic for vascular endothelial cells and are among the most potent inducers of neovascularization (17, 18). Basic fibroblast growth factor (bFGF) has been extracted from the subendothelial ECM produced *in vitro* (19) and from  
15       basement membranes of the cornea (20), suggesting that ECM may serve as a reservoir for bFGF. Immunohistochemical staining revealed the localization of bFGF in basement membranes of diverse tissues and blood vessels (21). Despite the ubiquitous presence of bFGF in normal tissues, endothelial cell proliferation in these tissues is usually very low, suggesting  
20       that bFGF is somehow sequestered from its site of action. Studies on the interaction of bFGF with ECM revealed that bFGF binds to HSPG in the ECM and can be released in an active form by HS degrading enzymes (15,

20, 22). It was demonstrated that heparanase activity expressed by platelets, mast cells, neutrophils, and lymphoma cells is involved in release of active bFGF from ECM and basement membranes (23), suggesting that heparanase activity may not only function in cell migration and invasion, but may also elicit an indirect neovascular response. These results suggest that the ECM HSPG provides a natural storage depot for bFGF and possibly other heparin-binding growth promoting factors (24, 25). Displacement of bFGF from its storage within basement membranes and ECM may therefore provide a novel mechanism for induction of neovascularization in normal and pathological situations.

Recent studies indicate that heparin and HS are involved in binding of bFGF to high affinity cell surface receptors and in bFGF cell signaling (26, 27). Moreover, the size of HS required for optimal effect was similar to that of HS fragments released by heparanase (28). Similar results were obtained with vascular endothelial cells growth factor (VEGF) (29), suggesting the operation of a dual receptor mechanism involving HS in cell interaction with heparin-binding growth factors. It is therefore proposed that restriction of endothelial cell growth factors in ECM prevents their systemic action on the vascular endothelium, thus maintaining a very low rate of endothelial cells turnover and vessel growth. On the other hand, release of bFGF from storage in ECM as a complex with HS fragment, may elicit localized endothelial cell proliferation and neovascularization in

processes such as wound healing, inflammation and tumor development (24, 25).

*Expression of heparanase by cells of the immune system:*

Heparanase activity correlates with the ability of activated cells of the immune system to leave the circulation and elicit both inflammatory and autoimmune responses. Interaction of platelets, granulocytes, T and B lymphocytes, macrophages and mast cells with the subendothelial ECM is associated with degradation of HS by a specific heparanase activity (6).

The enzyme is released from intracellular compartments (e.g., lysosomes, specific granules, etc.) in response to various activation signals (e.g., thrombin, calcium ionophore, immune complexes, antigens, mitogens, etc.), suggesting its regulated involvement in inflammation and cellular immunity.

*Some of the observations regarding the heparanase enzyme were reviewed in reference No. 6 and are listed hereinbelow:*

First, a proteolytic activity (plasminogen activator) and heparanase participate synergistically in sequential degradation of the ECM HSPG by inflammatory leukocytes and malignant cells.

Second, a large proportion of the platelet heparanase exists in a latent form, probably as a complex with chondroitin sulfate. The latent enzyme is activated by tumor cell-derived factor(s) and may then facilitate cell invasion through the vascular endothelium in the process of tumor metastasis.

Third, release of the platelet heparanase from  $\alpha$ -granules is induced by a strong stimulant (i.e., thrombin), but not in response to platelet activation on ECM.

Fourth, the neutrophil heparanase is preferentially and readily  
5 released in response to a threshold activation and upon incubation of the cells on ECM.

Fifth, contact of neutrophils with ECM inhibited release of noxious enzymes (proteases, lysozyme) and oxygen radicals, but not of enzymes (heparanase, gelatinase) which may enable diapedesis. This protective role  
10 of the subendothelial ECM was observed when the cells were stimulated with soluble factors but not with phagocytosable stimulants.

Sixth, intracellular heparanase is secreted within minutes after exposure of T cell lines to specific antigens.

Seventh, mitogens (Con A, LPS) induce synthesis and secretion of  
15 heparanase by normal T and B lymphocytes maintained *in vitro*. T lymphocyte heparanase is also induced by immunization with antigen *in vivo*.

Eighth, heparanase activity is expressed by pre-B lymphomas and B-lymphomas, but not by plasmacytomas and resting normal B  
20 lymphocytes.

Ninth, heparanase activity is expressed by activated macrophages during incubation with ECM, but there was little or no release of the

enzyme into the incubation medium. Similar results were obtained with human myeloid leukemia cells induced to differentiate to mature macrophages.

Tenth, T-cell mediated delayed type hypersensitivity and experimental autoimmunity are suppressed by low doses of heparanase inhibiting non-anticoagulant species of heparin (30).

Eleventh, heparanase activity expressed by platelets, neutrophils and metastatic tumor cells releases active bFGF from ECM and basement membranes. Release of bFGF from storage in ECM may elicit a localized neovascular response in processes such as wound healing, inflammation and tumor development.

Twelfth, among the breakdown products of the ECM generated by heparanase is a tri-sulfated disaccharide that can inhibit T-cell mediated inflammation *in vivo* (31). This inhibition was associated with an inhibitory effect of the disaccharide on the production of biologically active TNF $\alpha$  by activated T cells *in vitro* (31).

***Other potential therapeutic applications:*** Apart from its involvement in tumor cell metastasis, inflammation and autoimmunity, mammalian heparanase may be applied to modulate: bioavailability of heparin-binding growth factors (15); cellular responses to heparin-binding growth factors (e.g., bFGF, VEGF) and cytokines (IL-8) (31a, 29); cell interaction with plasma lipoproteins (32); cellular susceptibility to certain

viral and some bacterial and protozoa infections (33, 33a, 33b); and disintegration of amyloid plaques (34). Heparanase may thus prove useful for conditions such as wound healing, angiogenesis, restenosis, atherosclerosis, inflammation, neurodegenerative diseases and viral infections. Mammalian heparanase can be used to neutralize plasma heparin, as a potential replacement of protamine. Anti-heparanase antibodies may be applied for immunodetection and diagnosis of micrometastases, autoimmune lesions and renal failure in biopsy specimens, plasma samples, and body fluids. Common use in basic research is expected.

The identification of the *hpa* gene encoding for heparanase enzyme will enable the production of a recombinant enzyme in heterologous expression systems. Availability of the recombinant protein will pave the way for solving the protein structure function relationship and will provide a tool for developing new inhibitors.

***Viral Infection:*** The presence of heparan sulfate on cell surfaces have been shown to be the principal requirement for the binding of Herpes Simplex (33) and Dengue (33a) viruses to cells and for subsequent infection of the cells. Removal of the cell surface heparan sulfate by heparanase may therefore abolish virus infection. In fact, treatment of cells with bacterial heparitinase (degrading heparan sulfate) or heparinase (degrading heparan) reduced the binding of two related animal herpes viruses to cells and

rendered the cells at least partially resistant to virus infection (33). There are some indications that the cell surface heparan sulfate is also involved in HIV infection (33b).

*Neurodegenerative diseases:* Heparan sulfate proteoglycans were identified in the prion protein amyloid plaques of Genstmann-Straussler Syndrome, Creutzfeldt-Jakob disease and Scrape (34). Heparanase may disintegrate these amyloid plaques which are also thought to play a role in the pathogenesis of Alzheimer's disease.

*Restenosis and Atherosclerosis:* Proliferation of arterial smooth muscle cells (SMCs) in response to endothelial injury and accumulation of cholesterol rich lipoproteins are basic events in the pathogenesis of atherosclerosis and restenosis (35). Apart from its involvement in SMC proliferation (i.e., low affinity receptors for heparin-binding growth factors), HS is also involved in lipoprotein binding, retention and uptake (36). It was demonstrated that HSPG and lipoprotein lipase participate in a novel catabolic pathway that may allow substantial cellular and interstitial accumulation of cholesterol rich lipoproteins (32). The latter pathway is expected to be highly atherogenic by promoting accumulation of apoB and apoE rich lipoproteins (i.e. LDL, VLDL, chylomicrons), independent of feed back inhibition by the cellular sterol content. Removal of SMC HS by heparanase is therefore expected to inhibit both SMC proliferation and lipid



accumulation and thus may halt the progression of restenosis and atherosclerosis.

***Gene therapy:***

The ultimate goal in the management of inherited as well as acquired  
5 diseases is a rational therapy with the aim to eliminate the underlying biochemical defects associated with the disease rather than symptomatic treatment. Gene therapy is a promising candidate to meet these objectives. Initially it was developed for treatment of genetic disorders, however, the consensus view today is that it offers the prospect of providing therapy for a  
10 variety of acquired diseases, including cancer, viral infections, vascular diseases and neurodegenerative disorders.

The gene-based therapeutic can act either intracellularly, affecting only the cells to which it is delivered, or extracellularly, using the recipient cells as local endogenous factories for the therapeutic product(s). The  
15 application of gene therapy may follow any of the following strategies: (i) prophylactic gene therapy, such as using gene transfer to protect cells against viral infection; (ii) cytotoxic gene therapy, such as cancer therapy, where genes encode cytotoxic products to render the target cells vulnerable to attack by the normal immune response; (iii) biochemical correction,  
20 primarily for the treatment of single gene defects, where a normal copy of the gene is added to the affected or other cells.

To allow efficient transfer of the therapeutic genes, a variety of gene delivery techniques have been developed based on viral and non-viral vector systems. The most widely used and most efficient systems for delivering genetic material into target cells are viral vectors. So far, 329 clinical studies (phase I, I/II and II) with over 2,500 patients have been initiated Worldwide since 1989 (50).

The approach of gene addition pose serious barriers. The expression of many genes is tightly regulated and context dependent, so achieving the correct balance and function of expression is challenging. The gene itself is often quite large, containing many exons and introns. The delivery vector is usually a virus, which can infect with a high efficiency but may, on the other hand, induce immunological response and consequently decreases effectiveness, especially upon secondary administration. Most of the current expression vector-based gene therapy protocols fail to achieve clinically significant transgene expression required for treating genetic diseases. Apparently, it is difficult to deliver enough virus to the right cell type to elicit an effective and therapeutic effect (51)

Homologous recombination, which was initially considered to be of limited use for gene therapy because of its low frequency in mammalian cells, has recently emerged as a potential strategy for developing gene therapy. Different approaches have been used to study homologous recombination in mammalian cells; some involve DNA repair mechanisms.

These studies aimed at either gene disruption or gene correction and include RNA/DNA chimeric oligonucleotides, small or large homologous DNA fragments, or adeno-associated viral vectors. Most of these studies show a reasonable frequency of homologous recombination, which warrants further *in vivo* testing (52). Homologous recombination-based gene therapy has the potential to develop into a powerful therapeutic modality for genetic diseases. It can offer permanent expression and normal regulation of corrected genes in appropriate cells or organs and probably can be used for treating dominantly inherited diseases such as polycystic kidney disease.

***Genomic sequences function in regulation of gene expression:***

The efficient expression of therapeutic genes in target cells or tissues is an important component of efficient and safe gene therapy. The expression of genes is driven by the promoter region upstream of the coding sequence, although regulation of expression may be supplemented by farther upstream or downstream DNA sequences or DNA in the introns of the gene. Since this important information is embedded in the DNA, the description of gene structure is crucial to the analysis of gene regulation. Characterization of cell specific or tissue specific promoters, as well as other tissue specific regulatory elements enables the use of such sequences to direct efficient cell specific, or developmental stage specific gene expression. This information provides the basis for targeting individual genes and for control of their expression by exogenous agents, such as

drugs. Identification of transcription factors and other regulatory proteins required for proper gene expression will point at new potential targets for modulating gene expression, when so desired or required.

Efficient expression of many mammalian genes depends on the  
5 presence of at least one intron. The expression of mouse thymidylate synthase (TS) gene, for example, is greatly influenced by intron sequences. The addition of almost any of the introns from the mouse TS gene to an intronless TS minigene leads to a large increase in expression (42). The involvement of intron 1 in the regulation of expression was demonstrated  
10 for many other genes. In human factor IX (hFIX), intron 1 is able to increase the expression level about 3 fold more as compared to that of the hFIX cDNA (43). The expression enhancing activity of intron 1 is due to efficient functional splicing sequences, present in the precursor mRNA. By being efficiently assembled into spliceosome complexes, transcripts with  
15 splicing sequences may be better protected in the nucleus from random degradations, than those without such sequences (44).

A forward-inserted intron1-carrying hFIX expression cassette suggested to be useful for directed gene transfer, while for retroviral-mediated gene transfer system, reversely-inserted intron  
20 1-carrying hFIX expression cassette was considered (43).

A highly conserved cis-acting sequence element was identified in the first intron of the mouse and rat c-Ha-ras, and in the first exon of Ha- and

Ki-ras genes of human, mouse and rat. This cis-acting regulatory sequence confers strong transcription enhancer activity that is differentially modulated by steroid hormones in metastatic and nonmetastatic subpopulations. Perturbations in the regulatory activities of such cis-acting sequences may play an important role in governing oncogenic potency of Ha-ras through transcriptional control mechanisms (45).

Intron sequences affect tissue specific, as well as inducible gene expression. A 182 bp intron 1 DNA segment of the mouse Col2a1 gene contains the necessary information to confer high-level, temporally correct, chondrocyte expression on a reporter gene in intact mouse embryos, while Col2a1 promoter sequences are dispensable for chondrocyte expression (46). In Col1A1 gene the intron plays little or no role in constitutive expression of collagen in the skin, and in cultured cells derived from the skin, however, in the lungs of young mice, intron deletion results in decrease of expression to less than 50 % (47).

A classical enhancer activity was shown in the 2 kb intron fragment in bovine beta-casein gene. The enhancer activity was largely dependent on the lactogenic hormones, especially prolactin. It was suggested that several elements in the intron-1 of the bovine beta-casein gene cooperatively interact not only with each other but also with its promoter for hormonal induction (48).

Identification and characterization of regulatory elements in genomic non-coding sequences, such as introns, provides a tool for designing and constructing novel vectors for tissue specific, hormone regulated or any other defined expression pattern, for gene therapy. Such an expression cassette was developed, utilizing regulatory elements from the human cytokерatin 18 (K18) gene, including 5' genomic sequences and one of its introns. This cassette efficiently expresses reporter genes, as well as the human cystic fibrosis transmembrane conductance regulator (CFTR) gene, in cultured lung epithelial cells (49).

#### *Alternative splicing:*

Alternative splicing of pre mRNA is a powerful and versatile regulatory mechanism that can effect quantitative control of gene expression and functional diversification of proteins. It contributes to major developmental decisions and also to a fine-tuning of gene function. Genetic and biochemical approaches have identified cis-acting regulatory elements and trans-acting factors that control alternative splicing of specific mRNAs. This mechanism results in the generation of variant isoforms of various proteins from a single gene. These include cell surface molecules such as CD44, receptors, cytokines such as VEGF and enzymes. Products of alternatively spliced transcripts differ in their expression pattern, substrate specificity and other biological parameters.

The FGF receptor RNA undergoes alternative splicing which results in the production of several isoforms, which exhibit different ligand binding specificities. The alternative splicing is regulated in a cell specific manner (53).

5        Alternative spliced mRNAs are often correlated with malignancy. An increase in specific splice variant of tyrosinase was identified in murine melanomas (54). Multiple splicing variants of estrogen receptor are present in individual human breast tumors. CD44 has various isoform, some are characteristic of malignant tissues.

10       Identification of tumor specific alternative splice variants provide new tool for cancer diagnostics. CD44 variants have been used for detection of malignancy in urine samples from patients with urothelial cancer by competitive RT-PCR (55). CD44 exon 6 was suggested as prognostic indicator of metastasis in breast cancer (56).

15       Different enzymes or polypeptides generated by alternative splicing may have different function or catalytic specificity. The identification and characterization of the enzyme forms, which are involved in pathological processes, is crucial for the design of appropriate and efficient drugs.

***Modulation of gene expression – Antisense technology:***

20       An antisense oligonucleotide (e.g., antisense oligodeoxyribonucleotide) may bind its target nucleic acid either by Watson-Crick base pairing or Hoogsteen and anti-Hoogsteen base pairing

(64). According to the Watson-Crick base pairing, heterocyclic bases of the antisense oligonucleotide form hydrogen bonds with the heterocyclic bases of target single-stranded nucleic acids (RNA or single-stranded DNA), whereas according to the Hoogsteen base pairing, the heterocyclic bases of the target nucleic acid are double-stranded DNA, wherein a third strand is accommodated in the major groove of the B-form DNA duplex by Hoogsteen and anti-Hoogsteen base pairing to form a triple helix structure.

According to both the Watson-Crick and the Hoogsteen base pairing models, antisense oligonucleotides have the potential to regulate gene expression and to disrupt the essential functions of the nucleic acids in cells. Therefore, antisense oligonucleotides have possible uses in modulating a wide range of diseases in which gene expression is altered.

Since the development of effective methods for chemically synthesizing oligonucleotides, these molecules have been extensively used in biochemistry and biological research and have the potential use in medicine, since carefully devised oligonucleotides can be used to control gene expression by regulating levels of transcription, transcripts and/or translation.

Oligodeoxyribonucleotides as long as 100 base pairs (bp) are routinely synthesized by solid phase methods using commercially available, fully automated synthesis machines. The chemical synthesis of oligoribonucleotides, however, is far less routine. Oligoribonucleotides are



also much less stable than oligodeoxyribonucleotides, a fact which has contributed to the more prevalent use of oligodeoxyribonucleotides in medical and biological research, directed at, for example, the regulation of transcription or translation levels.

5        Gene expression involves few distinct and well regulated steps. The first major step of gene expression involves transcription of a messenger RNA (mRNA) which is an RNA sequence complementary to the antisense (i.e., -) DNA strand, or, in other words, identical in sequence to the DNA sense (i.e., +) strand, composing the gene. In eukaryotes, transcription  
10 occurs in the cell nucleus.

The second major step of gene expression involves translation of a protein (e.g., enzymes, structural proteins, secreted proteins, gene expression factors, etc.) in which the mRNA interacts with ribosomal RNA complexes (ribosomes) and amino acid activated transfer RNAs (tRNAs) to  
15 direct the synthesis of the protein coded for by the mRNA sequence.

Initiation of transcription requires specific recognition of a promoter DNA sequence located upstream to the coding sequence of a gene by an RNA-synthesizing enzyme -- RNA polymerase. This recognition is preceded by sequence-specific binding of one or more transcription factors  
20 to the promoter sequence. Additional proteins which bind at or close to the promoter sequence may trans upregulate transcription via cis elements known as enhancer sequences. Other proteins which bind to or close to the

promoter, but whose binding prohibits the action of RNA polymerase, are known as repressors.

There are also evidence that in some cases gene expression is downregulated by endogenous antisense RNA repressors that bind a complementary mRNA transcript and thereby prevent its translation into a functional protein.

Thus, gene expression is typically upregulated by transcription factors and enhancers and downregulated by repressors.

However, in many disease situation gene expression is impaired. In many cases, such as different types of cancer, for various reasons the expression of a specific endogenous or exogenous (e.g., of a pathogen such as a virus) gene is upregulated. Furthermore, in infectious diseases caused by pathogens such as parasites, bacteria or viruses, the disease progression depends on expression of the pathogen genes, this phenomenon may also be considered as far as the patient is concerned as upregulation of exogenous genes.

Most conventional drugs function by interaction with and modulation of one or more targeted endogenous or exogenous proteins, e.g., enzymes. Such drugs, however, typically are not specific for targeted proteins but interact with other proteins as well. Thus, a relatively large dose of drug must be used to effectively modulate a targeted protein.

Typical daily doses of drugs are from  $10^{-5}$  -  $10^{-1}$  millimoles per kilogram of body weight or  $10^{-3}$  - 10 millimoles for a 100 kilogram person.

If this modulation instead could be effected by interaction with and inactivation of mRNA, a dramatic reduction in the necessary amount of drug could likely be achieved, along with a corresponding reduction in side effects. Further reductions could be effected if such interaction could be rendered site-specific. Given that a functioning gene continually produces mRNA, it would thus be even more advantageous if gene transcription could be arrested in its entirety.

Given these facts, it would be advantageous if gene expression could be arrested or downmodulated at the transcription level.

The ability of chemically synthesizing oligonucleotides and analogs thereof having a selected predetermined sequence offers means for downmodulating gene expression. Three types of gene expression modulation strategies may be considered.

At the transcription level, antisense or sense oligonucleotides or analogs that bind to the genomic DNA by strand displacement or the formation of a triple helix, may prevent transcription (64).

At the transcript level, antisense oligonucleotides or analogs that bind target mRNA molecules lead to the enzymatic cleavage of the hybrid by intracellular RNase H (65). In this case, by hybridizing to the targeted mRNA, the oligonucleotides or oligonucleotide analogs provide a duplex

hybrid recognized and destroyed by the RNase H enzyme. Alternatively, such hybrid formation may lead to interference with correct splicing (66). As a result, in both cases, the number of the target mRNA intact transcripts ready for translation is reduced or eliminated.

5        At the translation level, antisense oligonucleotides or analogs that bind target mRNA molecules prevent, by steric hindrance, binding of essential translation factors (ribosomes), to the target mRNA, a phenomenon known in the art as hybridization arrest, disabling the translation of such mRNAs (67).

10       Thus, antisense sequences, which as described hereinabove may arrest the expression of any endogenous and/or exogenous gene depending on their specific sequence, attracted much attention by scientists and pharmacologists who were devoted at developing the antisense approach into a new pharmacological tool (68).

15       For example, several antisense oligonucleotides have been shown to arrest hematopoietic cell proliferation (69), growth (70), entry into the S phase of the cell cycle (71), reduced survival (72) and prevent receptor mediated responses (73). For use of antisense oligonucleotides as antiviral agents the reader is referred to reference 74.

20       For efficient *in vivo* inhibition of gene expression using antisense oligonucleotides or analogs, the oligonucleotides or analogs must fulfill the following requirements (i) sufficient specificity in binding to the target

sequence; (ii) solubility in water; (iii) stability against intra- and extracellular nucleases; (iv) capability of penetration through the cell membrane; and (v) when used to treat an organism, low toxicity.

Unmodified oligonucleotides are impractical for use as antisense sequences since they have short *in vivo* half-lives, during which they are degraded rapidly by nucleases. Furthermore, they are difficult to prepare in more than milligram quantities. In addition, such oligonucleotides are poor cell membrane penetrators (75).

Thus it is apparent that in order to meet all the above listed requirements, oligonucleotide analogs need to be devised in a suitable manner. Therefore, an extensive search for modified oligonucleotides has been initiated.

For example, problems arising in connection with double-stranded DNA (dsDNA) recognition through triple helix formation have been diminished by a clever "switch back" chemical linking, whereby a sequence of polypurine on one strand is recognized, and by "switching back", a homopurine sequence on the other strand can be recognized. Also, good helix formation has been obtained by using artificial bases, thereby improving binding conditions with regard to ionic strength and pH.

In addition, in order to improve half-life as well as membrane penetration, a large number of variations in polynucleotide backbones have been done, nevertheless with little success.

Oligonucleotides can be modified either in the base, the sugar or the phosphate moiety. These modifications include, for example, the use of methylphosphonates, monothiophosphates, dithiophosphates, phosphoramidates, phosphate esters, bridged phosphorothioates, bridged phosphoramidates, bridged methylenephosphonates, dephospho internucleotide analogs with siloxane bridges, carbonate bridges, carboxymethyl ester bridges, carbonate bridges, carboxymethyl ester bridges, acetamide bridges, carbamate bridges, thioether bridges, sulfoxy bridges, sulfono bridges, various "plastic" DNAs, o-anomeric bridges and borane derivatives. For further details the reader is referred to reference 76.

International patent application WO 89/12060 discloses various building blocks for synthesizing oligonucleotide analogs, as well as oligonucleotide analogs formed by joining such building blocks in a defined sequence. The building blocks may be either "rigid" (i.e., containing a ring structure) or "flexible" (i.e., lacking a ring structure). In both cases, the building blocks contain a hydroxy group and a mercapto group, through which the building blocks are said to join to form oligonucleotide analogs. The linking moiety in the oligonucleotide analogs is selected from the group consisting of sulfide (-S-), sulfoxide (-SO-), and sulfone (-SO<sub>2</sub>-). However, the application provides no data supporting the specific binding of an oligonucleotide analog to a target oligonucleotide.

International patent application WO 92/20702 describe an acyclic oligonucleotide which includes a peptide backbone on which any selected chemical nucleobases or analogs are stringed and serve as coding characters as they do in natural DNA or RNA. These new compounds, known as peptide nucleic acids (PNAs), are not only more stable in cells than their natural counterparts, but also bind natural DNA and RNA 50 to 100 times more tightly than the natural nucleic acids cling to each other (77). PNA oligomers can be synthesized from the four protected monomers containing thymine, cytosine, adenine and guanine by Merrifield solid-phase peptide synthesis. In order to increase solubility in water and to prevent aggregation, a lysine amide group is placed at the C-terminal.

Thus, antisense technology requires pairing of messenger RNA with an oligonucleotide to form a double helix that inhibits translation. The concept of antisense-mediated gene therapy was already introduced in 1978 for cancer therapy. This approach was based on certain genes that are crucial in cell division and growth of cancer cells. Synthetic fragments of genetic substance DNA can achieve this goal. Such molecules bind to the targeted gene molecules in RNA of tumor cells, thereby inhibiting the translation of the genes and resulting in dysfunctional growth of these cells. Other mechanisms has also been proposed. These strategies have been used, with some success in treatment of cancers, as well as other illnesses, including viral and other infectious diseases. Antisense oligonucleotides

are typically synthesized in lengths of 13-30 nucleotides. The life span of oligonucleotide molecules in blood is rather short. Thus, they have to be chemically modified to prevent destruction by ubiquitous nucleases present in the body. Phosphorothioates are very widely used modification in antisense oligonucleotide ongoing clinical trials (57). A new generation of antisense molecules consist of hybrid antisense oligonucleotide with a central portion of synthetic DNA while four bases on each end have been modified with 2'O-methyl ribose to resemble RNA. In preclinical studies in laboratory animals, such compounds have demonstrated greater stability to metabolism in body tissues and an improved safety profile when compared with the first-generation unmodified phosphorothioate (Hybridon Inc. news). Dosens of other nucleotide analogs have also been tested in antisense technology.

RNA oligonucleotides may also be used for antisense inhibition as they form a stable RNA-RNA duplex with the target, suggesting efficient inhibition. However, due to their low stability RNA oligonucleotides are typically expressed inside the cells using vectors designed for this purpose. This approach is favored when attempting to target a mRNA that encodes an abundant and long-lived protein (57).

Recent scientific publications have validated the efficacy of antisense compounds in animal models of hepatitis, cancers, coronary artery restenosis and other diseases. The first antisense drug was recently



approved by the FDA. This drug Fomivirsen, developed by Isis, is indicated for local treatment of cytomegalovirus in patients with AIDS who are intolerant of or have a contraindication to other treatments for CMV retinitis or who were insufficiently responsive to previous treatments for CMV retinitis (Pharmacotherapy News Network).

Several antisense compounds are now in clinical trials in the United States. These include locally administered antivirals, systemic cancer therapeutics. Antisense therapeutics has the potential to treat many life-threatening diseases with a number of advantages over traditional drugs.

Traditional drugs intervene after a disease-causing protein is formed. Antisense therapeutics, however, block mRNA transcription/translation and intervene before a protein is formed, and since antisense therapeutics target only one specific mRNA, they should be more effective with fewer side effects than current protein-inhibiting therapy.

A second option for disrupting gene expression at the level of transcription uses synthetic oligonucleotides capable of hybridizing with double stranded DNA. A triple helix is formed. Such oligonucleotides may prevent binding of transcription factors to the gene's promoter and therefore inhibit transcription. Alternatively, they may prevent duplex unwinding and, therefore, transcription of genes within the triple helical structure.

Another approach is the use of specific nucleic acid sequences to act as decoys for transcription factors. Since transcription factors bind specific

DNA sequences it is possible to synthesize oligonucleotides that will effectively compete with the native DNA sequences for available transcription factors *in vivo*. This approach requires the identification of gene specific transcription factor (57).

5 Indirect inhibition of gene expression was demonstrated for matrix metalloproteinase genes (MMP-1, -3, and -9), which are associated with invasive potential of human cancer cells. E1AF is a transcription activator of MMP genes. Expression of E1AF antisense RNA in HSC3AS cells showed decrease in mRNA and protein levels of MMP-1, -3, and -9. 10 Moreover, HSC3AS showed lower invasive potential *in vitro* and *in vivo*. These results imply that transfection of antisense inhibits tumor invasion by down-regulating MMP genes (58).

### ***Ribozymes:***

Ribozymes are being increasingly used for the sequence-specific 15 inhibition of gene expression by the cleavage of mRNAs encoding proteins of interest. The possibility of designing ribozymes to cleave any specific target RNA has rendered them valuable tools in both basic research and therapeutic applications. In the therapeutics area, ribozymes have been exploited to target viral RNAs in infectious diseases, dominant oncogenes 20 in cancers and specific somatic mutations in genetic disorders. Most notably, several ribozyme gene therapy protocols for HIV patients are already in Phase 1 trials (62). More recently, ribozymes have been used for

transgenic animal research, gene target validation and pathway elucidation.

Several ribozymes are in various stages of clinical trials. ANGIOZYME was the first chemically synthesized ribozyme to be studied in human clinical trials. ANGIOZYME specifically inhibits formation of the VEGF-r  
5 (Vascular Endothelial Growth Factor receptor), a key component in the angiogenesis pathway. Ribozyme Pharmaceuticals, Inc., as well as other firms have demonstrated the importance of anti-angiogenesis therapeutics in animal models. HEPTAZYME, a ribozyme designed to selectively destroy Hepatitis C Virus (HCV) RNA, was found effective in decreasing Hepatitis  
10 C viral RNA in cell culture assays (Ribozyme Pharmaceuticals, Incorporated - WEB home page).

*Gene disruption in animal models:*

The emergence of gene inactivation by homologous recombination methodology in embryonic stem cells has revolutionized the field of mouse  
15 genetics. The availability of a rapidly growing number of mouse null mutants has represented an invaluable source of knowledge on mammalian development, cellular biology and physiology, and has provided many models for human inherited diseases. Animal models are required for an effective drug delivery development program and evaluation of gene  
20 therapy approach. The improvement of the original knockout strategy, as well as exploitation of exogenous enzymatic systems that are active in the recombination process, has been considerably extended the range of genetic

manipulations that can be produced. Additional methods have been developed to provide versatile research tools: Double replacement method, sequential gene targeting, conditional cell type specific gene targeting, single copy integration method, inducible gene targeting, gene disruption by viral delivery, replacing one gene with another, the so called knock-in method and the induction of specific balanced chromosomal translocation. It is now possible to introduce a point mutation as a unique change in the entire genome, therefore allowing very fine dissection of gene function *in vivo*. Furthermore, the advent of methods allowing conditional gene targeting opens the way for analysis of consequence of a particular mutation in a defined organ and at a specific time during the life of the experimental animal (59).

#### ***DNA vaccination:***

Observations in the early 1990s that plasmid DNA could directly transfect animal cells *in vivo* sparked exploration of the use of DNA plasmids to induce immune response by direct injection into animal of DNA encoding antigenic protein. When a DNA vaccine plasmid enters the eukaryotic cell, the protein it encodes is transcribed and translated within the cell. In the case of pathogens, these proteins are presented to the immune system in their native form, mimicking the presentation of antigens during a natural infection. DNA vaccination is particularly useful for the induction of T cell activation. It was applied for viral and bacterial

infectious diseases, as well as for allergy and for cancer. The central hypothesis behind active specific immunotherapy for cancer is that tumor cells express unique antigens that should stimulate the immune system. The first DNA vaccine against tumor was carcino-embryonic antigen (CEA).  
5 DNA vaccinated animals expressed immunoprotection and immunotherapy of human CEA-expressing syngeneic mouse colon and breast carcinoma (61). In a mouse model of neuroblastoma, DNA immunization with HuD resulted in tumor growth inhibition with no neurological disease (60). Immunity to the brown locus protein, gp<sup>75</sup> tyrosinase-related protein-1,  
10 associated with melanoma, was investigated in a syngeneic mouse model. Priming with human gp75 DNA broke tolerance to mouse gp75. Immunity against mouse gp75 provided significant tumor protection (60).

*Glycosyl hydrolases:*

Glycosyl hydrolases are a widespread group of enzymes that  
15 hydrolyze the o-glycosidic bond between two or more carbohydrates or between a carbohydrate and a noncarbohydrate moiety. The enzymatic hydrolysis of glycosidic bond occurs by using major one or two mechanisms leading to overall retention or inversion of the anomeric configuration. In both mechanisms catalysis involves two residues: a proton donor and a  
20 nucleophile. Glycosyl hydrolyses have been classified into 58 families based on amino acid similarities. The glycosyl hydrolyses from families 1, 2, 5, 10, 17, 30, 35, 39 and 42 act on a large variety of substrates, however,

they all hydrolyze the glycosidic bond in a general acid catalysis mechanism, with retention of the anomeric configuration. The mechanism involves two glutamic acid residues, which are the proton donors and the nucleophile, with an asparagine always preceding the proton donor.

Analyses of a set of known 3D structures from this group revealed that their catalytic domains, despite the low level of sequence identity, adopt a similar ( $\alpha/\beta$ ) 8 fold with the proton donor and the nucleophile located at the C-terminal ends of strands  $\beta 4$  and  $\beta 7$ , respectively. Mutations in the functional conserved amino acids of lysosomal glycosyl hydrolases were identified in lysosomal storage diseases.

Lysosomal glycosyl hydrolases including  $\beta$ -glucuronidase,  $\beta$ -mannosidase,  $\beta$ -glucocerebrosidase,  $\beta$ -galactosidase and  $\alpha$ -L iduronidase, are all exo-glycosyl hydrolases, belong to the GH-A clan and share a similar catalytic site. However, many endo-glucanases from various organisms, such as bacterial and fungal xylanases and cellulases share this catalytic domain.

#### ***Genomic sequence of hpa gene and its implications:***

It is well established that heparanase activity is correlated with cancer metastasis. This correlation was demonstrated at the level of enzymatic activity as well as the levels of protein and *hpa* cDNA expression in highly metastatic cancer cells as compared with non-metastatic cells. As such, inhibition of heparanase activity is desirable, and has been attempted

by several means. The genomic region, encoding the *hpa* gene and the surrounding, provides a new powerful tool for regulation of heparanase activity at the level of gene expression. Regulatory sequences may reside in noncoding regions both upstream and downstream the transcribed region as well as in intron sequences. A DNA sequence upstream of the transcription start site contains the promoter region and potential regulatory elements. Regulatory factors, which interact with the promoter region may be identified and be used as potential drugs for inhibition of cancer, metastasis and inflammation. The promoter region can be used to screen for inhibitors of heparanase gene expression. Furthermore, the *hpa* promoter can be used to direct cell specific, particularly cancer cell specific, expression of foreign genes, such as cytotoxic or apoptotic genes, in order to specifically destroy cancer cells.

Cancer and yet unknown related genetic disorders may involve rearrangements and mutations in the heparanase gene, either in coding or non-coding regions. Such mutations may affect expression level or enzymatic activity. The genomic sequence of *hpa* enables the amplification of specific genomic DNA fragments, identification and diagnosis of mutations.

There is thus a widely recognized need for, and it would be highly advantageous to have genomic, cDNA and composite polynucleotides encoding a polypeptide having heparanase activity, vectors including same,

genetically modified cells expressing heparanase and a recombinant protein having heparanase activity, as well as antisense oligonucleotides, constructs and ribozymes which can be used for down regulation heparanase activity.

## 5 SUMMARY OF THE INVENTION

Cloning of the human *hpa* gene which encodes heparanase, and expression of recombinant heparanase by transfected host cells is reported herein, as well as downregulation of heparanase activity by antisense technology.

10 A purified preparation of heparanase isolated from human hepatoma cells was subjected to tryptic digestion and microsequencing. The YGPDVGQPR (SEQ ID NO:8) sequence revealed was used to screen EST databases for homology to the corresponding back translated DNA sequence. Two closely related EST sequences were identified and were  
15 thereafter found to be identical. Both clones contained an insert of 1020 bp which included an open reading frame of 973 bp followed by a 27 bp of 3' untranslated region and a Poly A tail. Translation start site was not identified.

Cloning of the missing 5' end of *hpa* was performed by PCR  
20 amplification of DNA from placenta Marathon RACE cDNA composite using primers selected according to the EST clones sequence and the linkers of the composite. A 900 bp PCR fragment, partially overlapping with the



identified 3' encoding EST clones was obtained. The joined cDNA fragment (*hpa*), 1721 bp long (SEQ ID NO:9), contained an open reading frame which encodes a polypeptide of 543 amino acids (SEQ ID NO:10) with a calculated molecular weight of 61,192 daltons.

5 Cloning an extended 5' sequence was enabled from the human SK-hep1 cell line by PCR amplification using the Marathon RACE. The 5' extended sequence of the SK-hep1 *hpa* cDNA was assembled with the sequence of the *hpa* cDNA isolated from human placenta (SEQ ID NO:9). The assembled sequence contained an open reading frame, SEQ ID NOs: 13  
10 and 15, which encodes, as shown in SEQ ID NOs:14 and 15, a polypeptide of 592 amino acids with a calculated molecular weight of 66,407 daltons.

The ability of the *hpa* gene product to catalyze degradation of heparan sulfate in an *in vitro* assay was examined by expressing the entire open reading frame of *hpa* in insect cells, using the Baculovirus expression  
15 system. Extracts and conditioned media of cells infected with virus containing the *hpa* gene, demonstrated a high level of heparan sulfate degradation activity both towards soluble ECM-derived HSPG and intact ECM. This degradation activity was inhibited by heparin, which is another substrate of heparanase. Cells infected with a similar construct containing  
20 no *hpa* gene had no such activity, nor did non-infected cells. The ability of heparanase expressed from the extended 5' clone towards heparin was demonstrated in a mammalian expression system.

The expression pattern of *hpa* RNA in various tissues and cell lines was investigated using RT-PCR. It was found to be expressed only in tissues and cells previously known to have heparanase activity.

A panel of monochromosomal human/CHO and human/mouse  
5 somatic cell hybrids was used to localize the human heparanase gene to human chromosome 4. The newly isolated heparanase sequence can be used to identify a chromosome region harboring a human heparanase gene in a chromosome spread.

A human genomic library was screened and the human locus  
10 harboring the heparanase gene isolated, sequenced and characterized. Alternatively spliced heparanase mRNAs were identified and characterized. The human heparanase promoter has been isolated, identified and positively tested for activity. The mouse heparanase promoter has been isolated and identified as well. Antisense heparanase constructs were prepared and their  
15 influence on cells *in vitro* tested. A predicted heparanase active site was identified. And finally, the presence of sequences hybridizing with human heparanase sequences was demonstrated for a variety of mammals and for an avian.

According to one aspect of the present invention there is provided an  
20 isolated nucleic acid comprising a genomic, complementary or composite polynucleotide sequence encoding a polypeptide having heparanase catalytic activity.

According to further features in preferred embodiments of the invention described below, the polynucleotide or a portion thereof is hybridizable with SEQ ID NOs: 9, 13, 42, 43 or a portion thereof at 68 °C in 6 x SSC, 1 % SDS, 5 x Denharts, 10 % dextran sulfate, 100 µg/ml salmon sperm DNA, and <sup>32</sup>p labeled probe and wash at 68 °C with 3 x SSC and 0.1 % SDS.

According to still further features in the described preferred embodiments the polynucleotide or a portion thereof is at least 60 % identical with SEQ ID NOs: 9, 13, 42, 43 or portions thereof as determined using the Bestfit procedure of the DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the university of Wisconsin (gap creation penalty - 12, gap extension penalty - 4).

According to still further features in the described preferred embodiments the polypeptide is as set forth in SEQ ID NOs: 10, 14, 44 or portions thereof.

According to still further features in the described preferred embodiments the polypeptide is at least 60 % homologous to SEQ ID NOs: 10, 14, 44 or portions thereof as determined with the Smith-Waterman algorithm, using the Bioaccelerator platform developed by Compugene (gapop: 10.0, gapext: 0.5, matrix: blosum62).

According to additional aspects of the present invention there are provided a nucleic acid construct (vector) comprising the isolated nucleic acid described herein and a host cell comprising the construct.

According to a further aspect of the present invention there is  
5 provided an antisense oligonucleotide comprising a polynucleotide or a polynucleotide analog of at least 10 bases being hybridizable *in vivo*, under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity.

According to an additional aspect of the present invention there is  
10 provided a method of *in vivo* downregulating heparanase activity comprising the step of *in vivo* administering the antisense oligonucleotide herein described.

According to yet an additional aspect of the present invention there is provided a pharmaceutical composition comprising the antisense  
15 oligonucleotide herein described and a pharmaceutically acceptable carrier.

According to still an additional aspect of the present invention there is provided a ribozyme comprising the antisense oligonucleotide described herein and a ribozyme sequence.

According to a further aspect of the present invention there is  
20 provided an antisense nucleic acid construct comprising a promoter sequence and a polynucleotide sequence directing the synthesis of an antisense RNA sequence of at least 10 bases being hybridizable *in vivo*,

under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity.

According to further features in preferred embodiments of the invention described below, the polynucleotide strand encoding the polypeptide having heparanase catalytic activity is as set forth in SEQ ID  
5 NOs: 9, 13, 42 or 43.

According to still further features in the described preferred embodiments the polypeptide having heparanase catalytic activity is as set forth in SEQ ID NOs: 10, 14 or 44.

10 According to still a further aspect of the present invention there is provided a method of *in vivo* downregulating heparanase activity comprising the step of *in vivo* administering the antisense nucleic acid construct herein described.

According to yet a further aspect of the present invention there is  
15 provided a pharmaceutical composition comprising the antisense nucleic acid construct herein described and a pharmaceutically acceptable carrier.

According to a further aspect of the present invention there is provided a nucleic acid construct comprising a polynucleotide sequence functioning as a promoter, the polynucleotide sequence is derived from SEQ  
20 ID NO:42 and includes at least nucleotides 2535-2635 thereof or from SEQ ID NO:43 and includes at least nucleotides 320-420.

According to a further aspect of the present invention there is provided a method of expressing a polynucleotide sequence comprising the step of ligating the polynucleotide sequence into the nucleic acid construct described above, downstream of the polynucleotide sequence derived from  
5 SEQ ID NOs:42 or 43.

According to a further aspect of the present invention there is provided a recombinant protein comprising a polypeptide having heparanase catalytic activity.

According to further features in preferred embodiments of the  
10 invention described below, the polypeptide includes at least a portion of SEQ ID NOs:10, 14 or 44.

According to still further features in the described preferred embodiments the protein is encoded by a polynucleotide hybridizable with SEQ ID NOs: 9, 13, 42, 43 or a portion thereof at 68 °C in 6 x SSC, 1 %  
15 SDS, 5 x Denharts, 10 % dextran sulfate, 100 µg/ml salmon sperm DNA, and <sup>32</sup>p labeled probe and wash at 68 °C with 3 x SSC and 0.1 % SDS.

According to still further features in the described preferred embodiments the protein is encoded by a polynucleotide at least 60 % identical with SEQ ID NOs: 9, 13, 42, 43 or portions thereof as determined  
20 using the Bestfit procedure of the DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the university of Wisconsin (gap creation penalty - 12, gap extension penalty - 4).

According to a further aspect of the present invention there is provided a pharmaceutical composition comprising, as an active ingredient, the recombinant protein herein described.

According to a further aspect of the present invention there is  
5 provided a method of identifying a chromosome region harboring a heparanase gene in a chromosome spread comprising the steps of (a) hybridizing the chromosome spread with a tagged polynucleotide probe encoding heparanase; (b) washing the chromosome spread, thereby removing excess of non-hybridized probe; and (c) searching for signals  
10 associated with the hybridized tagged polynucleotide probe, wherein detected signals being indicative of a chromosome region harboring a heparanase gene.

According to a further aspect of the present invention there is provided a method of *in vivo* eliciting anti-heparanase antibodies  
15 comprising the steps of administering a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*. Accordingly, there is provided also a DNA vaccine for *in vivo* eliciting anti-heparanase antibodies comprising a nucleic  
20 acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*.

The present invention can be used to develop new drugs to inhibit tumor cell metastasis, inflammation and autoimmunity. The identification of the *hpa* gene encoding for heparanase enzyme enables the production of a recombinant enzyme in heterologous expression systems. Additional features, advantages, uses and applications of the present invention in biological science and in diagnostic and therapeutic medicine are described hereinafter.

#### BRIEF DESCRIPTION OF THE DRAWINGS

The invention herein described, by way of example only, with reference to the accompanying drawings, wherein:

FIG. 1 presents nucleotide sequence and deduced amino acid sequence of *hpa* cDNA. A single nucleotide difference at position 799 (A to T) between the EST (Expressed Sequence Tag) and the PCR amplified cDNA (reverse transcribed RNA) and the resulting amino acid substitution (Tyr to Phe) are indicated above and below the substituted unit, respectively. Cysteine residues and the poly adenylation consensus sequence are underlined. The asterisk denotes the stop codon TGA.

FIG. 2 demonstrates degradation of soluble sulfate labeled HSPG substrate by lysates of High Five cells infected with pF*hpa2* virus. Lysates of High Five cells that were infected with pF*hpa2* virus (●) or control pF2 virus (□) were incubated (18 h, 37 °C) with sulfate labeled ECM-derived



soluble HSPG (peak I). The incubation medium was then subjected to gel filtration on Sepharose 6B. Low molecular weight HS degradation fragments (peak II) were produced only during incubation with the pFhpa2 infected cells, but there was no degradation of the HSPG substrate (✧) by lysates of pF2 infected cells.

FIGs. 3a-b demonstrate degradation of soluble sulfate labeled HSPG substrate by the culture medium of pFhpa2 and pFhpa4 infected cells. Culture media of High Five cells infected with pFhpa2 (3a) or pFhpa4 (3b) viruses (●), or with control viruses (□) were incubated (18 h, 37 °C) with sulfate labeled ECM-derived soluble HSPG (peak I, ✧). The incubation media were then subjected to gel filtration on Sepharose 6B. Low molecular weight HS degradation fragments (peak II) were produced only during incubation with the *hpa* gene containing viruses. There was no degradation of the HSPG substrate by the culture medium of cells infected with control viruses.

FIG. 4 presents size fractionation of heparanase activity expressed by pFhpa2 infected cells. Culture medium of pFhpa2 infected High Five cells was applied onto a 50 kDa cut-off membrane. Heparanase activity (conversion of the peak I substrate, ✧ into peak II HS degradation fragments) was found in the high (> 50 kDa) (●), but not low (< 50 kDa) (○) molecular weight compartment.

FIGs. 5a-b demonstrate the effect of heparin on heparanase activity expressed by pFhpa2 and pFhpa4 infected High Five cells. Culture media of pFhpa2 (5a) and pFhpa4 (5b) infected High Five cells were incubated (18 h, 37 °C) with sulfate labeled ECM-derived soluble HSPG (peak I,  $\diamond$ ) in the absence ( $\circ$ ) or presence ( $\Delta$ ) of 10  $\mu$ g/ml heparin. Production of low molecular weight HS degradation fragments was completely abolished in the presence of heparin, a potent inhibitor of heparanase activity (6, 7).

FIGs. 6a-b demonstrate degradation of sulfate labeled intact ECM by virus infected High Five and Sf21 cells. High Five (6a) and Sf21 (6b) cells were plated on sulfate labeled ECM and infected (48 h, 28 °C) with pFhpa4 ( $\circ$ ) or control pF1 ( $\square$ ) viruses. Control non-infected Sf21 cells ( $\text{R}$ ) were plated on the labeled ECM as well. The pH of the cultured medium was adjusted to 6.0 - 6.2 followed by 24 h incubation at 37 °C. Sulfate labeled material released into the incubation medium was analyzed by gel filtration on Sepharose 6B. HS degradation fragments were produced only by cells infected with the *hpa* containing virus.

FIG. 7a-b demonstrate degradation of sulfate labeled intact ECM by virus infected cells. High Five (7a) and Sf21 (7b) cells were plated on sulfate labeled ECM and infected (48 h, 28 °C) with pFhpa4 ( $\circ$ ) or control pF1 ( $\square$ ) viruses. Control non-infected Sf21 cells ( $\text{R}$ ) were plate on labeled ECM as well. The pH of the cultured medium was adjusted to 6.0 - 6.2, followed by 48 h incubation at 28 °C. Sulfate labeled degradation

fragments released into the incubation medium was analyzed by gel filtration on Sepharose 6B. HS degradation fragments were produced only by cells infected with the *hpa* containing virus.

FIGs. 8a-b demonstrate degradation of sulfate labeled intact ECM by the culture medium of pF*hpa4* infected cells. Culture media of High Five (8a) and Sf21 (8b) cells that were infected with pF*hpa4* (●) or control pF1 (□) viruses were incubated (48 h, 37 °C, pH 6.0) with intact sulfate labeled ECM. The ECM was also incubated with the culture medium of control non-infected Sf21 cells (○). Sulfate labeled material released into the reaction mixture was subjected to gel filtration analysis. Heparanase activity was detected only in the culture medium of pF*hpa4* infected cells.

FIGs. 9a-b demonstrate the effect of heparin on heparanase activity in the culture medium of pF*hpa4* infected cells. Sulfate labeled ECM was incubated (24 h, 37 °C, pH 6.0) with culture medium of pF*hpa4* infected High Five (9a) and Sf21 (9b) cells in the absence (●) or presence (V) of 10 µg/ml heparin. Sulfate labeled material released into the incubation medium was subjected to gel filtration on Sepharose 6B. Heparanase activity (production of peak II HS degradation fragments) was completely inhibited in the presence of heparin.

FIGs. 10a-b demonstrate purification of recombinant heparanase on heparin-Sepharose. Culture medium of Sf21 cells infected with pF*hpa4* virus was subjected to heparin-Sepharose chromatography. Elution of

fractions was performed with 0.35 - 2 M NaCl gradient (◊). Heparanase activity in the eluted fractions is demonstrated in Figure 10a (○). Fractions 15-28 were subjected to 15 % SDS-polyacrylamide gel electrophoresis followed by silver nitrate staining. A correlation is demonstrated between a major protein band (MW ~ 63,000) in fractions 19 - 24 and heparanase activity.

FIGs. 11a-b demonstrate purification of recombinant heparanase on a Superdex 75 gel filtration column. Active fractions eluted from heparin-Sepharose (Figure 10a) were pooled, concentrated and applied onto Superdex 75 FPLC column. Fractions were collected and aliquots of each fraction were tested for heparanase activity (C, Figure 11a) and analyzed by SDS-polyacrylamide gel electrophoresis followed by silver nitrate staining (Figure 11b). A correlation is seen between the appearance of a major protein band (MW ~ 63,000) in fractions 4 - 7 and heparanase activity.

FIGs. 12a-e demonstrate expression of the *hpa* gene by RT-PCR with total RNA from human embryonal tissues (12a), human extra-embryonal tissues (12b) and cell lines from different origins (12c-e). RT-PCR products using *hpa* specific primers (I), primers for GAPDH housekeeping gene (II), and control reactions without reverse transcriptase demonstrating absence of genomic DNA or other contamination in RNA samples (III). M- DNA molecular weight marker VI (Boehringer Mannheim). For 12a: lane 1 - neutrophil cells (adult), lane 2 - muscle, lane 3 - thymus, lane 4 - heart, lane

5 - adrenal. For 12b: lane 1 - kidney, lane 2 - placenta (8 weeks), lane 3 - placenta (11 weeks), lanes 4-7 - mole (complete hydatidiform mole), lane 8 - cytotrophoblast cells (freshly isolated), lane 9 - cytotrophoblast cells (1.5 h *in vitro*), lane 10 - cytotrophoblast cells (6 h *in vitro*), lane 11 - cytotrophoblast cells (18 h *in vitro*), lane 12 - cytotrophoblast cells (48 h *in vitro*). For 12c: lane 1 - JAR bladder cell line, lane 2 - NCITT testicular tumor cell line, lane 3 - SW-480 human hepatoma cell line, lane 4 - HTR (cytotrophoblasts transformed by SV40), lane 5 - HPTLP-I hepatocellular carcinoma cell line, lane 6 - EJ-28 bladder carcinoma cell line. For 12d: lane 1 - SK-hep-1 human hepatoma cell line, lane 2 - DAMI human megakaryocytic cell line, lane 3 - DAMI cell line + PMA, lane 4 - CHRF cell line + PMA, lane 5 - CHRF cell line. For 12e: lane 1 - ABAE bovine aortic endothelial cells, lane 2 - 1063 human ovarian cell line, lane 3 - human breast carcinoma MDA435 cell line, lane 4 - human breast carcinoma MDA231 cell line.

FIG. 13 presents a comparison between nucleotide sequences of the human *hpa* and a mouse EST cDNA fragment (SEQ ID NO:12) which is 80 % homologous to the 3' end (starting at nucleotide 1066 of SEQ ID NO:9) of the human *hpa*. The aligned termination codons are underlined.

FIG. 14 demonstrates the chromosomal localization of the *hpa* gene. PCR products of DNA derived from somatic cell hybrids and of genomic DNA of hamster, mouse and human of were separated on 0.7 % agarose gel

following amplification with *hpa* specific primers. Lane 1 – Lambda DNA digested with *Bst*EII, lane 2 – no DNA control, lanes 3 – 29, PCR amplification products. Lanes 3-5 – human, mouse and hamster genomic DNA, respectively. Lanes 6-29, human monochromosomal somatic cell hybrids representing chromosomes 1-22 and X and Y, respectively. Lane 30 – Lambda DNA digested with *Bst*EII. An amplification product of approximately 2.8 Kb is observed only in lanes 5 and 9, representing human genomic DNA and DNA derived from cell hybrid carrying human chromosome 4, respectively. These results demonstrate that the *hpa* gene is localized in human chromosome 4.

FIG. 15 demonstrates the genomic exon-intron structure of the human *hpa* locus (top) and the relative positions of the lambda clones used as sequencing templates to sequence the locus (below). The vertical rectangles represent exons (E) and the horizontal lines therebetween represent introns (I), upstream (U) and downstream (D) regions. Continuous lines represent DNA fragments, which were used for sequence analysis. The discontinuous line in lambda 6 represent a region, which overlaps with lambda 8 and hence was not analyzed. The plasmid contains a PCR product, which bridges the gap between L3 and L6.

FIG. 16 presents the nucleotide sequence of the genomic region of the *hpa* gene. Exon sequences appear in upper case and intron sequences in lower case. The deduced amino acid sequence of the exons is printed below

the nucleotide sequence. Two predicted transcription start sites are shown in bold.

FIG. 17 presents an alignment of the amino acid sequences of human heparanase, mouse and partial sequences of rat homologues. The human and the mouse sequences were determined by sequence analysis of the isolated cDNAs. The rat sequence is derived from two different EST clones, which represent two different regions (5' and 3') of the rat *hpa* cDNA. The human sequence and the amino acids in the mouse and rat homologues, which are identical to the human sequence, appear in bold.

FIG. 18 presents a heparanase Zoo blot. Ten micrograms of genomic DNA from various sources were digested with *EcoRI* and separated on 0.7 % agarose – TBE gel. Following electrophoresis, the was gel treated with HCl and than with NaOH and the DNA fragments were downward transferred to a nylon membrane (Hybond N+, Amersham) with 0.4 N NaOH. The membrane was hybridized with a 1.6 Kb DNA probe that contained the entire *hpa* cDNA. Lane order: H – Human; M – Mouse; Rt – Rat; P – Pig; Cw – Cow; Hr – Horse; S – Sheep; Rb – Rabbit; D – Dog; Ch – Chicken; F – Fish. Size markers (Lambda *Bst*II) are shown on the left

FIG. 19 demonstrates the secondary structure prediction for heparanase performed using the PHD server – Profile network Prediction Heidelberg. H – helix, E – extended (beta strand), The glutamic acid

predicted as the proton donor is marked by asterisk and the possible nucleophiles are underlined.

### DESCRIPTION OF THE PREFERRED EMBODIMENTS

5       The present invention is of a polynucleotide or nucleic acid, referred to hereinbelow interchangeably as *hpa*, *hpa* cDNA or *hpa* gene or identified by its SEQ ID NOs, encoding a polypeptide having heparanase activity, vectors or nucleic acid constructs including same and which are used for over-expression or antisense inhibition of heparanase, genetically modified  
10 cells expressing same, recombinant protein having heparanase activity, antisense oligonucleotides and ribozymes for heparanase modulation, and heparanase promoter sequences which can be used to direct the expression of desired genes.

Before explaining at least one embodiment of the invention in detail,  
15 it is to be understood that the invention is not limited in its application to the details of construction and the arrangement of the components set forth in the following description or illustrated in the drawings. The invention is capable of other embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology  
20 employed herein is for the purpose of description and should not be regarded as limiting.



Cloning of the human and mouse *hpa* genes, cDNAs and genomic sequence (for human), encoding heparanase and expressing recombinant heparanase by transfected cells is reported herein. These are the first mammalian heparanase genes to be cloned.

5 A purified preparation of heparanase isolated from human hepatoma cells was subjected to tryptic digestion and microsequencing.

The YGPDVGQPR (SEQ ID NO:8) sequence revealed was used to screen EST databases for homology to the corresponding back translated DNA sequences. Two closely related EST sequences were identified and  
10 were thereafter found to be identical.

Both clones contained an insert of 1020 bp which includes an open reading frame of 973 bp followed by a 3' untranslated region of 27 bp and a Poly A tail, whereas a translation start site was not identified.

Cloning of the missing 5' end was performed by PCR amplification  
15 of DNA from placenta Marathon RACE cDNA composite using primers selected according to the EST clones sequence and the linkers of the composite.

A 900 bp PCR fragment, partially overlapping with the identified 3' encoding EST clones was obtained. The joined cDNA fragment (*hpa*),  
20 1721 bp long (SEQ ID NO:9), contained an open reading frame which encodes, as shown in Figure 1 and SEQ ID NO:11, a polypeptide of 543

amino acids (SEQ ID NO:10) with a calculated molecular weight of 61,192 daltons.

A single nucleotide difference at position 799 (A to T) between the EST clones and the PCR amplified cDNA was observed. This difference results in a single amino acid substitution (Tyr to Phe) (Figure 1). Furthermore, the published EST sequences contained an unidentified nucleotide, which following DNA sequencing of both the EST clones was resolved into two nucleotides (G and C at positions 1630 and 1631 in SEQ ID NO:9, respectively).

The ability of the *hpa* gene product to catalyze degradation of heparan sulfate in an *in vitro* assay was examined by expressing the entire open reading frame in insect cells, using the Baculovirus expression system.

Extracts and conditioned media of cells infected with virus containing the *hpa* gene, demonstrated a high level of heparan sulfate degradation activity both towards soluble ECM-derived HSPG and intact ECM, which was inhibited by heparin, while cells infected with a similar construct containing no *hpa* gene had no such activity, nor did non-infected cells.

The expression pattern of *hpa* RNA in various tissues and cell lines was investigated using RT-PCR. It was found to be expressed only in tissues and cells previously known to have heparanase activity.

Cloning an extended 5' sequence was enabled from the human SK-hep1 cell line by PCR amplification using the Marathon RACE. The 5' extended sequence of the SK-hep1 *hpa* cDNA was assembled with the sequence of the *hpa* cDNA isolated from human placenta (SEQ ID NO:9).

5 The assembled sequence contained an open reading frame, SEQ ID NOs: 13 and 15, which encodes, as shown in SEQ ID NOs:14 and 15, a polypeptide of 592 amino acids, with a calculated molecular weight of 66,407 daltons. This open reading frame was shown to direct the expression of catalytically active heparanase in a mammalian cell expression system. The expressed  
10 heparanase was detectable by anti heparanase antibodies in Western blot analysis.

A panel of monochromosomal human/CHO and human/mouse somatic cell hybrids was used to localize the human heparanase gene to human chromosome 4. The newly isolated heparanase sequence can  
15 therefore be used to identify a chromosome region harboring a human heparanase gene in a chromosome spread.

The *hpa* cDNA was then used as a probe to screen a human genomic library. Several phages were positive. These phages were analyzed and were found to cover most of the *hpa* locus, except for a small  
20 portion which was recovered by bridging PCR. The *hpa* locus covers about 50,000 bp. The *hpa* gene includes 12 exons separated by 11 introns.

RT-PCR performed on a variety of cells revealed alternatively spliced *hpa* transcripts.

The amino acid sequence of human heparanase was used to search for homologous sequences in the DNA and protein databases. Several human EST's were identified, as well as mouse sequences highly homologous to human heparanase. The following mouse EST's were identified AA177901, AA674378, AA67997, AA047943, AA690179, AI122034, all sharing an identical sequence and correspond to amino acids 336-543 of the human heparanase sequence. The entire mouse heparanase cDNA was cloned, based on the nucleotide sequence of the mouse EST's using Marathon cDNA libraries. The mouse and the human *hpa* genes share an average homology of 78 % between the nucleotide sequences and 81 % similarity between the deduced amino acid sequences. *hpa* homologous sequences from rat were also uncovered (EST's AI060284 and AI237828).

Homology search of heparanase amino acid sequence against the DNA and the protein databases and prediction of its protein secondary structure enabled to identify candidate amino acids that participate in the heparanase active site.

Expression of *hpa* antisense in mammalian cell lines resulted in about five fold decrease in the number of recoverable cells as compared to controls.

Human *Hpa* cDNA was shown to hybridize with genomic DNAs of a variety of mammalian species and with an avian.

The human and mouse *hpa* promoters were identified and the human promoter was tested positive in directing the expression of a reporter gene.

5 Thus, according to the present invention there is provided an isolated nucleic acid comprising a genomic, complementary or composite polynucleotide sequence encoding a polypeptide having heparanase catalytic activity.

The phrase "composite polynucleotide sequence" refers to a sequence  
10 which includes exonal sequences required to encode the polypeptide having heparanase activity, as well as any number of intronal sequences. The intronal sequences can be of any source and typically will include conserved splicing signal sequences. Such intronal sequences may further include cis acting expression regulatory elements.

15 The term "heparanase catalytic activity" or its equivalent term "heparanase activity" both refer to a mammalian endoglycosidase hydrolyzing activity which is specific for heparan or heparan sulfate proteoglycan substrates, as opposed to the activity of bacterial enzymes (heparinase I, II and III) which degrade heparin or heparan sulfate by means  
20 of  $\beta$ -elimination (37).

According to a preferred embodiment of the present invention the polynucleotide or a portion thereof is hybridizable with SEQ ID NOs: 9, 13,

42, 43 or a portion thereof at 68 °C in 6 x SSC, 1 % SDS, 5 x Denharts, 10 % dextran sulfate, 100 µg/ml salmon sperm DNA, and <sup>32</sup>p labeled probe and wash at 68 °C with 3, 2, 1, 0.5 or 0.1 x SSC and 0.1 % SDS.

According to another preferred embodiment of the present invention  
5 the polynucleotide or a portion thereof is at least 60 %, preferably at least 65 %, more preferably at least 70 %, more preferably at least 75 %, more preferably at least 80 %, more preferably at least 85 %, more preferably at least 90 %, most preferably, 95-100 % identical with SEQ ID NOs: 9, 13, 42, 43 or portions thereof as determined using the Bestfit procedure of the  
10 DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the university of Wisconsin (gap creation penalty - 12, gap extension penalty - 4 - which are the default parameters).

According to another preferred embodiment of the present invention  
the polypeptide encoded by the polynucleotide sequence is as set forth in  
15 SEQ ID NOs:10, 14, 44 or portions thereof having heparanase catalytic activity. Such portions are expected to include amino acids Asp-Glu 224-225 (SEQ ID NO:10), which can serve as proton donors and glutamic acid 343 or 396 which can serve as a nucleophile.

According to another preferred embodiment of the present invention  
20 the polypeptide encoded by the polynucleotide sequence is at least 60 %, preferably at least 65 %, more preferably at least 70 %, more preferably at least 75 %, more preferably at least 80 %, more preferably at least 85 %, more preferably at least 90 %, most preferably, 95-100 % identical with SEQ ID NOs: 10, 14, 44 or portions thereof as determined using the Bestfit procedure of the

more preferably at least 90 %, most preferably, 95-100 % homologous (both similar and identical acids) to SEQ ID NOs:10, 14, 44 or portions thereof as determined with the Smith-Waterman algorithm, using the Bioaccelerator platform developed by Compugene (gapop: 10.0, gapext: 0.5, matrix: 5 blosum62, see also the description to Figure 17).

Further according to the present invention there is provided a nucleic acid construct comprising the isolated nucleic acid described herein. The construct may and preferably further include an origin of replication and trans regulatory elements, such as promoter and enhancer sequences.

10 The construct or vector can be of any type. It may be a phage which infects bacteria or a virus which infects eukaryotic cells. It may also be a plasmid, phagemid, cosmid, bacmid or an artificial chromosome.

Further according to the present invention there is provided a host cell comprising the nucleic acid construct described herein. The host cell 15 can be of any type. It may be a prokaryotic cell, an eukaryotic cell, a cell line, or a cell as a portion of an organism. The polynucleotide encoding heparanase can be permanently or transiently present in the cell. In other words, genetically modified cells obtained following stable or transient transfection, transformation or transduction are all within the scope of the 20 present invention. The polynucleotide can be present in the cell in low copy (say 1-5 copies) or high copy number (say 5-50 copies or more). It may be

integrated in one or more chromosomes at any location or be present as an extrachromosomal material.

The present invention is further directed at providing a heparanase over-expression system which includes a cell overexpressing heparanase catalytic activity. The cell may be a genetically modified host cell transiently or stably transfected or transformed with any suitable vector which includes a polynucleotide sequence encoding a polypeptide having heparanase activity and a suitable promoter and enhancer sequences to direct over-expression of heparanase. However, the overexpressing cell may also be a product of an insertion (e.g., via homologous recombination) of a promoter and/or enhancer sequence downstream to the endogenous heparanase gene of the expressing cell, which will direct over-expression from the endogenous gene.

The term "over-expression" as used herein in the specification and claims below refers to a level of expression which is higher than a basal level of expression typically characterizing a given cell under otherwise identical conditions.

According to another aspect the present invention provides an antisense oligonucleotide comprising a polynucleotide or a polynucleotide analog of at least 10, preferably 11-15, more preferably 16-17, more preferably 18, more preferably 19-25, more preferably 26-35, most preferably 35-100 bases being hybridizable *in vivo*, under physiological



conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity. The antisense oligonucleotide can be used for downregulating heparanase activity by *in vivo* administration thereof to a patient. As such, the antisense oligonucleotide according to the present invention can be used to treat types of cancers which are characterized by impaired (over) expression of heparanase, and are dependent on the expression of heparanase for proliferating or forming metastases.

The antisense oligonucleotide can be DNA or RNA or even include nucleotide analogs, examples of which are provided in the Background section hereinabove. The antisense oligonucleotide according to the present invention can be synthetic and is preferably prepared by solid phase synthesis. In addition, it can be of any desired length which still provides specific base pairing (e.g., 8 or 10, preferably more, nucleotides long) and it can include mismatches that do not hamper base pairing under physiological conditions.

Further according to the present invention there is provided a pharmaceutical composition comprising the antisense oligonucleotide herein described and a pharmaceutically acceptable carrier. The carrier can be, for example, a liposome loadable with the antisense oligonucleotide.

According to a preferred embodiment of the present invention the antisense oligonucleotide further includes a ribozyme sequence. The

ribozyme sequence serves to cleave a heparanase RNA molecule to which the antisense oligonucleotide binds, to thereby downregulate heparanase expression.

Further according to the present invention there is provided an  
5 antisense nucleic acid construct comprising a promoter sequence and a polynucleotide sequence directing the synthesis of an antisense RNA sequence of at least 10 bases being hybridizable *in vivo*, under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity. Like the antisense oligonucleotide, the  
10 antisense construct can be used for downregulating heparanase activity by *in vivo* administration thereof to a patient. As such, the antisense construct, like the antisense oligonucleotide, according to the present invention can be used to treat types of cancers which are characterized by impaired (over) expression of heparanase, and are dependent on the expression of  
15 heparanase for proliferating or forming metastases.

Thus, further according to the present invention there is provided a pharmaceutical composition comprising the antisense construct herein described and a pharmaceutically acceptable carrier. The carrier can be, for example, a liposome loadable with the antisense construct.

20 Formulations for topical administration may include, but are not limited to, lotions, ointments, gels, creams, suppositories, drops, liquids, sprays and powders. Conventional pharmaceutical carriers, aqueous,

powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, stents, active pads, and other medical devices may also be useful. Compositions for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, sachets, capsules or tablets. Thickeners, diluents, flavorings, dispersing aids, emulsifiers or binders may be desirable. Formulations for parenteral administration may include, but are not limited to, sterile aqueous solutions which may also contain buffers, diluents and other suitable additives.

Dosing is dependent on severity and responsiveness of the condition to be treated, but will normally be one or more doses per day, week or month with course of treatment lasting from several days to several months or until a cure is effected or a diminution of disease state is achieved. Persons ordinarily skilled in the art can easily determine optimum dosages, dosing methodologies and repetition rates.

Further according to the present invention there is provided a nucleic acid construct comprising a polynucleotide sequence functioning as a promoter, the polynucleotide sequence is derived from SEQ ID NO:42 and includes at least nucleotides 2135-2635, preferably 2235-2635, more preferably 2335-2635, more preferably 2435-2635, most preferably 2535-2635 thereof, or SEQ ID NO:43 and includes at least nucleotides 1-420, preferably 120-420, more preferably 220-420, most preferably 320-420, thereof. These nucleotides are shown in the example section that

follows to direct the synthesis of a reporter gene in transformed cells. Thus, further according to the present invention there is provided a method of expressing a polynucleotide sequence comprising the step of ligating the polynucleotide sequence downstream to either of the promoter sequences  
5 described herein. Heparanase promoters can be isolated from a variety of mammalian and other species by cloning genomic regions present 5' to the coding sequence thereof. This can be readily achievable by one ordinarily skilled in the art using the heparanase polynucleotides described herein, which are shown in the Examples section that follows to participate in  
10 efficient cross species hybridization.

Further according to the present invention there is provided a recombinant protein comprising a polypeptide having heparanase catalytic activity. The protein according to the present invention include modifications known as post translational modifications, including, but not  
15 limited to, proteolysis (e.g., removal of a signal peptide and of a pro- or preprotein sequence), methionine modification, glycosylation, alkylation (e.g., methylation), acetylation, etc. According to preferred embodiments the polypeptide includes at least a portion of SEQ ID NOs:10, 14 or 44, the portion has heparanase catalytic activity. According to preferred  
20 embodiments of the present invention the protein is encoded by any of the above described isolated nucleic acids. Further according to the present

invention there is provided a pharmaceutical composition comprising, as an active ingredient, the recombinant protein described herein.

The recombinant protein may be purified by any conventional protein purification procedure close to homogeneity and/or be mixed with additives.

5 The recombinant protein may be manufactured using any of the genetically modified cells described above, which include any of the expression nucleic acid constructs described herein. The recombinant protein may be in any form. It may be in a crystallized form, a dehydrated powder form or in solution. The recombinant protein may be useful in obtaining pure  
10 heparanase, which in turn may be useful in eliciting anti-heparanase antibodies, either poly or monoclonal antibodies, and as a screening active ingredient in an anti-heparanase inhibitors or drugs screening assay or system.

Further according to the present invention there is provided a method  
15 of identifying a chromosome region harboring a human heparanase gene in a chromosome spread. the method is executed implementing the following method steps, in which in a first step the chromosome spread (either interphase or metaphase spread) is hybridized with a tagged polynucleotide probe encoding heparanase. The tag is preferably a fluorescent tag. In a  
20 second step according to the method the chromosome spread is washed, thereby excess of non-hybridized probe is removed. Finally, signals associated with the hybridized tagged polynucleotide probe are searched for,

wherein detected signals being indicative of a chromosome region harboring the human heparanase gene. One ordinarily skilled in the art would know how to use the sequences disclosed herein in suitable labeling reactions and how to use the tagged probes to detect, using *in situ* hybridization, a chromosome region harboring a human heparanase gene.

Further according to the present invention there is provided a method of *in vivo* eliciting anti-heparanase antibodies comprising the steps of administering a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*. Accordingly, there is provided also a DNA vaccine for *in vivo* eliciting anti-heparanase antibodies comprising a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*. The vaccine optionally further includes a pharmaceutically acceptable carrier, such as a virus, liposome or an antigen presenting cell. Alternatively, the vaccine is employed as a naked DNA vaccine

The present invention can be used to develop treatments for various diseases, to develop diagnostic assays for these diseases and to provide new tools for basic research especially in the fields of medicine and biology.

Specifically, the present invention can be used to develop new drugs to inhibit tumor cell metastasis, inflammation and autoimmunity. The identification of the *hpa* gene encoding for the heparanase enzyme enables the production of a recombinant enzyme in heterologous expression systems.

Furthermore, the present invention can be used to modulate bioavailability of heparin-binding growth factors, cellular responses to heparin-binding growth factors (e.g., bFGF, VEGF) and cytokines (e.g., IL-8), cell interaction with plasma lipoproteins, cellular susceptibility to viral, protozoa and some bacterial infections, and disintegration of neurodegenerative plaques. Recombinant heparanase offers a potential treatment for wound healing, angiogenesis, restenosis, atherosclerosis, inflammation, neurodegenerative diseases (such as, for example, Genstmann-Straussler Syndrome, Creutzfeldt-Jakob disease, Scrape and Alzheimer's disease) and certain viral and some bacterial and protozoa infections. Recombinant heparanase can be used to neutralize plasma heparin, as a potential replacement of protamine.

As used herein, the term "modulate" includes substantially inhibiting, slowing or reversing the progression of a disease, substantially ameliorating clinical symptoms of a disease or condition, or substantially preventing the appearance of clinical symptoms of a disease or condition. A "modulator" therefore includes an agent which may modulate a disease or condition.

Modulation of viral, protozoa and bacterial infections includes any effect which substantially interrupts, prevents or reduces any viral, bacterial or protozoa activity and/or stage of the virus, bacterium or protozoon life cycle, or which reduces or prevents infection by the virus, bacterium or protozoon in a subject, such as a human or lower animal.

As used herein, the term "wound" includes any injury to any portion of the body of a subject including, but not limited to, acute conditions such as thermal burns, chemical burns, radiation burns, burns caused by excess exposure to ultraviolet radiation such as sunburn, damage to bodily tissues such as the perineum as a result of labor and childbirth, including injuries sustained during medical procedures such as episiotomies, trauma-induced injuries including cuts, those injuries sustained in automobile and other mechanical accidents, and those caused by bullets, knives and other weapons, and post-surgical injuries, as well as chronic conditions such as pressure sores, bedsores, conditions related to diabetes and poor circulation, and all types of acne, etc.

Anti-heparanase antibodies, raised against the recombinant enzyme, would be useful for immunodetection and diagnosis of micrometastases, autoimmune lesions and renal failure in biopsy specimens, plasma samples, and body fluids. Such antibodies may also serve as neutralizing agents for heparanase activity.



The genomic heparanase sequences described herein can be used to construct knock-in and knock-out constructs. Such constructs include a fragment of 10-20 Kb of a heparanase locus and a negative and a positive selection markers and can be used to provide heparanase knock-in and knock-out animal models by methods known to the skilled artisan. Such animal models can be used for studying the function of heparanase in developmental processes, and in normal as well as pathological processes. They can also serve as an experimental model for testing drugs and gene therapy protocols. The complementary heparanase sequence (cDNA) can be used to derive transgenic animals, overexpressing heparanase for same. Alternatively, if cloned in the antisense orientation, the complementary heparanase sequence (cDNA) can be used to derive transgenic animals under-expressing heparanase for same.

The heparanase promoter sequences described herein and other cis regulatory elements linked to the heparanase locus can be used to regulated the expression of genes. For example, these promoters can be used to direct the expression of a cytotoxic protein, such as TNF, in tumor cells. It will be appreciated that heparanase itself is abnormally expressed under the control of its own promoter and other cis acting elements in a variety of tumors, and its expression is correlated with metastasis. It is also abnormally highly expressed in inflammatory cells. The introns of the heparanase gene can be used for the same purpose, as it is known that

introns, especially upstream introns include cis acting element which affect expression. A heparanase promoter fused to a reporter protein can be used to study/monitor its activity.

The polynucleotide sequences described herein can also be used to provide DNA vaccines which will elicit in vivo anti heparanase antibodies. Such vaccines can therefore be used to combat inflammatory and cancer.

Antisense oligonucleotides derived according to the heparanase sequences described herein, especially such oligonucleotides supplemented with ribozyme activity, can be used to modulate heparanase expression.

Such oligonucleotides can be from the coding region, from the introns or promoter specific. Antisense heparanase nucleic acid constructs can similarly function, as well known in the art.

The heparanase sequences described herein can be used to study the catalytic mechanism of heparanase. Carefully selected site directed mutagenesis can be employed to provide modified heparanase proteins having modified characteristics in terms of, for example, substrate specificity, sensitivity to inhibitors, etc.

While studying heparanase expression in a variety of cell types alternatively spliced transcripts were identified. Such transcripts if found characteristic of certain pathological conditions can be used as markers for such conditions. Such transcripts are expected to direct the synthesis of heparanases with altered functions.

Additional objects, advantages, and novel features of the present invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting. Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

### EXAMPLES

Generally, the nomenclature used herein and the laboratory procedures in recombinant DNA technology described below are those well known and commonly employed in the art. Standard techniques are used for cloning, DNA and RNA isolation, amplification and purification. Generally enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like are performed according to the manufacturers' specifications. These techniques and various other techniques are generally performed according to Sambrook et al., Molecular Cloning--A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1989), which is incorporated herein by reference. Other general references are provided throughout this document. The procedures therein are believed to be well known in the art and are provided for the convenience of the reader. All the information contained therein is incorporated herein by reference.

The following protocols and experimental details are referenced in the Examples that follow:

*Purification and characterization of heparanase from a human*

5 *hepatoma cell line and human placenta:* A human hepatoma cell line (Sk-hep-1) was chosen as a source for purification of a human tumor-derived heparanase. Purification was essentially as described in U.S. Pat. No. 5,362,641 to Fuks, which is incorporated by reference as if fully set forth herein. Briefly, 500 liter,  $5 \times 10^{11}$  cells were grown in suspension and  
10 the heparanase enzyme was purified about 240,000 fold by applying the following steps: (i) cation exchange (CM-Sephadex) chromatography performed at pH 6.0, 0.3-1.4 M NaCl gradient; (ii) cation exchange (CM-Sephadex) chromatography performed at pH 7.4 in the presence of 0.1% CHAPS, 0.3-1.1 M NaCl gradient; (iii) heparin-Sepharose  
15 chromatography performed at pH 7.4 in the presence of 0.1% CHAPS, 0.35-1.1 M NaCl gradient; (iv) ConA-Sepharose chromatography performed at pH 6.0 in buffer containing 0.1 % CHAPS and 1 M NaCl, elution with 0.25 M  $\alpha$ -methyl mannoside; and (v) HPLC cation exchange (Mono-S) chromatography performed at pH 7.4 in the presence of 0.1 % CHAPS,  
20 0.25-1 M NaCl gradient.

Active fractions were pooled, precipitated with TCA and the precipitate subjected to SDS polyacrylamide gel electrophoresis and/or

tryptic digestion and reverse phase HPLC. Tryptic peptides of the purified protein were separated by reverse phase HPLC (C8 column) and homogeneous peaks were subjected to amino acid sequence analysis.

The purified enzyme was applied to reverse phase HPLC and  
5 subjected to N-terminal amino acid sequencing using the amino acid sequencer (Applied Biosystems).

*Cells:* Cultures of bovine corneal endothelial cells (BCECs) were established from steer eyes as previously described (19, 38). Stock cultures were maintained in DMEM (1 g glucose/liter) supplemented with 10 %  
10 newborn calf serum and 5 % FCS. bFGF (1 ng/ml) was added every other day during the phase of active cell growth (13, 14).

*Preparation of dishes coated with ECM:* BCECs (second to fifth passage) were plated into 4-well plates at an initial density of  $2 \times 10^5$  cells/ml, and cultured in sulfate-free Fisher medium plus 5 % dextran T-40  
15 for 12 days.  $\text{Na}_2^{35}\text{SO}_4$  (25  $\mu\text{Ci/ml}$ ) was added on day 1 and 5 after seeding and the cultures were incubated with the label without medium change. The subendothelial ECM was exposed by dissolving (5 min., room temperature) the cell layer with PBS containing 0.5 % Triton X-100 and 20 mM  $\text{NH}_4\text{OH}$ , followed by four washes with PBS. The ECM remained intact, free of  
20 cellular debris and firmly attached to the entire area of the tissue culture dish (19, 22).

To prepare soluble sulfate labeled proteoglycans (peak I material), the ECM was digested with trypsin (25  $\mu\text{g/ml}$ , 6 h, 37  $^{\circ}\text{C}$ ), the digest was concentrated by reverse dialysis and the concentrated material was applied onto a Sepharose 6B gel filtration column. The resulting high molecular weight material ( $K_{av} < 0.2$ , peak I) was collected. More than 80 % of the labeled material was shown to be composed of heparan sulfate proteoglycans (11, 39).

*Heparanase activity:* Cells ( $1 \times 10^6$ /35-mm dish), cell lysates or conditioned media were incubated on top of  $^{35}\text{S}$ -labeled ECM (18 h, 37  $^{\circ}\text{C}$ ) in the presence of 20 mM phosphate buffer (pH 6.2). Cell lysates and conditioned media were also incubated with sulfate labeled peak I material (10-20  $\mu\text{l}$ ). The incubation medium was collected, centrifuged (18,000  $\times g$ , 4  $^{\circ}\text{C}$ , 3 min.), and sulfate labeled material analyzed by gel filtration on a Sepharose CL-6B column (0.9  $\times$  30 cm). Fractions (0.2 ml) were eluted with PBS at a flow rate of 5 ml/h and counted for radioactivity using Bio-fluor scintillation fluid. The excluded volume ( $V_o$ ) was marked by blue dextran and the total included volume ( $V_t$ ) by phenol red. The latter was shown to comigrate with free sulfate (7, 11, 23). Degradation fragments of HS side chains were eluted from Sepharose 6B at  $0.5 < K_{av} < 0.8$  (peak II) (7, 11, 23). A nearly intact HSPG released from ECM by trypsin - and, to a lower extent, during incubation with PBS alone - was eluted next to  $V_o$  ( $K_{av} < 0.2$ , peak I). Recoveries of labeled material applied on the columns

ranged from 85 to 95 % in different experiments (11). Each experiment was performed at least three times and the variation of elution positions ( $K_{av}$  values) did not exceed  $\pm 15$  %.

**Cloning of *hpa* cDNA:** cDNA clones 257548 and 260138 were  
5 obtained from the I.M.A.G.E Consortium (2130 Memorial Parkway SW, Hunstville, AL 35801). The cDNAs were originally cloned in *Eco*RI and *Not*I cloning sites in the plasmid vector pT3T7D-Pac. Although these clones are reported to be somewhat different, DNA sequencing demonstrated that these clones are identical to one another. Marathon  
10 RACE (rapid amplification of cDNA ends) human placenta (poly-A) cDNA composite was a gift of Prof. Yossi Shiloh of Tel Aviv University. This composite is vector free, as it includes reverse transcribed cDNA fragments to which double, partially single stranded adapters are attached on both sides. The construction of the specific composite employed is described in  
15 reference 39a.

Amplification of hp3 PCR fragment was performed according to the protocol provided by Clontech laboratories. The template used for amplification was a sample taken from the above composite. The primers used for amplification were:

20 First step: 5'-primer: AP1: 5'-CCATCCTAATACGACTCACT ATAGGGC-3', SEQ ID NO:1; 3'-primer: HPL229: 5'-GTAGTGATGCCA TGTA ACTGAATC-3', SEQ ID NO:2.

Second step: nested 5'-primer: AP2: 5'-ACTCACTATAGGGGCTCG  
 AGCGGC-3', SEQ ID NO:3; nested 3'- primer: HPL171:  
 5'-GCATCTTAGCCGTCTTTCTTCG-3', SEQ ID NO:4. The HPL229 and  
 HPL171 were selected according to the sequence of the EST clones. They  
 5 include nucleotides 933-956 and 876-897 of SEQ ID NO:9, respectively.

PCR program was 94 °C - 4 min., followed by 30 cycles of 94 °C -  
 40 sec., 62 °C - 1 min., 72 °C - 2.5 min. Amplification was performed with  
 Expand High Fidelity (Boehringer Mannheim). The resulting ca. 900 bp  
 hp3 PCR product was digested with *Bfr*I and *Pvu*II. Clone 257548 (*phpa*1)  
 10 was digested with *Eco*RI, followed by end filling and was then further  
 digested with *Bfr*I. Thereafter the *Pvu*II - *Bfr*I fragment of the hp3 PCR  
 product was cloned into the blunt end - *Bfr*I end of clone *phpa*1 which  
 resulted in having the entire cDNA cloned in pT3T7-pac vector, designated  
*phpa*2.

15 **RT-PCR:** RNA was prepared using TRI-Reagent (Molecular  
 research center Inc.) according to the manufacturer instructions. 1.25 µg  
 were taken for reverse transcription reaction using MuMLV Reverse  
 transcriptase (Gibco BRL) and Oligo (dT)<sub>15</sub> primer, SEQ ID NO:5,  
 (Promega). Amplification of the resultant first strand cDNA was  
 20 performed with *Taq* polymerase (Promega). The following primers were  
 used:

HPU-355: 5'-TTCGATCCCAAGAAGGAATCAAC-3', SEQ ID NO:6,



nucleotides 372-394 in SEQ ID NOs:9 or 11.

HPL-229: 5'-GTAGTGATGCCATGTAAGTGAATC-3', SEQ ID NO:7,

nucleotides 933-956 in SEQ ID NOs:9 or 11.

PCR program: 94 °C - 4 min., followed by 30 cycles of 94 °C - 40  
5 sec., 62 °C - 1 min., 72 °C - 1 min.

Alternatively, total RNA was prepared from cell cultures using  
Tri-reagent (Molecular Research Center, Inc.) according to the  
manufacturer recommendation. Poly A+ RNA was isolated from total RNA  
using mRNA separator (Clontech). Reverse transcription was performed  
10 with total RNA using Superscript II (GibcoBRL). PCR was performed with  
Expand high fidelity (Boehringer Mannheim). Primers used for  
amplification were as follows:

Hpu-685, 5'-GAGCAGCCAGGTGAGCCCAAGAT-3', SEQ ID NO:24

Hpu-355, 5'-TTCGATCCCAAGAAGGAATCAAC-3', SEQ ID NO:25

15 Hpu 565, 5'-AGCTCTGTAGATGTGCTATACAC-3', SEQ ID NO:26

Hpl 967, 5'-TCAGATGCAAGCAGCAACTTTGGC-3', SEQ ID NO:27

Hpl 171, 5'-GCATCTTAGCCGTCTTTCTTCG-3', SEQ ID NO:28

Hpl 229, 5'-GTAGTGATGCCATGTAAGTGAATC-3', SEQ ID NO:29

PCR reaction was performed as follows: 94 °C 3 minutes, followed  
20 by 32 cycles of 94 °C 40 seconds, 64 °C 1 minute, 72 °C 3 minutes, and one  
cycle 72 °C, 7 minutes.

**Expression of recombinant heparanase in insect cells:** Cells, High Five and Sf21 insect cell lines were maintained as monolayer cultures in SF900II-SFM medium (GibcoBRL).

**Recombinant Baculovirus:** Recombinant virus containing the *hpa* gene was constructed using the Bac to Bac system (GibcoBRL). The transfer vector pFastBac was digested with *SalI* and *NotI* and ligated with a 1.7 kb fragment of *phpa2* digested with *XhoI* and *NotI*. The resulting plasmid was designated pFast*hpa2*. An identical plasmid designated pFast*hpa4* was prepared as a duplicate and both independently served for further experimentations. Recombinant bacmid was generated according to the instructions of the manufacturer with pFast*hpa2*, pFast*hpa4* and with pFastBac. The latter served as a negative control. Recombinant bacmid DNAs were transfected into Sf21 insect cells. Five days after transfection recombinant viruses were harvested and used to infect High Five insect cells,  $3 \times 10^6$  cells in T-25 flasks. Cells were harvested 2 - 3 days after infection.  $4 \times 10^6$  cells were centrifuged and resuspended in a reaction buffer containing 20 mM phosphate citrate buffer, 50 mM NaCl. Cells underwent three cycles of freeze and thaw and lysates were stored at  $-80^\circ\text{C}$ . Conditioned medium was stored at  $4^\circ\text{C}$ .

**Partial purification of recombinant heparanase:** Partial purification of recombinant heparanase was performed by heparin-Sepharose column chromatography followed by Superdex 75

column gel filtration. Culture medium (150 ml) of Sf21 cells infected with pFhpa4 virus was subjected to heparin-Sepharose chromatography. Elution of 1 ml fractions was performed with 0.35 - 2 M NaCl gradient in presence of 0.1 % CHAPS and 1 mM DTT in 10 mM sodium acetate buffer, pH 5.0.

5 A 25  $\mu$ l sample of each fraction was tested for heparanase activity. Heparanase activity was eluted at the range of 0.65 - 1.1 M NaCl (fractions 18-26, Figure 10a). 5  $\mu$ l of each fraction was subjected to 15 % SDS-polyacrylamide gel electrophoresis followed by silver nitrate staining. Active fractions eluted from heparin-Sepharose (Figure 10a) were pooled  
10 and concentrated (x 6) on YM3 cut-off membrane. 0.5 ml of the concentrated material was applied onto 30 ml Superdex 75 FPLC column equilibrated with 10 mM sodium acetate buffer, pH 5.0, containing 0.8 M NaCl, 1 mM DTT and 0.1 % CHAPS. Fractions (0.56 ml) were collected at a flow rate of 0.75 ml/min. Aliquots of each fraction were tested for  
15 heparanase activity and were subjected to SDS-polyacrylamide gel electrophoresis followed by silver nitrate staining (Figure 11b).

*PCR amplification of genomic DNA:* 94 °C 3 minutes, followed by 32 cycles of 94 °C 45 seconds, 64 °C 1 minute, 68 °C 5 minutes, and one cycle at 72 °C, 7 minutes. Primers used for amplification of genomic DNA  
20 included:

GHpu-L3 5'-AGGCACCCTAGAGATGTTCCAG-3', SEQ ID NO:30

GHpl-L6 5'-GAAGATTTCTGTTTCCATGACGTG-3', SEQ ID NO:31.

**Screening of genomic libraries:** A human genomic library in Lambda phage EMBLE3 SP6/T7 (Clontech, Paulo Alto, CA) was screened.  $5 \times 10^5$  plaques were plated at  $5 \times 10^4$  pfu/plate on NZCYM agar/top agarose plates. Phages were absorbed on nylon membranes in duplicates (Qiagen). Hybridization was performed at 65 °C in 5 x SSC, 5 x Denhart's, 10 % dextran sulfate, 100 µg/ml Salmon sperm,  $^{32}\text{p}$  labeled probe ( $10^6$  cpm/ml). A 1.6 kb fragment, containing the entire *hpa* cDNA was labeled by random priming (Boehringer Mannheim). Following hybridization membranes were washed once with 2 x SSC, 0.1 % SDS at 65 °C for 20 minutes, and twice with 0.2 x SSC, 0.1 % SDS at 65 °C for 15 minutes. Hybridizing plaques were picked, and plated at 100 pfu/plate. Hybridization was performed as above and single isolated positive plaques were picked.

Phage DNA was extracted using a Lambda DNA extraction kit (Qiagen). DNA was digested with *Xho*I and *Eco*RI, separated on 0.7 % agarose gel and transferred to nylon membrane Hybond N+ (Amersham). Hybridization and washes were performed as above.

**cDNA Sequence analysis:** Sequence determinations were performed with vector specific and gene specific primers, using an automated DNA sequencer (Applied Biosystems, model 373A). Each nucleotide was read from at least two independent primers.

**Genomic sequence analysis:** Large-scale sequencing was performed by Commonwealth Biotechnology Incorporation.

**Isolation of mouse *hpa*:** Mouse *hpa* cDNA was amplified from either Marathon ready cDNA library of mouse embryo or from mRNA isolated from mouse melanoma cell line BL6, using the Marathon RACE kit from Clontech. Both procedures were performed according to the manufacturer's recommendation.

***Primers used for PCR amplification of mouse *hpa*:***

Mhpl773 5'-CCACACTGAATGTAATACTGAAGTG-3', SEQ ID NO:32

10 MHpl736 5'-CGAAGCTCTGGAAGCTCGGCAAG-3', SEQ ID NO:33

MHpl83 5'-GCCAGCTGCAAAGGTGTTGGAC-3', SEQ ID NO:34

Mhpl152 5'-AACACCTGCCTCATCACGACTTC-3', SEQ ID NO:35

Mhpl114 5'-GCCAGGCTGGCGTCGATGGTGA-3', SEQ ID NO:36

MHpl103 5'-GTCGATGGTGATGGACAGGAAC-3', SEQ ID NO:37

15 Ap1 5'-GTAATACGACTCACTATAGGGC-3', SEQ ID NO:38 -

(Genome walker)

Ap2 5'-ACTATAGGGCACGCGTGGT-3', SEQ ID NO:39 -

(Genome walker)

Ap1 5'-CCATCCTAATACGACTCACTATAGGGC-3', SEQ ID NO:40 -

20 (Marathon RACE)

Ap2 5'-ACTCACTATAGGGCTCGAGCGGC-3', SEQ ID NO:41 -

(Marathon RACE)

***Southern analysis of genomic DNA:*** Genomic DNA was extracted from animal or from human blood using Blood and cell culture DNA maxi kit (Qiagene). DNA was digested with *EcoRI*, separated by gel electrophoresis and transferred to a nylon membrane Hybond N+ (Amersham). Hybridization was performed at 68 °C in 6 x SSC, 1 % SDS, 5 x Denharts, 10 % dextran sulfate, 100 µg/ml salmon sperm DNA, and <sup>32</sup>p labeled probe. A 1.6 kb fragment, containing the entire *hpa* cDNA was used as a probe. Following hybridization, the membrane was washed with 3 x SSC, 0.1 % SDS, at 68 °C and exposed to X-ray film for 3 days. Membranes were then washed with 1 x SSC, 0.1 % SDS, at 68 °C and were reexposed for 5 days.

***Construction of hpa promoter-GFP expression vector:*** Lambda DNA of phage L3, was digested with *SacI* and *BglII*, resulting in a 1712 bp fragment which contained the *hpa* promoter (877-2688 of SEQ ID NO:42). The pEGFP-1 plasmid (Clontech) was digested with *BglII* and *SacI* and ligated with the 1712 bp fragment of the *hpa* promoter sequence. The resulting plasmid was designated phpEGL. A second *hpa* promoter-GFP plasmid was constructed containing a shorter fragment of the *hpa* promoter region: phpEGL was digested with *HindIII*, and the resulting 1095 bp fragment (nucleotides 1593-2688 of SEQ ID NO:42) was ligated with *HindIII* digested pEGFP-1. The resulting plasmid was designated phpEGS.

*Computer analysis of sequences:* Homology searches were

performed using several computer servers, and various databases. Blast 2.0 service, at the NCBI server was used to screen the protein database swplus and DNA databases such as GenBank, EMBL, and the EST databases.

5 Blast 2.0 search was performed using the basic search option of the NCBI server. Sequence analysis and alignments were done using the DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the university of Wisconsin. Alignments of two sequences were performed using Bestfit (gap creation penalty - 12, gap extension  
10 penalty - 4). Protein homology search was performed with the Smith-Waterman algorithm, using the Bioaccelerator platform developed by Compugene. The protein database swplus was searched using the following parameters: gapop: 10.0, gapext: 0.5, matrix: blosum62. Blocks homology was performed using the Blocks WWW server developed at Fred  
15 Hutchinson Cancer Research Center in Seattle, Washington, USA. Secondary structure prediction was performed using the PHD server – Profile network Prediction Heidelberg. Fold recognition (threading) was performed using the UCLA-DOE structure prediction server. The method used for prediction was gonnet+predss. Alignment of three sequences was  
20 performed using the pileup application (gap creation penalty - 5, gap extension penalty - 1). Promoter analysis was performed using TSSW and

TSSG programs (BCM Search Launcher Human Genome Center, Baylor College of Medicine, Houston TX).

### EXAMPLE 1

#### 5                    *Cloning of human hpa cDNA*

Purified fraction of heparanase isolated from human hepatoma cells (SK-hep-1) was subjected to tryptic digestion and microsequencing. EST (Expressed Sequence Tag) databases were screened for homology to the back translated DNA sequences corresponding to the obtained peptides.

10    Two EST sequences (accession Nos. N41349 and N45367) contained a DNA sequence encoding the peptide YGPDVGQPR (SEQ ID NO:8). These two sequences were derived from clones 257548 and 260138 (I.M.A.G.E Consortium) prepared from 8 to 9 weeks placenta cDNA library (Soares). Both clones which were found to be identical contained an insert

15    of 1020 bp which included an open reading frame (ORF) of 973 bp followed by a 3' untranslated region of 27 bp and a Poly A tail. No translation start site (AUG) was identified at the 5' end of these clones.

Cloning of the missing 5' end was performed by PCR amplification of DNA from a placenta Marathon RACE cDNA composite. A 900 bp

20    fragment (designated hp3), partially overlapping with the identified 3' encoding EST clones was obtained.



The joined cDNA fragment, 1721 bp long (SEQ ID NO:9), contained an open reading frame which encodes, as shown in Figure 1 and SEQ ID NO:11, a polypeptide of 543 amino acids (SEQ ID NO:10) with a calculated molecular weight of 61,192 daltons. The 3' end of the partial cDNA inserts  
5 contained in clones 257548 and 260138 started at nucleotide G<sup>721</sup> of SEQ ID NO:9 and Figure 1.

As further shown in Figure 1, there was a single sequence discrepancy between the EST clones and the PCR amplified sequence, which led to an amino acid substitution from Tyr<sup>246</sup> in the EST to Phe<sup>246</sup> in  
10 the amplified cDNA. The nucleotide sequence of the PCR amplified cDNA fragment was verified from two independent amplification products. The new gene was designated *hpa*.

As stated above, the 3' end of the partial cDNA inserts contained in EST clones 257548 and 260138 started at nucleotide 721 of *hpa* (SEQ ID  
15 NO:9). The ability of the *hpa* cDNA to form stable secondary structures, such as stem and loop structures involving nucleotide stretches in the vicinity of position 721 was investigated using computer modeling. It was found that stable stem and loop structures are likely to be formed involving nucleotides 698-724 (SEQ ID NO:9). In addition, a high GC content, up to  
20 70 %, characterizes the 5' end region of the *hpa* gene, as compared to about only 40 % in the 3' region. These findings may explain the immature termination and therefore lack of 5' ends in the EST clones.

To examine the ability of the *hpa* gene product to catalyze degradation of heparan sulfate in an *in vitro* assay the entire open reading frame was expressed in insect cells, using the Baculovirus expression system. Extracts of cells, infected with virus containing the *hpa* gene, demonstrated a high level of heparan sulfate degradation activity, while cells infected with a similar construct containing no *hpa* gene had no such activity, nor did non-infected cells. These results are further demonstrated in the following Examples.

## EXAMPLE 2

### *Degradation of soluble ECM-derived HSPG*

Monolayer cultures of High Five cells were infected (72 h, 28 °C) with recombinant Baculovirus containing the pFast*hpa* plasmid or with control virus containing an insert free plasmid. The cells were harvested and lysed in heparanase reaction buffer by three cycles of freezing and thawing. The cell lysates were then incubated (18 h, 37 °C) with sulfate labeled, ECM-derived HSPG (peak I), followed by gel filtration analysis (Sephacrose 6B) of the reaction mixture.

As shown in Figure 2, the substrate alone included almost entirely high molecular weight (Mr) material eluted next to  $V_0$  (peak I, fractions 5-20,  $K_{av} < 0.35$ ). A similar elution pattern was obtained when the HSPG substrate was incubated with lysates of cells that were infected with control

virus. In contrast, incubation of the HSPG substrate with lysates of cells infected with the *hpa* containing virus resulted in a complete conversion of the high Mr substrate into low Mr labeled degradation fragments (peak II, fractions 22-35,  $0.5 < K_{av} < 0.75$ ).

5           Fragments eluted in peak II were shown to be degradation products of heparan sulfate, as they were (i) 5- to 6-fold smaller than intact heparan sulfate side chains ( $K_{av}$  approx. 0.33) released from ECM by treatment with either alkaline borohydride or papain; and (ii) resistant to further digestion with papain or chondroitinase ABC, and susceptible to deamination by  
10   nitrous acid (6, 11).           Similar results (not shown) were obtained with Sf21 cells. Again, heparanase activity was detected in cells infected with the *hpa* containing virus (pF*hpa*), but not with control virus (pF). This result was obtained with two independently generated recombinant viruses. Lysates of control not infected High Five cells failed to degrade the HSPG  
15   substrate.

In subsequent experiments, the labeled HSPG substrate was incubated with medium conditioned by infected High Five or Sf21 cells.

As shown in Figures 3a-b, heparanase activity, reflected by the conversion of the high Mr peak I substrate into the low Mr peak II which  
20   represents HS degradation fragments, was found in the culture medium of cells infected with the pF*hpa2* or pF*hpa4* viruses, but not with the control

pF1 or pF2 viruses. No heparanase activity was detected in the culture medium of control non-infected High Five or Sf21 cells.

The medium of cells infected with the pF*hpa4* virus was passed through a 50 kDa cut off membrane to obtain a crude estimation of the molecular weight of the recombinant heparanase enzyme. As demonstrated in Figure 4, all the enzymatic activity was retained in the upper compartment and there was no activity in the flow through (<50 kDa) material. This result is consistent with the expected molecular weight of the *hpa* gene product.

In order to further characterize the *hpa* product the inhibitory effect of heparin, a potent inhibitor of heparanase mediated HS degradation (40) was examined.

As demonstrated in Figures 5a-b, conversion of the peak I substrate into peak II HS degradation fragments was completely abolished in the presence of heparin.

Altogether, these results indicate that the heparanase enzyme is expressed in an active form by insect cells infected with Baculovirus containing the newly identified human *hpa* gene.

### EXAMPLE 3

#### *Degradation of HSPG in intact ECM*

Next, the ability of intact infected insect cells to degrade HS in intact, naturally produced ECM was investigated. For this purpose, High Five or Sf21 cells were seeded on metabolically sulfate labeled ECM followed by infection (48 h, 28 °C) with either the pFhpa4 or control pF2 viruses. The pH of the medium was then adjusted to pH 6.2-6.4 and the cells further incubated with the labeled ECM for another 48 h at 28 °C or 24 h at 37 °C. Sulfate labeled material released into the incubation medium was analyzed by gel filtration on Sepharose 6B.

As shown in Figures 6a-b and 7a-b, incubation of the ECM with cells infected with the control pF2 virus resulted in a constant release of labeled material that consisted almost entirely (>90%) of high Mr fragments (peak I) eluted with or next to  $V_0$ . It was previously shown that a proteolytic activity residing in the ECM itself and/or expressed by cells is responsible for release of the high Mr material (6). This nearly intact HSPG provides a soluble substrate for subsequent degradation by heparanase, as also indicated by the relatively large amount of peak I material accumulating when the heparanase enzyme is inhibited by heparin (6, 7, 12, Figure 9). On the other hand, incubation of the labeled ECM with cells infected with the pFhpa4 virus resulted in release of 60-70% of the ECM-associated radioactivity in the form of low Mr sulfate-labeled fragments (peak II, 0.5

<Kav< 0.75), regardless of whether the infected cells were incubated with the ECM at 28 °C or 37 °C. Control intact non-infected Sf21 or High Five cells failed to degrade the ECM HS side chains.

In subsequent experiments, as demonstrated in Figures 8a-b, High Five and Sf21 cells were infected (96 h, 28 °C) with pFhpa4 or control pF1 viruses and the culture medium incubated with sulfate-labeled ECM. Low Mr HS degradation fragments were released from the ECM only upon incubation with medium conditioned by pFhpa4 infected cells. As shown in Figure 9, production of these fragments was abolished in the presence of heparin. No heparanase activity was detected in the culture medium of control, non-infected cells. These results indicate that the heparanase enzyme expressed by cells infected with the pFhpa4 virus is capable of degrading HS when complexed to other macromolecular constituents (i.e. fibronectin, laminin, collagen) of a naturally produced intact ECM, in a manner similar to that reported for highly metastatic tumor cells or activated cells of the immune system (6, 7).

#### EXAMPLE 4

##### *Purification of recombinant human heparanase*

The recombinant heparanase was partially purified from medium of pFhpa4 infected Sf21 cells by Heparin-Sepharose chromatography (Figure 10a) followed by gel filtration of the pooled active fractions over an FPLC

Superdex 75 column (Figure 11a). A ~ 63 kDa protein was observed, whose quantity, as was detected by silver stained SDS-polyacrylamide gel electrophoresis, correlated with heparanase activity in the relevant column fractions (Figures 10b and 11b, respectively). This protein was not detected  
5 in the culture medium of cells infected with the control pF1 virus and was subjected to a similar fractionation on heparin-Sepharose (not shown).

### EXAMPLE 5

#### *Expression of the human hpa cDNA in various cell types, organs and 10 tissues*

Referring now to Figures 12a-e, RT-PCR was applied to evaluate the expression of the *hpa* gene by various cell types and tissues. For this purpose, total RNA was reverse transcribed and amplified. The expected 585 bp long cDNA was clearly demonstrated in human kidney, placenta (8  
15 and 11 weeks) and mole tissues, as well as in freshly isolated and short termed (1.5-48 h) cultured human placental cytotrophoblastic cells (Figure 12a), all known to express a high heparanase activity (41). The *hpa* transcript was also expressed by normal human neutrophils (Figure 12b). In contrast, there was no detectable expression of the *hpa* mRNA in embryonic  
20 human muscle tissue, thymus, heart and adrenal (Figure 12b). The *hpa* gene was expressed by several, but not all, human bladder carcinoma cell lines (Figure 12c), SK hepatoma (SK-hep-1), ovarian carcinoma (OV 1063),

breast carcinoma (435, 231), melanoma and megakaryocytic (DAMI, CHRF) human cell lines (Figures 12d-e).

The above described expression pattern of the *hpa* transcript was determined to be in a very good correlation with heparanase activity levels

determined in various tissues and cell types (not shown).

## EXAMPLE 6

### *Isolation of an extended 5' end of hpa cDNA from human SK-hep1 cell line*

The 5' end of *hpa* cDNA was isolated from human SK-hep1 cell line by PCR amplification using the Marathon RACE (rapid amplification of cDNA ends) kit (Clontech). Total RNA was prepared from SK-hep1 cells using the TRI-Reagent (Molecular research center Inc.) according to the manufacturer instructions. Poly A<sup>+</sup> RNA was isolated using the mRNA separator kit (Clontech).

The Marathon RACE SK-hep1 cDNA composite was constructed according to the manufacturer recommendations. First round of amplification was performed using an adaptor specific primer AP1: 5'-CCATCCTAATACG ACTCACTATAGGGC-3', SEQ ID NO:1, and a *hpa* specific antisense primer hpl-629: 5'-CCCCAGGAGCAGCAGCATCAG-3', SEQ ID NO:17, corresponding to nucleotides 119-99 of SEQ ID NO:9. The resulting PCR product was



subjected to a second round of amplification using an adaptor specific nested primer AP2: 5'-ACTCACTATAGGGCTCGAGCGGC-3', SEQ ID NO:3, and a *hpa* specific antisense nested primer hpl-666 5'-AGGCTTCGAGCGCAGCAGCAT-3', SEQ ID NO:18, corresponding to nucleotides 83-63 of SEQ ID NO:9. The PCR program was as follows: a hot start of 94 °C for 1 minute, followed by 30 cycles of 90 °C - 30 seconds, 68 °C - 4 minutes. The resulting 300 bp DNA fragment was extracted from an agarose gel and cloned into the vector pGEM-T Easy (Promega). The resulting recombinant plasmid was designated pHPSK1.

The nucleotide sequence of the pHPSK1 insert was determined and it was found to contain 62 nucleotides of the 5' end of the placenta *hpa* cDNA (SEQ ID NO:9) and additional 178 nucleotides upstream, the first 178 nucleotides of SEQ ID NOs:13 and 15.

A single nucleotide discrepancy was identified between the SK-hep1 cDNA and the placenta cDNA. The "T" derivative at position 9 of the placenta cDNA (SEQ ID NO:9), is replaced by a "C" derivative at the corresponding position 187 of the SK-hep1 cDNA (SEQ ID NO:13).

The discrepancy is likely to be due to a mutation at the 5' end of the placenta cDNA clone as confirmed by sequence analysis of several additional cDNA clones isolated from placenta, which like the SK-hep1 cDNA contained C at position 9 of SEQ ID NO:9.

The 5' extended sequence of the SK-hepI *hpa* cDNA was assembled with the sequence of the *hpa* cDNA isolated from human placenta (SEQ ID NO:9). The assembled sequence contained an open reading frame which encodes, as shown in SEQ ID NOs:14 and 15, a polypeptide of 592 amino acids with a calculated molecular weight of 66,407 daltons. The open reading frame is flanked by 93 bp 5' untranslated region (UTR).

### EXAMPLE 7

#### *Isolation of the upstream genomic region of the hpa gene*

The upstream region of the *hpa* gene was isolated using the Genome Walker kit (Clontech) according to the manufacturer recommendations. The kit includes five human genomic DNA samples each digested with a different restriction endonuclease creating blunt ends: *EcoRV*, *ScaI*, *DraI*, *PvuII* and *SspI*.

The blunt ended DNA fragments are ligated to partially single stranded adaptors. The Genomic DNA samples were subjected to PCR amplification using the adaptor specific primer and a gene specific primer. Amplification was performed with Expand High Fidelity (Boehringer Mannheim).

A first round of amplification was performed using the ap1 primer: 5'-G TAATACGACTCACTATAGGGC-3', SEQ ID NO:19, and the *hpa* specific antisense primer hpl-666:

5'-AGGCTTCGAGCGCAGCAGCAT-3', SEQ ID NO:18, corresponding to nucleotides 83 – 63 of SEQ ID NO:9. The PCR program was as follows: a hot start of 94 °C - 3 minutes, followed by 36 cycles of 94 °C - 40 seconds, 67 °C - 4 minutes.

5        The PCR products of the first amplification were diluted 1:50. One µl of the diluted sample was used as a template for a second amplification using a nested adaptor specific primer ap2: 5'-ACTATAGGGCACGCGTGGT-3', SEQ ID NO:20, and a *hpa* specific antisense primer hpl-690, 5'-CTTGGGCTCACC TGGCTGCTC-3', SEQ ID  
10 NO:21, corresponding to nucleotides 62-42 of SEQ ID NO:9. The resulting amplification products were analyzed using agarose gel electrophoresis. Five different PCR products were obtained from the five amplification reactions. A DNA fragment of approximately 750 bp which was obtained from the *SspI* digested DNA sample was gel extracted. The purified  
15 fragment was ligated into the plasmid vector pGEM-T Easy (Promega). The resulting recombinant plasmid was designated pGHP6905 and the nucleotide sequence of the *hpa* insert was determined.

A partial sequence of 594 nucleotides is shown in SEQ ID NO:16. The last nucleotide in SEQ ID NO:13 corresponds to nucleotide 93 in SEQ  
20 ID:13. The DNA sequence in SEQ ID NO:16 contains the 5' region of the *hpa* cDNA and 501 nucleotides of the genomic upstream region which are predicted to contain the promoter region of the *hpa* gene.

**EXAMPLE 8*****Expression of the 592 amino acids HPA polypeptide in a human 293 cell line***

The 592 amino acids open reading frame (SEQ ID NOs:13 and 15) was constructed by ligation of the 110 bp corresponding to the 5' end of the SK-hep1 *hpa* cDNA with the placenta cDNA. More specifically the Marathon RACE - PCR amplification product of the placenta *hpa* DNA was digested with *SacI* and an approximately 1 kb fragment was ligated into a *SacI*-digested pGHP6905 plasmid. The resulting plasmid was digested with *EarI* and *AatII*. The *EarI* sticky ends were blunted and an approximately 280 bp *EarI*/blunt-*AatII* fragment was isolated. This fragment was ligated with pFast*hpa* digested with *EcoRI* which was blunt ended using Klenow fragment and further digested with *AatII*. The resulting plasmid contained a 1827 bp insert which includes an open reading frame of 1776 bp, 31 bp of 3' UTR and 21 bp of 5' UTR. This plasmid was designated pFast*Lhpa*.

A mammalian expression vector was constructed to drive the expression of the 592 amino acids heparanase polypeptide in human cells. The *hpa* cDNA was excised from pFast*Lhpa* with *BssHII* and *NotI*. The resulting 1850 bp *BssHII*-*NotI* fragment was ligated to a mammalian expression vector pSI (Promega) digested with *MluI* and *NotI*. The resulting recombinant plasmid, pSI*hpa*Met2 was transfected into a human 293 embryonic kidney cell line.

Transient expression of the 592 amino-acids heparanase was examined by western blot analysis and the enzymatic activity was tested using the gel shift assay. Both these procedures are described in length in U.S. Pat. application No. 09/071,739, filed May 1, 1998, which is  
5 incorporated by reference as if fully set forth herein. Cells were harvested 3 days following transfection. Harvested cells were re-suspended in lysis buffer containing 150 mM NaCl, 50 mM Tris pH 7.5, 1% Triton X-100, 1 mM PMSF and protease inhibitor cocktail (Boehringer Mannheim). 40 µg protein extract samples were used for separation on a SDS-PAGE. Proteins  
10 were transferred onto a PVDF Hybond-P membrane (Amersham). The membrane was incubated with an affinity purified polyclonal anti heparanase antibody, as described in U.S. Pat. application No. 09/071,739. A major band of approximately 50 kDa was observed in the transfected cells as well as a minor band of approximately 65 kDa. A similar pattern  
15 was observed in extracts of cells transfected with the *pShpa* as demonstrated in U.S. Pat. application No. 09/071,739. These two bands probably represent two forms of the recombinant heparanase protein produced by the transfected cells. The 65 kDa protein probably represents a heparanase precursor, while the 50 kDa protein is suggested herein to be the  
20 processed or mature form.

The catalytic activity of the recombinant protein expressed in the *pShpaMet2* transfected cells was tested by gel shift assay. Cell extracts of

transfected and of mock transfected cells were incubated overnight with heparin (6  $\mu$ g in each reaction) at 37 °C, in the presence of 20 mM phosphate citrate buffer pH 5.4, 1 mM  $\text{CaCl}_2$ , 1 mM DTT and 50 mM NaCl. Reaction mixtures were then separated on a 10 % polyacrylamide gel. The catalytic activity of the recombinant heparanase was clearly demonstrated by a faster migration of the heparin molecules incubated with the transfected cell extract as compared to the control. Faster migration indicates the disappearance of high molecular weight heparin molecules and the generation of low molecular weight degradation products.

## EXAMPLE 9

### *Chromosomal localization of the hpa gene*

Chromosomal mapping of the *hpa* gene was performed utilizing a panel of monochromosomal human/CHO and human/mouse somatic cell hybrids, obtained from the UK HGMP Resource Center (Cambridge, England).

40 ng of each of the somatic cell hybrid DNA samples were subjected to PCR amplification using the *hpa* primers: hpu565 5'-AGCTCTGTAGATGTGC TATACAC-3', SEQ ID NO:22, corresponding to nucleotides 564-586 of SEQ ID NO:9 and an antisense primer hpl171 5'-GCATCTTAGCCGTCTTTCTTCG-3', SEQ ID NO:23, corresponding to nucleotides 897-876 of SEQ ID NO:9.

The PCR program was as follows: a hot start of 94 °C – 3 minutes, followed by 7 cycles of 94 °C – 45 seconds, 66 °C – 1 minute, 68 °C – 5 minutes, followed by 30 cycles of 94 °C – 45 seconds, 62 °C – 1 minute, 68 °C – 5 minutes, and a 10 minutes final extension at 72 °C.

5        The reactions were performed with Expand long PCR (Boehringer Mannheim). The resulting amplification products were analyzed using agarose gel electrophoresis. As demonstrated in Figure 14, a single band of approximately 2.8 Kb was obtained from chromosome 4, as well as from the control human genomic DNA. A 2.8 kb amplification product is expected  
10    based on amplification of the genomic *hpa* clone (data not shown). No amplification products were obtained neither in the control DNA samples of hamster and mouse nor in somatic hybrids of other human chromosome.

## EXAMPLE 10

### 15        *Human genomic clone encoding heparanase*

Five plaques were isolated following screening of a human genomic library and were designated L3-1, L5-1, L8-1, L10-1 and L6-1. The phage DNAs were analyzed by Southern hybridization and by PCR with *hpa* specific and vector specific primers. Southern analysis was performed with  
20    three fragments of *hpa* cDNA: a *PvuII-BamHI* fragment (nucleotides 32-450, SEQ ID NO:9), a *BamHI-NdeI* fragment (nucleotides 451-1102,

SEQ ID NO:9) and an *NdeI-XhoI* fragment (nucleotides 1103-1721, SEQ ID NO:9).

Following Southern analysis, phages L3, L6, L8 were selected for further analysis. A scheme of the genomic region and the relative position of the three phage clones is depicted in Figure 15. A 2 kb DNA fragment containing the gap between phages L6 and L3 was PCR amplified from human genomic DNA with two gene specific primers GHpuL3 and GHplL6. The PCR product was cloned into the plasmid vector pGEM-T-easy (Promega).

Large scale DNA sequencing of the three Lambda clones and the amplified fragment was performed with Lambda purified DNA by primer walking. A nucleotide sequence of 44,898 bp was analyzed (Figure 16, SEQ ID NO:42). Comparison of the genomic sequence with that of *hpa* cDNA revealed 12 exons separated by 11 introns (Figures 15 and 16). The genomic organization of the *hpa* gene is depicted in Figure 15 (top). The sequence include the coding region from the first ATG to the stop codon which spans 39,113 nucleotides, 2742 nucleotides upstream of the first ATG and 3043 nucleotides downstream of the stop codon. Splice site consensus sequences were identified at exon/intron junctions.



**EXAMPLE 11*****Alternative splicing***

Several minor RT-PCR products were obtained from various cell types, following amplification with *hpa* specific primers. Each one found to contain a deletion of one or two exons. Some of these PCR products contain ORFs, which encode potential shorter proteins.

Table 1 below summarizes the alternative spliced products isolated from various cell lines.

Fragments of similar sizes were obtained following amplification with two cell lines, placenta and platelets.

Cell type	Nucleotides deleted	Exons deleted	ORF
Platelets	1047-1267	8, 9	+
Platelets	1154-1267	9	-
Platelets	289-435, 562-735	2, 4	-
Sk-hep1, platelets, Zr75	562-735	4	+
Sk-hep1 (hepatoma)	561-904	4, 5	-
Zr75 (breast carcinoma)	96-203	1 (partial)	+

**EXAMPLE 12*****Mouse and rat hpa***

EST databases were screened for sequences homologous to the *hpa* gene. Three mouse EST's were identified (accession No. Aa177901, from mouse spleen, Aa067997 from mouse skin, Aa47943 from mouse embryo), assembled into a 824 bp cDNA fragment which contains a partial open reading frame (lacking a 5' end) of 629 bp and a 3' untranslated region of 195 bp (SEQ ID NO:12). As shown in Figure 13, the coding region is 80 %

similar to the 3' end of the *hpa* cDNA sequence. These EST's are probably cDNA fragments of the mouse *hpa* homolog that encodes for the mouse heparanase.

Searching for consensus protein domains revealed an amino terminal  
5 homology between the heparanase and several precursor proteins such as Procollagen Alpha 1 precursor, Tyrosine-protein kinase-RYK, Fibulin-1, Insulin-like growth factor binding protein and several others. The amino terminus is highly hydrophobic and contains a potential trans-membrane domain. The homology to known signal peptide sequences suggests that it  
10 could function as a signal peptide for protein localization.

The amino acid sequence of human heparanase was used to search for homologous sequences in the DNA and protein databases. Several human EST's were identified, as well as mouse sequences highly homologous to human heparanase. The following mouse EST's were  
15 identified AA177901, AA674378, AA67997, AA047943, AA690179, AI122034, all sharing an identical sequence and correspond to amino acids 336-543 of the human heparanase sequence. The entire mouse heparanase cDNA was cloned, based on the nucleotide sequence of the mouse EST's. PCR primers were designed and a Marathon RACE was performed using a  
20 Marathon cDNA library from 15 days mouse embryo (Clontech) and from BL6 mouse melanoma cell line. The mouse *hpa* homologous cDNA was isolated following several amplification steps. A 1.1 kb fragment was

amplified from mouse embryo Marathon cDNA library. The first cycle of amplification was performed with primers mhpl773 and Ap1 and the second cycle with primers mhpl736 and AP2. A 1.1 kb fragment was then amplified from BL6 Marathon cDNA library. The first cycle of  
5 amplification was performed with the primers mhpl152 and Ap1, and the second with mhpl83 and AP2. The combined sequence was homologous to nucleotides 157 - 1702 of the human *hpa* cDNA, which encode amino acids 33-543. The 5' end of the mouse *hpa* gene was isolated from a mouse genomic DNA library using the Genome Walker kit (Clontech). An 0.9 kb  
10 fragment was amplified from a *Dra*I digested Genome walker DNA library. The first cycle of amplification was performed with primers mhpl114 and Ap1 and the second with primers mhpl103 and AP2. The assembled sequence (SEQ ID NOs:43, 45) is 2396 nucleotides long. It contains an open reading frame of 1605 nucleotides, which encode a polypeptide of 535  
15 amino acids (SEQ ID NOs:44, 45), 196 nucleotides of 3' untranslated region (UTR), and anupstream sequence which includes the promoter region and the 5'-UTR of the mouse *hpa* cDNA.. According to two promoter predicting programs TSSW and TSSG, the transcription start site is localized to nucleotide 431 of SEQ ID NOs:43, 45, 163 nucleotides  
20 upstream of the first ATG codon. The 431 upstream genomic sequence contains the promoter region. A TATA box is predicted at position 394 of SEQ ID NOs:43, 45. The mouse and the human *hpa* genes share an

average homology of 78 % between the nucleotide sequences and 81 % similarity between the deduced amino acid sequences.

Search for *hpa* homologous sequences, using the Blast 2.0 server revealed two EST's from rat: AI060284 (385 nucleotides, SEQ ID NO:46) which is homologous to the amino terminus (68 % similarity to amino acids 12-136) of human heparanase and AI237828 (541 nucleotides, SEQ ID NO:47) which is homologous to the carboxyl terminus (81 % similarity to amino acids 500-543) of human heparanase, and contains a 3'-UTR. A comparison between the human heparanase and the mouse and rat homologous sequences is demonstrated in Figure 17.

### EXAMPLE 13

#### *Prediction of heparanase active site*

Homology search of heparanase amino acid sequence against the DNA and the protein databases revealed no significant homologies. The protein secondary structure as predicted by the PHD program consists of alternating alpha helices and beta sheets. The fold recognition server of UCLA predicted alpha/beta barrel structure, with under-threshold confidence.

Five of 15 proteins, which were predicted to have most similar folds, were glycosyl hydrolases from various organisms: 1xyza – xylanase from *Clostridium Thermocellum*, 1pbga – 6-phospho-beta- $\delta$ -galactosidase from

Lactococcus Lactis, 1amy – alpha-amylase from Barley, 1ecea – endocellulase from Acidothermus Cellulolyticus and 1qbc – hexosaminidase alpha chain, glycosyl hydrolase.

Protein homology search using the bioaccelerator pulled out several  
 5 proteins, including glycosyl hydrolyses such as beta-fructofuranosidase from *Vicia faba* (broad bean) and from potato, lactase phlorizin hydrolase from human, xylanases from *Clostridium thermocellum* and from *Streptomyces halstedii* and cellulase from *Clostridium thermocellum*. Blocks 9.3 database pulled out the active site of glycosyl hydrolases family  
 10 five, which includes cellulases from various bacteria and fungi. Similar active site motif is shared by several lysosomal acid hydrolases (63) and other glycosyl hydrolases. The common mechanism shared by these enzymes involves two glutamic acid residues, a proton donor and a nucleophile.

15 Despite the lack of an overall homology between the heparanase and other glycosyl hydrolases, the amino acid couple Asp-Glu (NE), which is characteristic of the proton donor of glycosyl hydrolyses of the GH-A clan, was found at positions 224-225 of the human heparanase protein sequence. As in other clan members, this NE couple is located at the end of a  $\beta$  sheet.

20 Considering the relative location of the proton donor and the predicted secondary structure, the glutamic acid that functions as nucleophile is most likely located at position 343, or at position 396.

Identification of the active site and the amino acids directly involved in hydrolysis opens the way for expression of the defined catalytic domain. In addition, it will provide the tools for rational design of enzyme activity either by modification of the microenvironment or catalytic site itself.

5

#### EXAMPLE 14

##### *Expression of hpa antisense in mammalian cell lines*

A mammalian expression vector Hpa2Kepcdna3 was constructed in order to express *hpa* antisense in mammalian cells. *hpa* cDNA (1.7 kb  
10 *Eco*RI fragment) was cloned into the plasmid pCDNA3 in 3'>5' (antisense) orientation. The construct was used to transfect MBT2-T50 and T24P cell lines.  $2 \times 10^5$  cells in 35 mm plates were transfected using the Fugene protocol (Boehringer Mannheim). 48 hours after transfection cells were trypsinized and seeded in six well plates. 24 hours later G418 was added to  
15 initiate selection. The number of colonies per 35 mm plate following 3 weeks:

	Antisense	No insert
T24P	15	60
20 MBT-T50	1	6

The lower number of colonies obtained after transfection with *hpa* antisense, as compared with the control plasmid suggests that the introduction of *hpa* antisense interfere with cell growth. This experiment demonstrates the use of complementary antisense *hpa* DNA sequence to control heparanase expression in cells. This approach may be used to inhibit expression of heparanase *in vivo*, in, for example, cancer cells and in other pathological processes in which heparanase is involved.

## EXAMPLE 15

### *Zoo blot*

*Hpa* cDNA was used as a probe to detect homologous sequences in human DNA and in DNA of various animals. The autoradiogram of the Southern analysis is presented in Figure 18. Several bands were detected in human DNA, which correlated with the accepted pattern according to the genomic *hpa* sequence. Several intense bands were detected in all mammals, while faint bands were detected in chicken. This correlates with the phylogenetic relation between human and the tested animals. The intense bands indicate that *hpa* is conserved among mammals as well as in more genetically distant organisms. The multiple bands patterns suggest that in all animals, like in human, the *hpa* locus occupy large genomic region. Alternatively, the various bands could represent homologous sequences and suggest the existence of a gene family, which can be isolated

based on their homology to the human *hpa* reported herein. This conservation was actually found, between the isolated human *hpa* cDNA and the mouse homologue.

## EXAMPLE 16

### *Characterization of the hpa promoter*

The DNA sequence upstream of the *hpa* first ATG was subjected to computational analysis in order to localize the predicted transcription start site and to identify potential transcription factors binding sites. Recognition of human PolII promoter region and start of transcription were predicted using the TSSW and TSSG programs. Both programs identified a promoter region upstream of the coding region. TSSW pointed at nucleotide 2644 and TSSG at 2635 of SEQ ID NO:42. These two predicted transcription start sites are located 4 and 13 nucleotides upstream of the longest *hpa* cDNA isolated by RACE.

A *hpa* promoter-GFP reporter vector was constructed in order to investigate the regulation of *hpa* transcription. Two constructs were made, containing 1.8 kb and 1.1 kb of the *hpa* promoter region. The reporter vector was transfected into T50-mouse bladder carcinoma cells. Cells transfected with both constructs exhibited green fluorescence, which indicated the promoter activity of the genomic sequence upstream of the *hpa*-coding region. This reporter vector, enables the monitoring of *hpa*



promoter activity, at various conditions and in different cell types and to characterize the factors involved regulation of *hpa* expression.

Although the invention has been described in conjunction with  
5 specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims.

LIST OF REFERENCES

1. Wight, T.N., Kinsella, M.G., and Qwarnstromn, E.E. (1992). The role of proteoglycans in cell adhesion, migration and proliferation. *Curr. Opin. Cell Biol.*, 4, 793-801.
2. Jackson, R.L., Busch, S.J., and Cardin, A.L. (1991). Glycosaminoglycans: Molecular properties, protein interactions and role in physiological processes. *Physiol. Rev.*, 71, 481-539.
3. Wight, T.N. (1989). Cell biology of arterial proteoglycans. *Arteriosclerosis*, 9, 1-20.
4. Kjellen, L., and Lindahl, U. (1991). Proteoglycans: structures and interactions. *Annu. Rev. Biochem.*, 60, 443-475.
5. Ruoslahti, E., and Yamaguchi, Y. (1991). Proteoglycans as modulators of growth factor activities. *Cell*, 64, 867-869.
6. Vlodavsky, I., Eldor, A., Haimovitz-Friedman, A., Matzner, Y., Ishai-Michaeli, R., Levi, E., Bashkin, P., Lider, O., Naparstek, Y., Cohen, I.R., and Fuks, Z. (1992). Expression of heparanase by platelets and

circulating cells of the immune system: Possible involvement in diapedesis and extravasation. *Invasion & Metastasis*, 12, 112-127.

7. Vlodavsky, I., Mohsen, M., Lider, O., Ishai-Michaeli, R., Ekre, H.-P., Svahn, C.M., Vigoda, M., and Peretz, T. (1995). Inhibition of tumor metastasis by heparanase inhibiting species of heparin. *Invasion & Metastasis*, 14, 290-302.

8. Nakajima, M., Irimura, T., and Nicolson, G.L. (1988). Heparanase and tumor metastasis. *J. Cell. Biochem.*, 36, 157-167.

9. Nicolson, G.L. (1988). Organ specificity of tumor metastasis: Role of preferential adhesion, invasion and growth of malignant cells at specific secondary sites. *Cancer Met. Rev.*, 7, 143-188.

10. Liotta, L.A., Rao, C.N., and Barsky, S.H. (1983). Tumor invasion and the extracellular matrix. *Lab. Invest.*, 49, 639-649.

11. Vlodavsky, I., Fuks, Z., Bar-Ner, M., Ariav, Y., and Schirmacher, V. (1983). Lymphoma cell mediated degradation of sulfated proteoglycans in the subendothelial extracellular matrix: Relationship to tumor cell metastasis. *Cancer Res.*, 43, 2704-2711.

p12p. Vlodavsky, I., Ishai-Michaeli, R., Bar-Ner, M., Fridman, R., Horowitz, A.T., Fuks, Z. and Biran, S. (1988). Involvement of heparanase in tumor metastasis and angiogenesis. *Is. J. Med.*, 24, 464-470.

13. Vlodavsky, I., Liu, G.M., and Gospodarowicz, D. (1980). Morphological appearance, growth behavior and migratory activity of human tumor cells maintained on extracellular matrix vs. plastic. *Cell*, 19, 607-616.

14. Gospodarowicz, D., Delgado, D., and Vlodavsky, I. (1980). Permissive effect of the extracellular matrix on cell proliferation in-vitro. *Proc. Natl. Acad. Sci. USA.*, 77, 4094-4098.

15. Bashkin, P., Doctrow, S., Klagsbrun, M., Svahn, C.M., Folkman, J., and Vlodavsky, I. (1989). Basic fibroblast growth factor binds to subendothelial extracellular matrix and is released by heparitinase and heparin-like molecules. *Biochemistry*, 28, 1737-1743.

16. Parish, C.R., Coombe, D.R., Jakobsen, K.B., and Underwood, P.A. (1987). Evidence that sulphated polysaccharides inhibit tumor metastasis by blocking tumor cell-derived heparanase. *Int. J. Cancer*, 40, 511-517.

- 16a. Vlodavsky, I., Hua-Quan Miao., Benezra, M., Lider, O., Bar-Shavit, R., Schmidt, A., and Peretz, T. (1997). Involvement of the extracellular matrix, heparan sulfate proteoglycans and heparan sulfate degrading enzymes in angiogenesis and metastasis. In: Tumor Angiogenesis. Eds. C.E. Lewis, R. Bicknell & N. Ferrara. Oxford University Press, Oxford UK, pp. 125-140.
17. Burgess, W.H., and Maciag, T. (1989). The heparin-binding (fibroblast) growth factor family of proteins. *Annu. Rev. Biochem.*, 58, 575-606.
18. Folkman, J., and Klagsbrun, M. (1987). Angiogenic factors. *Science*, 235, 442-447.
19. Vlodavsky, I., Folkman, J., Sullivan, R., Fridman, R., Ishai-Michaelli, R., Sasse, J., and Klagsbrun, M. (1987). Endothelial cell-derived basic fibroblast growth factor: Synthesis and deposition into subendothelial extracellular matrix. *Proc. Natl. Acad. Sci. USA*, 84, 2292-2296.
20. Folkman, J., Klagsbrun, M., Sasse, J., Wadzinski, M., Ingber, D., and Vlodavsky, I. (1980). A heparin-binding angiogenic protein - basic

fibroblast growth factor - is stored within basement membrane. *Am. J. Pathol.*, 130, 393-400.

21. Cardon-Cardo, C., Vlodavsky, I., Haimovitz-Friedman, A., Hicklin, D., and Fuks, Z. (1990). Expression of basic fibroblast growth factor in normal human tissues. *Lab. Invest.*, 63, 832-840.

22. Ishai-Michaeli, R., Svahn, C.-M., Chajek-Shaul, T., Korner, G., Ekre, H.-P., and Vlodavsky, I. (1992). Importance of size and sulfation of heparin in release of basic fibroblast factor from the vascular endothelium and extracellular matrix. *Biochemistry*, 31, 2080-2088.

23. Ishai-Michaeli, R., Eldor, A., and Vlodavsky, I. (1990). Heparanase activity expressed by platelets, neutrophils and lymphoma cells releases active fibroblast growth factor from extracellular matrix. *Cell Reg.*, 1, 833-842.

24. Vlodavsky, I., Bar-Shavit, R., Ishai-Michaeli, R., Bashkin, P., and Fuks, Z. (1991). Extracellular sequestration and release of fibroblast growth factor: a regulatory mechanism? *Trends Biochem. Sci.*, 16, 268-271.

25. Vlodavsky, I., Bar-Shavit, R., Korner, G., and Fuks, Z. (1993). Extracellular matrix-bound growth factors, enzymes and plasma proteins. In Basement membranes: Cellular and molecular aspects (eds. D.H. Rohrbach and R. Timpl), pp327-343. Academic press Inc., Orlando, Fl.
26. Yayon, A., Klagsbrun, M., Esko, J.D., Leder, P., and Ornitz, D.M. (1991). Cell surface, heparin-like molecules are required for binding of basic fibroblast growth factor to its high affinity receptor. *Cell*, 64, 841-848.
27. Spivak-Kroizman, T., Lemmon, M.A., Dikic, I., Ladbury, J.E., Pinchasi, D., Huang, J., Jaye, M., Crumley, G., Schlessinger, J., and Lax, I. (1994). Heparin-induced oligomerization of FGF molecules is responsible for FGF receptor dimerization, activation, and cell proliferation. *Cell*, 79, 1015-1024.
28. Ornitz, D.M., Herr, A.B., Nilsson, M., West, a., J., Svahn, C.-M., and Waksman, G. (1995). FGF binding and FGF receptor activation by synthetic heparan-derived di- and trisaccharides. *Science*, 268, 432-436.

29. Gitay-Goren, H., Soker, S., Vlodavsky, I., and Neufeld, G. (1992). Cell surface associated heparin-like molecules are required for the binding of vascular endothelial growth factor (VEGF) to its cell surface receptors. *J. Biol. Chem.*, 267, 6093-6098.
30. Lider, O., Baharav, E., Mekori, Y., Miller, T., Naparstek, Y., Vlodavsky, I., and Cohen, I.R. (1989). Suppression of experimental autoimmune diseases and prolongation of allograft survival by treatment of animals with heparinoid inhibitors of T lymphocyte heparanase. *J. Clin. Invest.*, 83, 752-756.
31. Lider, O., Cahalon, L., Gilat, D., HersHKovitz, R., Siegel, D., Margalit, R., Shoseyov, O., and Cohn, I.R. (1995). A disaccharide that inhibits tumor necrosis factor  $\alpha$  is formed from the extracellular matrix by the enzyme heparanase. *Proc. Natl. Acad. Sci. USA.*, 92, 5037-5041.
- 31a. Rapraeger, A., Krufka, A., and Olwin, B.R. (1991). Requirement of heparan sulfate for bFGF-mediated fibroblast growth and myoblast differentiation. *Science*, 252, 1705-1708.
32. Eisenberg, S., Sehayek, E., Olivecrona, T., and Vlodavsky, I. (1992). Lipoprotein lipase enhances binding of lipoproteins to heparan



sulfate on cell surfaces and extracellular matrix. *J. Clin. Invest.*, 90, 2013-2021.

þ33þ. Shieh, M-T., Wundunn, D., Montgomery, R.I., Esko, J.D., and Spear, P.G. J. (1992). Cell surface receptors for herpes simplex virus are heparan sulfate proteoglycans. *J Cell Biol.*, 116, 1273-1281.

33a. Chen, Y., Maguire, T., Hileman, R.E., Fromm, J.R., Esko, J.D., Linhardt, R.J., and Marks, R.M. (1997). Dengue virus infectivity depends on envelope protein binding to target cell heparan sulfate. *Nature Medicine* 3, 866-871.

33b. Putnak, J.R., Kanesa-Thasan, N., and Innis, B.L. (1997). A putative cellular receptor for dengue viruses. *Nature Medicine* 3, 828-829.

þ34þ. Narindrasorasak, S., Lowery, D., Gonzalez-DeWhitt, P., Poorman, R.A., Greenberg, B., Kisilevsky, R. (1991). High affinity interactions between the Alzheimer's beta-amyloid precursor protein and the basement membrane form of theparan sulfate proteoglycan. *J. Biol. Chem.*, 266, 12878-83.

35. Ross, R. (1993). The pathogenesis of atherosclerosis: a perspective for the 1990s. *Nature (Lond.)*, 362:801-809.
36. Zhong-Sheng, J., Walter, J., Brecht, R., Miranda, D., Mahmood Hussain, M., Innerarity, T.L. and Mahley, W.R. (1993). Role of heparan sulfate proteoglycans in the binding and uptake of apolipoprotein E-enriched remnant lipoproteins by cultured cells. *J. Biol. Chem.*, 268, 10160-10167.
37. Ernst, S., Langer, R., Cooney, Ch.L., and Sasisekharan, R. (1995). Enzymatic degradation of glycosaminoglycans. *Critical Reviews in Biochemistry and Molecular Biology*, 30(5), 387-444.
38. Gospodarowicz, D., Mescher, AL., Birdwell, CR. (1977). Stimulation of corneal endothelial cell proliferation in vitro by fibroblast and epidermal growth factors. *Exp Eye Res* 25, 75-89.
39. Haimovitz-Friedman, A., Falcone, D.J., Eldor, A., Schirmacher, V., Vlodavsky, I., and Fuks, Z. (1991) Activation of platelet heparitinase by tumor cell-derived factors. *Blood*, 78, 789-796.

- 39a. Savitsky, K., Platzer, M., Uziel, T., Gilad, S., Sartiel, A., Rosental, A., Elroy-Stein, O., Siloh, Y. and Rotman, G. (1997). Ataxia-telangiectasia: structural diversity of untranslated sequences suggests complex post-translational regulation of ATM gene expression. *Nucleic Acids Res.* 25(9), 1678-1684.
40. Bar-Ner, M., Eldor, A., Wasserman, L., Matzner, Y., and Vlodavsky, I. (1987). Inhibition of heparanase mediated degradation of extracellular matrix heparan sulfate by modified and non-anticoagulant heparin species. *Blood*, 70, 551-557.
41. Goshen, R., Hochberg, A., Korner, G., Levi, E., Ishai-Michaeli, R., Elkin, M., de Grot, N., and Vlodavsky, I. (1996). Purification and characterization of placental heparanase and its expression by cultured cytotrophoblasts. *Mol. Human Reprod.*, 2, 679-684.
42. Korb M., Ke Y. and Johnson L.F. (1993) Stimulation of gene expression by introns: conversion of an inhibitory intron to a stimulatory intron by alteration of the splice donor sequence. *Nucleic Acids Res.*, 21(25):5901-8.

43. Zheng B., Qiu X.Y., Tan M., Xing Y.N., Lo D., Xue J.L. and Qiu X.F. (1997) Increment of hFIX expression with endogenous intron 1 in vitro. *Cell Res.*, 7(1):21-29.
44. Kurachi S., Hitomi Y., Furukawa M. and Kurachi K. (1995) Role of intron I in expression of the human factor IX gene. *J. Biol. Chem.* 10, 270(10):5276-5281.
45. Shekhar P.V. and Miller F.R. (1994-5) Correlation of differences in modulation of ras expression with metastatic competence of mouse mammary tumor subpopulations. *Invasion Metastasis*, 14(1-6):27-37.
46. Zhou G., Garofalo S., Mukhopadhyay K., Lefebvre V., Smith C.N., Eberspacher H. and de Crombrughe B. (1995) A 182 bp fragment of the mouse pro alpha 1(II) collagen gene is sufficient to direct chondrocyte expression in transgenic mice. *J. Cell Sci.*, 108 ( Pt 12):3677-3684.
47. Hormuzdi S.G., Penttinen R., Jaenisch R. and Bornstein P. (1998) A gene-targeting approach identifies a function for the first intron in expression of the alpha1(I) collagen gene. *Mol. Cell*, 18(6):3368-3375.

48. Kang Y.K., Lee C.S., Chung A.S. and Lee K.K. (1998) Prolactin-inducible enhancer activity of the first intron of the bovine beta-casein gene. *Mol. Cells*, 30;8(3):259-265.
49. Chow Y.H., O'Brodovich H., Plumb J., Wen Y., Sohn K.J., Lu Z., Zhang F., Lukacs G.L., Tanswell A.K., Hui C.C., Buchwald M. and Hu J. (1997) Development of an epithelium-specific expression cassette with human DNA regulatory elements for transgene expression in lung airways. *Proc. Natl. Acad. Sci. USA*, 23;94(26):14695-14700.
50. Gottschalk U. and Chan S. (1998) Somatic gene therapy. Present situation and future perspective. *Arzneimittelforschung*, 48(11):1111-1120.
51. Ye S., Cole-Strauss A.C., Frank B. and Kmiec E.B. (1998) Targeted gene correction: a new strategy for molecular medicine. *Mol. Med. Today*, 4(10):431-437.
52. Lai L., and Lien Y. (1999) Homologous recombination based gene therapy. *Exp. Nephrol.*, 7(1):11-14.

53. Yazaki N., Fujita H., Ohta M., Kawasaki T. and Itoh N. (1993) The structure and expression of the FGF receptor-1 mRNA isoforms in rat tissues. *Biochim. Biophys. Acta.*, 20;1172(1-2):37-42.
54. Le Fur N., Kelsall S.R., Silvers W.K. and Mintz B. (1997) Selective increase in specific alternative splice variants of tyrosinase in murine melanomas: a projected basis for immunotherapy. *Proc. Natl. Acad. Sci. USA*, 13;94(10):5332-5337.
55. Miyake H., Okamoto I., Hara I., Gohji K., Yamanaka K., Arakawa S., Kamidono S. and Saya H. (1998) Highly specific and sensitive detection of malignancy in urine samples from patients with urothelial cancer by CD44v8-10/CD44v10 competitive RT-PCR. *Int. J. Cancer*, 18;79(6):560-564.
56. Guriec N., Marcellin L., Gairard B., Calderoli H., Wilk A., Renaud R., Bergerat J.P. and Oberling F. (1996) CD44 exon 6 expression as a possible early prognostic factor in primary node negative breast carcinoma. *Clin. Exp. Metastasis*, 14(5):434-439.

57. Gewirtz A.M., Sokol D.L. and Ratajczak M.Z. (1998) Nucleic acid therapeutics: state of the art and future prospects. *Blood*, 1;92(3):712-736.
58. Hida K., Shindoh M., Yasuda M., Hanzawa M., Funaoka K., Kohgo T., Amemiya A., Totsuka Y., Yoshida K. and Fujinaga K (1997) Antisense E1AF transfection restrains oral cancer invasion by reducing matrix metalloproteinase activities. *Am. J. Pathol.* 150(6):2125-2132.
59. Shastry B.S. (1998) Gene disruption in mice: models of development and disease. *Mol. Cell. Biochem.* 1998 Apr;181(1-2):163-179.
60. Carpentier A.F., Rosenfeld M.R., Delattre J.Y., Whalen R.G., Posner J.B. and Dalmau J. (1998) DNA vaccination with HuD inhibits growth of a neuroblastoma in mice. *Clin.Cancer Res.*, 4(11):2819-2824.
61. Lai W.C. and Bennett M. (1998) DNA vaccines. *Crit. Rev. Immunol.*, 18(5):449-484.
62. Welch P.J., Barber J.R., and Wong-Staal F. (1998) Expression of ribozymes in gene transfer systems to modulate target RNA levels. *Curr. Opin. Biotechnol.*, 9(5):486-496.

63. Durand P., Lehn P., Callebaunt I., Fabrega S., Henrissat B. and Mornon J.P. (1997) Active-site motifs of lysosomal acid hydrolyses: invariant features of clan GH-A glycosyl hydrolases deduced from hydrophobic cluster analysis. *Glycobiology*, 7(2):277-284.
64. Thuong and Helene (1993) Sequence specific recognition and modification of double helical DNA by oligonucleotides *Angev. Chem. Int. Ed. Engl.* 32:666
65. Dash P., Lotan I., Knapp M., Kandel E.R. and Goelet P. (1987) Selective elimination of mRNAs in vivo: complementary oligodeoxynucleotides promote RNA degradation by an RNase H-like activity. *Proc. Natl. Acad. Sci. USA*, 84:7896.
66. Chiang M.Y., Chan H., Zounes M.A., Freier S.M., Lima W.F. and Bennett C.F. (1991) Antisense oligonucleotides inhibit intercellular adhesion molecule 1 expression by two distinct mechanisms. *J. Biol. Chem.* 266:18162-71.



67. Paterson B.M, Roberts B.E and Kuff EL . (1977) Structural gene identification and mapping by DNA-mRNA hybrid-arrested cell-free translation. *Proc. Natl. Acad. Sci. USA*, 74:4370.
68. Cohen (1992) Oligonucleotide therapeutics. *Trends in Biotechnology*, 10:87.
69. Szczylik et al (1991) Selective inhibition of leukemia cell proliferation by BCR-ABL antisense oligodeoxynucleotides. *Science* 253:562.
70. Calabretta et al. (1991) Normal and leukemic hematopoietic cell manifest differential sensitivity to inhibitory effects of c-myc antisense oligodeoxynucleotides: an in vitro study relevant to bone marrow purging. *Proc. Natl. Acad. Sci. USA* 88:2351.
71. Heikhila et al. (1987) A c-myc antisense oligodeoxynucleotide inhibits entry into S phase but not progress from G(0) to G(1). *Nature*, 328:445.
72. Reed et al. (1990) Antisense mediated inhibition of BCL2 protooncogene expression and leukemic cell growth and survival: comparison

of phosphodiester and phosphorothioate oligodeoxynucleotides. Cancer Res. 50:6565.

73. Burch and Mahan (1991) Oligodeoxynucleotides antisense to the interleukin I receptor m RNA block the effects of interleukin I in cultured murine and human fibroblasts and in mice. J. Clin. Invest. 88:1190.

74. Agrawal (1992) Antisense oligonucleotides as antiviral agents. TIBTECH 10:152.

75. Uhlmann et al. (1990) Chem. Rev. 90:544.

76. Cook (1991) Medicinal chemistry of antisense oligonucleotides - future opportunities. Anti-Cancer Drug Design 6:585.

77. Biotechnology research news (1993) Can DNA mimics improve on the real thing? Science 262:1647.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Iris Pecker, Israel Vlodavsky and Elena Feinstein
- (ii) TITLE OF INVENTION: POLYNUCLEOTIDE ENCODING A POLYPEPTIDE HAVING HEPARANASE ACTIVITY AND EXPRESSION OF SAME IN GENETICALLY MODIFIED CELLS
- (iii) NUMBER OF SEQUENCES: 47
- (iv) CORRESPONDENCE ADDRESS:
- (A) ADDRESSEE: Mark M. Friedman c/o Anthony Castorina
- (B) STREET: 2001 Jefferson Davis Highway, Suite 207
- (C) CITY: Arlington
- (D) STATE: Virginia
- (E) COUNTRY: United States of America
- (F) ZIP: 22202
- (v) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: 1.44 megabyte, 3.5" microdisk
- (B) COMPUTER: Twinhead\* Slimnote-890TX
- (C) OPERATING SYSTEM: MS DOS version 6.2, Windows version 3.11
- (D) SOFTWARE: Word for Windows version 2.0 converted to an ASCII file
- (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER:
- (B) FILING DATE:
- (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: 08/922,170
- (B) FILING DATE: 2 SEP 1997
- (A) APPLICATION NUMBER: 09/109,386
- (B) FILING DATE: 10 JUL 1998
- (A) APPLICATION NUMBER: PCT/US98/17954
- (B) FILING DATE: 31 AUG 1998
- (viii) ATTORNEY/AGENT INFORMATION:
- (A) NAME: Friedmam, Mark M.
- (B) REGISTRATION NUMBER: 33,883
- (C) REFERENCE/DOCKET NUMBER: 910/14
- (ix) TELECOMMUNICATION INFORMATION:
- (A) TELEPHONE: 972-3-5625553
- (B) TELEFAX: 972-3-5625554
- (C) TELEX:

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 27
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:  
CCATCCTAAT ACGACTCACT ATAGGGC 27

## (2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GTAGTGATGC CATGTAAC TG AATC 24

- (2) INFORMATION FOR SEQ ID NO:3:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 23
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:  
ACTCACTATA GGGCTCGAGC GGC 23
- (2) INFORMATION FOR SEQ ID NO:4:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 22
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:  
GCATCTTAGC CGTCTTCTT CG 22
- (2) INFORMATION FOR SEQ ID NO:5:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 15
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:  
TTTTTTTTTT TTTT 15
- (2) INFORMATION FOR SEQ ID NO:6:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 23
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:  
TTCGATCCCA AGAAGGAATC AAC 23
- (2) INFORMATION FOR SEQ ID NO:7:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:  
GTAGTGATGC CATGTAAC TG AATC 24
- (2) INFORMATION FOR SEQ ID NO:8:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 9
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:  
Tyr Gly Pro Asp Val Gly Gln Pro Arg

## (2) INFORMATION FOR SEQ ID NO:9:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1721  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CTAGAGCTTT CGACTCTCCG CTGCGCGGCA GCTGGCGGGG GGAGCAGCCA GGTGAGCCCA 60  
 AGATGCTGCT GCGCTCGAAG CCTGCGCTGC CGCCGCCGCT GATGCTGCTG CTCCTGGGGC 120  
 CGCTGGGTCC CCTCTCCCTT GCGGCCCTGC CCCGACCTGC GCAAGCACAG GACGTCGTGG 180  
 ACCTGGACTT cTTCACCCAG GAGCCGCTGC ACCTGGTGAG CCCCTCGTTC CTGTCCGTCA 240  
 CCATTGACGC CAACCTGGCC ACGGACCCGC GGTTCCTCAT CCTCCTGGGT TCTCCAAAGC 300  
 TTCGTACCTT GGCCAGAGGC TTGTCTCCTG CGTACCTGAG GTTTGGTGGC ACCAAGACAG 360  
 ACTTCCTAAT TTTGATCCC AAGAAGGAAT CAACCTTTGA AGAGAGAAGT TACTGGCAAT 420  
 CTCAAGTCAA CCAGGATATT TGCAAATATG GATCCATCCC TCCTGATGTG GAGGAGAAGT 480  
 TACGGTTGGA ATGGCCCTAC CAGGAGCAAT TGCTACTCCG AGAACACTAC CAGAAAAAGT 540  
 TCAAGAACAG CACCTACTCA AGAAGCTCTG TAGATGTGCT ATACACTTTT GCAAACCTGCT 600  
 CAGGACTGGA CTTGATCTTT GGCCTAAATG CGTTATTAAG AACAGCAGAT TTGCAGTGGA 660  
 ACAGTTCTAA TGCTCAGTTT CTCTGGACT ACTGCTCTTC CAAGGGGTAT AACATTTCTT 720  
 GGGAACTAGG CAATGAACCT AACAGTTTCC TTAAGAAGGC TGATATTTTC ATCAATGGGT 780  
 CGCAGTTAGG AGAAGATTAT ATTCAATTGC ATAAACTTCT AAGAAAGTCC ACCTTCAAAA 840  
 ATGCAAAACT CTATGGTCCT GATGTTGGTC AGCCTCGAAG AAAGACGGCT AAGATGCTGA 900  
 AGAGCTTCCT GAAGGCTGGT GGAGAAGTGA TTGATTCACT TACATGGCAT CACTACTATT 960  
 TGAATGGACG GACTGCTACC AGGGAAGATT TTCTAAACCC TGATGTATTG GACATTTTAA 1020  
 TTTCATCTGT GCAAAAAGTT TTCCAGGTGG TTGAGAGCAC CAGGCCTGGC AAGAAGGTCT 1080  
 GGTTAGGAGA AACAAAGCTT GCATATGGAG GCGGAGCGCC CTTGCTATCC GACACCTTTG 1140  
 CAGCTGGCTT TATGTGGCTG GATAAATTGG GCCTGTCAGC CCGAATGGGA ATAGAAGTGG 1200  
 TGATGAGGCA AGTATTCTTT GGAGCAGGAA ACTACCATT AGTGATGAA AACTTCGATC 1260  
 CTTTACCTGA TTATTGGCTA TCTCTTCTGT TCAAGAAATT GGTGGGCACC AAGGTGTTAA 1320  
 TGGAAGCGT GCAAGGTTCA AAGAGAAGGA AGCTTCGAGT ATACCTTCAT TGCACAAACA 1380  
 CTGACAAATC AAGGTATAAA GAAGGAGATT TAACTCTGTA TGCCATAAAC CTCCATAACG 1440  
 TCACCAAGTA CTTGCGGTTA CCTATCCTT TTTCTAACAA GCAAGTGGAT AAATACCTTC 1500  
 TAAGACCTTT GGGACCTCAT GGATTACTTT CCAAATCTGT CCAACTCAAT GGTCTAACTC 1560  
 TAAAGATGGT GGATGATCAA ACCTTGCCAC CTTTAATGGA AAAACCTCTC CGGCCAGGAA 1620  
 GTTCACTGGG CTTGCCAGCT TTCTCATATA GTTTTCTTGT GATAAGAAAT GCCAAAGTTG 1680  
 CTGCTTGCAT CTGAAATATA AATATACTAG TCCTGACACT G 1721

## (2) INFORMATION FOR SEQ ID NO:10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 543  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Leu Leu Arg Ser Lys Pro Ala Leu Pro Pro Pro Leu Met Leu Leu  
 5 10 15

Leu Leu Gly Pro Leu Gly Pro Leu Ser Pro Gly Ala Leu Pro Arg Pro  
 20 25 30

Ala Gln Ala Gln Asp Val Val Asp Leu Asp Phe Phe Thr Gln Glu Pro  
 35 40 45

Leu His Leu Val Ser Pro Ser Phe Leu Ser Val Thr Ile Asp Ala Asn  
 50 55 60

Leu Ala Thr Asp Pro Arg Phe Leu Ile Leu Leu Gly Ser Pro Lys Leu

## 130

65	70	75	80
Arg Thr Leu Ala Arg Gly Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly	85	90	95
Thr Lys Thr Asp Phe Leu Ile Phe Asp Pro Lys Lys Glu Ser Thr Phe	100	105	110
Glu Glu Arg Ser Tyr Trp Gln Ser Gln Val Asn Gln Asp Ile Cys Lys	115	120	125
Tyr Gly Ser Ile Pro Pro Asp Val Glu Glu Lys Leu Arg Leu Glu Trp	130	135	140
Pro Tyr Gln Glu Gln Leu Leu Arg Glu His Tyr Gln Lys Lys Phe	145	150	155
Lys Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Val Leu Tyr Thr Phe	165	170	175
Ala Asn Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu Asn Ala Leu Leu	180	185	190
Arg Thr Ala Asp Leu Gln Trp Asn Ser Ser Asn Ala Gln Leu Leu Leu	195	200	205
Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile Ser Trp Glu Leu Gly Asn	210	215	220
Glu Pro Asn Ser Phe Leu Lys Lys Ala Asp Ile Phe Ile Asn Gly Ser	225	230	235
Gln Leu Gly Glu Asp Tyr Ile Gln Leu His Lys Leu Leu Arg Lys Ser	245	250	255
Thr Phe Lys Asn Ala Lys Leu Tyr Gly Pro Asp Val Gly Gln Pro Arg	260	265	270
Arg Lys Thr Ala Lys Met Leu Lys Ser Phe Leu Lys Ala Gly Gly Glu	275	280	285
Val Ile Asp Ser Val Thr Trp His His Tyr Tyr Leu Asn Gly Arg Thr	290	295	300
Ala Thr Arg Glu Asp Phe Leu Asn Pro Asp Val Leu Asp Ile Phe Ile	305	310	315
Ser Ser Val Gln Lys Val Phe Gln Val Val Glu Ser Thr Arg Pro Gly	325	330	335
Lys Lys Val Trp Leu Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala	340	345	350
Pro Leu Leu Ser Asp Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys	355	360	365
Leu Gly Leu Ser Ala Arg Met Gly Ile Glu Val Val Met Arg Gln Val			

370	375	380
Phe Phe Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe Asp Pro		
385	390	395 400
Leu Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu Val Gly Thr		
	405	410 415
Lys Val Leu Met Ala Ser Val Gln Gly Ser Lys Arg Arg Lys Leu Arg		
	420	425 430
Val Tyr Leu His Cys Thr Asn Thr Asp Asn Pro Arg Tyr Lys Glu Gly		
	435	440 445
Asp Leu Thr Leu Tyr Ala Ile Asn Leu His Asn Val Thr Lys Tyr Leu		
	450	455 460
Arg Leu Pro Tyr Pro Phe Ser Asn Lys Gln Val Asp Lys Tyr Leu Leu		
	465	470 475 480
Arg Pro Leu Gly Pro His Gly Leu Leu Ser Lys Ser Val Gln Leu Asn		
	485	490 495
Gly Leu Thr Leu Lys Met Val Asp Asp Gln Thr Leu Pro Pro Leu Met		
	500	505 510
Glu Lys Pro Leu Arg Pro Gly Ser Ser Leu Gly Leu Pro Ala Phe Ser		
	515	520 525
Tyr Ser Phe Phe Val Ile Arg Asn Ala Lys Val Ala Ala Cys Ile		
	530	535 540 543

## (2) INFORMATION FOR SEQ ID NO:11:

## (i) SEQUENCE CHARACTERISTICS:

(A)	LENGTH:	1721
(B)	TYPE:	nucleic acid
(C)	STRANDEDNESS:	double
(D)	TOPOLOGY:	linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CT AGA GCT TTC GAC	14
TCT CCG CTG CGC GGC AGC TGG CGG GGG GAG CAG CCA GGT GAG CCC AAG	62
ATG CTG CTG CGC TCG AAG CCT GCG CTG CCG CCG CCG CTG ATG CTG CTG	110
Met Leu Leu Arg Ser Lys Pro Ala Leu Pro Pro Pro Leu Met Leu Leu	
5 10 15	
CTC CTG GGG CCG CTG GGT CCC CTC TCC CCT GGC GCC CTG CCC CGA CCT	158
Leu Leu Gly Pro Leu Gly Pro Leu Ser Pro Gly Ala Leu Pro Arg Pro	
20 25 30	
GCG CAA GCA CAG GAC GTC GTG GAC CTG GAC TTC TTC ACC CAG GAG CCG	206
Ala Gln Ala Gln Asp Val Val Asp Leu Asp Phe Phe Thr Gln Glu Pro	
35 40 45	
CTG CAC CTG GTG AGC CCC TCG TTC CTG TCC GTC ACC ATT GAC GCC AAC	254

Leu	His	Leu	Val	Ser	Pro	Ser	Phe	Leu	Ser	Val	Thr	Ile	Asp	Ala	Asn	
50						55					60					
CTG	GCC	ACG	GAC	CCG	CGG	TTC	CTC	ATC	CTC	CTG	GGT	TCT	CCA	AAG	CTT	302
Leu	Ala	Thr	Asp	Pro	Arg	Phe	Leu	Ile	Leu	Leu	Gly	Ser	Pro	Lys	Leu	
65					70					75					80	
CGT	ACC	TTG	GCC	AGA	GGC	TTG	TCT	CCT	GCG	TAC	CTG	AGG	TTT	GGT	GGC	350
Arg	Thr	Leu	Ala	Arg	Gly	Leu	Ser	Pro	Ala	Tyr	Leu	Arg	Phe	Gly	Gly	
				85					90					95		
ACC	AAG	ACA	GAC	TTC	CTA	ATT	TTC	GAT	CCC	AAG	AAG	GAA	TCA	ACC	TTT	398
Thr	Lys	Thr	Asp	Phe	Leu	Ile	Phe	Asp	Pro	Lys	Lys	Glu	Ser	Thr	Phe	
			100					105					110			
GAA	GAG	AGA	AGT	TAC	TGG	CAA	TCT	CAA	GTC	AAC	CAG	GAT	ATT	TGC	AAA	446
Glu	Glu	Arg	Ser	Tyr	Trp	Gln	Ser	Gln	Val	Asn	Gln	Asp	Ile	Cys	Lys	
		115				120					125					
TAT	GGA	TCC	ATC	CCT	CCT	GAT	GTG	GAG	GAG	AAG	TTA	CGG	TTG	GAA	TGG	494
Tyr	Gly	Ser	Ile	Pro	Pro	Asp	Val	Glu	Glu	Lys	Leu	Arg	Leu	Glu	Trp	
	130					135					140					
CCC	TAC	CAG	GAG	CAA	TTG	CTA	CTC	CGA	GAA	CAC	TAC	CAG	AAA	AAG	TTC	542
Pro	Tyr	Gln	Glu	Gln	Leu	Leu	Leu	Arg	Glu	His	Tyr	Gln	Lys	Lys	Phe	
145					150					155					160	
AAG	AAC	AGC	ACC	TAC	TCA	AGA	AGC	TCT	GTA	GAT	GTG	CTA	TAC	ACT	TTT	590
Lys	Asn	Ser	Thr	Tyr	Ser	Arg	Ser	Ser	Val	Asp	Val	Leu	Tyr	Thr	Phe	
				165					170					175		
GCA	AAC	TGC	TCA	GGA	CTG	GAC	TTG	ATC	TTT	GGC	CTA	AAT	GCG	TTA	TTA	638
Ala	Asn	Cys	Ser	Gly	Leu	Asp	Leu	Ile	Phe	Gly	Leu	Asn	Ala	Leu	Leu	
			180					185					190			
AGA	ACA	GCA	GAT	TTG	CAG	TGG	AAC	AGT	TCT	AAT	GCT	CAG	TTG	CTC	CTG	686
Arg	Thr	Ala	Asp	Leu	Gln	Trp	Asn	Ser	Ser	Asn	Ala	Gln	Leu	Leu	Leu	
		195					200					205				
GAC	TAC	TGC	TCT	TCC	AAG	GGG	TAT	AAC	ATT	TCT	TGG	GAA	CTA	GGC	AAT	734
Asp	Tyr	Cys	Ser	Ser	Lys	Gly	Tyr	Asn	Ile	Ser	Trp	Glu	Leu	Gly	Asn	
	210					215					220					
GAA	CCT	AAC	AGT	TTC	CTT	AAG	AAG	GCT	GAT	ATT	TTC	ATC	AAT	GGG	TCG	782
Glu	Pro	Asn	Ser	Phe	Leu	Lys	Lys	Ala	Asp	Ile	Phe	Ile	Asn	Gly	Ser	
225					230					235				240		
CAG	TTA	GGA	GAA	GAT	TAT	ATT	CAA	TTG	CAT	AAA	CTT	CTA	AGA	AAG	TCC	830
Gln	Leu	Gly	Glu	Asp	Tyr	Ile	Gln	Leu	His	Lys	Leu	Leu	Arg	Lys	Ser	
			245					250					255			
ACC	TTC	AAA	AAT	GCA	AAA	CTC	TAT	GGT	CCT	GAT	GTT	GGT	CAG	CCT	CGA	878
Thr	Phe	Lys	Asn	Ala	Lys	Leu	Tyr	Gly	Pro	Asp	Val	Gly	Gln	Pro	Arg	
			260					265					270			
AGA	AAG	ACG	GCT	AAG	ATG	CTG	AAG	AGC	TTC	CTG	AAG	GCT	GGT	GGA	GAA	926
Arg	Lys	Thr	Ala	Lys	Met	Leu	Lys	Ser	Phe	Leu	Lys	Ala	Gly	Gly	Glu	



275	280	285	
GTG ATT GAT TCA GTT ACA TGG CAT CAC TAC TAT TTG AAT GGA CGG ACT	974		
Val Ile Asp Ser Val Thr Trp His His Tyr Tyr Leu Asn Gly Arg Thr			
290	295	300	
GCT ACC AGG GAA GAT TTT CTA AAC CCT GAT GTA TTG GAC ATT TTT ATT	1022		
Ala Thr Arg Glu Asp Phe Leu Asn Pro Asp Val Leu Asp Ile Phe Ile			
305	310	315	320
TCA TCT GTG CAA AAA GTT TTC CAG GTG GTT GAG AGC ACC AGG CCT GGC	1070		
Ser Ser Val Gln Lys Val Phe Gln Val Val Glu Ser Thr Arg Pro Gly			
325	330	335	
AAG AAG GTC TGG TTA GGA GAA ACA AGC TCT GCA TAT GGA GGC GGA GCG	1118		
Lys Lys Val Trp Leu Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala			
340	345	350	
CCC TTG CTA TCC GAC ACC TTT GCA GCT GGC TTT ATG TGG CTG GAT AAA	1166		
Pro Leu Leu Ser Asp Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys			
355	360	365	
TTG GGC CTG TCA GCC CGA ATG GGA ATA GAA GTG GTG ATG AGG CAA GTA	1214		
Leu Gly Leu Ser Ala Arg Met Gly Ile Glu Val Val Met Arg Gln Val			
370	375	380	
TTC TTT GGA GCA GGA AAC TAC CAT TTA GTG GAT GAA AAC TTC GAT CCT	1262		
Phe Phe Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe Asp Pro			
385	390	395	400
TTA CCT GAT TAT TGG CTA TCT CTT CTG TTC AAG AAA TTG GTG GGC ACC	1310		
Leu Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu Val Gly Thr			
405	410	415	
AAG GTG TTA ATG GCA AGC GTG CAA GGT TCA AAG AGA AGG AAG CTT CGA	1358		
Lys Val Leu Met Ala Ser Val Gln Gly Ser Lys Arg Arg Lys Leu Arg			
420	425	430	
GTA TAC CTT CAT TGC ACA AAC ACT GAC AAT CCA AGG TAT AAA GAA GGA	1406		
Val Tyr Leu His Cys Thr Asn Thr Asp Asn Pro Arg Tyr Lys Glu Gly			
435	440	445	
GAT TTA ACT CTG TAT GCC ATA AAC CTC CAT AAC GTC ACC AAG TAC TTG	1454		
Asp Leu Thr Leu Tyr Ala Ile Asn Leu His Asn Val Thr Lys Tyr Leu			
450	455	460	
CGG TTA CCC TAT CCT TTT TCT AAC AAG CAA GTG GAT AAA TAC CTT CTA	1502		
Arg Leu Pro Tyr Pro Phe Ser Asn Lys Gln Val Asp Lys Tyr Leu Leu			
465	470	475	480
AGA CCT TTG GGA CCT CAT GGA TTA CTT TCC AAA TCT GTC CAA CTC AAT	1550		
Arg Pro Leu Gly Pro His Gly Leu Leu Ser Lys Ser Val Gln Leu Asn			
485	490	495	
GGT CTA ACT CTA AAG ATG GTG GAT GAT CAA ACC TTG CCA CCT TTA ATG	1598		
Gly Leu Thr Leu Lys Met Val Asp Asp Gln Thr Leu Pro Pro Leu Met			
500	505	510	

GAA AAA CCT CTC CGG CCA GGA AGT TCA CTG GGC TTG CCA GCT TTC TCA 1646  
 Glu Lys Pro Leu Arg Pro Gly Ser Ser Leu Gly Leu Pro Ala Phe Ser  
 515 520 525

TAT AGT TTT TTT GTG ATA AGA AAT GCC AAA GTT GCT GCT TGC ATC TGA 1694  
 Tyr Ser Phe Phe Val Ile Arg Asn Ala Lys Val Ala Ala Cys Ile  
 530 535 540 543

AAA TAA AAT ATA CTA GTC CTG ACA CTG 1721

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 824  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12

CTGGCAAGAA GGTCTGGTTG GGAGAGACGA GCTCAGCTTA CGGTGGCGGT GCACCCCTTGC 60  
 TGTCCAACAC CTTTGCAGCT GGCTTTATGT GGCTGGATAA ATTGGGCCTG TCAGCCCAGA 120  
 TGGGCATAGA AGTCGTGATG AGGCAGGTGT TCTTCGGAGC AGGCAACTAC CACTTAGTGG 180  
 ATGAAACTTT TGAGCCTTTA CTTGATTACT GGCTCTCTCT TCTGTTCAAG AACTGGTAG 240  
 GTCCAGGGT GTTACTGTCA AGAGTGAAAG GCCCAGACAG GAGCAAACCTC CGAGTGTATC 300  
 TCCACTGCAC TAACGTCTAT CACCCACGAT ATCAGGAAGG AGATCTAACT CTGTATGTCC 360  
 TGAACCTCCA TAATGTCACC AAGCACTTGA AGGTACCGCC TCCGTTGTTT AGGAAACCAG 420  
 TGATACGTA CCTTCTGAAG CCTTCGGGGC CGGATGGATT ACTTTCCAA TCTGTCCAAC 480  
 TGAACGGTCA AATTCTGAAG ATGGTGGATG AGCAGACCCT GCCAGCTTTG ACAGAAAAAC 540  
 CTCTCCCCGC AGGAAGTGCA CTAAGCCTGC CTGCCTTTTC CTATGGTTTT TTTGTCATAA 600  
 GAAATGCCAA AATCGTGCT TGTATATGAA AATAAAGGC ATACGGTACC CCTGAGACAA 660  
 AAGCCGAGGG GGGTGTATT CATAAAACAA AACCTAGTT TAGGAGGCCA CCTCCTTGCC 720  
 GAGTTCAGA GCTTCGGGAG GGTGGGGTAC ACTTCAGTAT TACATTCACT GTGGTGTCT 780  
 CTCTAAGAAG AATACTGCAG GTGGTGACAG TTAATAGCAC TGTG 824

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1899  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13

GGGAAAGCGA GCAAGGAAGT AGGAGAGAGC CGGGCAGGCG GGGCGGGGTT GGATTGGGAG 60  
 CAGTGGGAGG GATGCAGAAG AGGAGTGGGA GGGATGGAGG GCGCAGTGGG AGGGGTGAGG 120  
 AGGCGTAACG GGGCGGAGGA AAGGAGAAAA GGGCGCTGGG GCTCGGCGGG AGGAAGTGCT 180  
 AGAGCTCTCG ACTCTCCGCT GCGCGGCAGC TGGCGGGGGG AGCAGCCAGG TGAGCCCAAG 240  
 ATGCTGCTGC GCTCGAAGCC TGCCTGCCG CCGCCGCTGA TGCTGCTGCT CCTGGGGCCG 300  
 CTGGGTCCCC TCTCCCTGG CGCCCTGCC CGACCTGCGC AAGCACAGGA CGTCGTGGAC 360  
 CTGGACTTCT TCACCCAGGA GCCCTGCAC CTGGTGAGCC CCTCGTTCCT GTCCGTCACC 420  
 ATTGACGCCA ACCTGGCCAC GGACCCGCGG TTCCTCATCC TCCTGGGTTT TCCAAAGCTT 480  
 CGTACCTTGG CCAGAGGCTT GTCTCCTGCG TACCTGAGGT TTGGTGGCAC CAAGACAGAC 540  
 TTCCTAATTT TCGATCCCAA GAAGGAATCA ACCTTTGAAG AGAGAAGTTA CTGGCAATCT 600  
 CAAGTCAACC AGGATATTTG CAAATATGGA TCCATCCCTC CTGATGTGGA GGAGAAGTTA 660  
 CGGTTGGAAT GGCCCTACCA GGAGCAATTG CTACTCCGAG AACACTACCA GAAAAAGTTC 720  
 AAGAACAGCA CCTACTCAAG AAGCTCTGTA GATGTGCTAT ACACTTTTCG AACTGCTCA 780  
 GGACTGGACT TGATCTTTGG CCTAAATGCG TTATTAAGAA CAGCAGATTT GCAGTGGAAC 840  
 AGTTCTAATG CTCAGTTGCT CCTGGACTAC TGCTCTTCCA AGGGGTATTA CATTTCTTGG 900

GAACTAGGCA ATGAACCTAA CAGTTTCCTT AAGAAGGCTG ATATTTTCAT CAATGGGTCG 960  
 CAGTTAGGAG AAGATTATAT TCAATTGCAT AAACCTCTAA GAAAGTCCAC CTTCAAAAAT 1020  
 GCAAAACTCT ATGGTCCTGA TGTGGTCAG CCTCGAAGAA AGACGGCTAA GATGCTGAAG 1080  
 AGCTTCCTGA AGGCTGGTGG AGAAGTGATT GATTCAGTTA CATGGCATCA CTACTATTTG 1140  
 AATGGACGGA CTGCTACCAG GGAAGATTTT CTAACCCCTG ATGTATTGGA CATTTTTATT 1200  
 TCATCTGTGC AAAAAGTTTT CCAGGTGGTT GAGAGCACCA GGCCTGGCAA GAAGGTCTGG 1260  
 TTAGGAGAAA CAAGCTCTGC ATATGGAGGC GGAGCGCCCT TGCTATCCGA CACCTTTGCA 1320  
 GCTGGCTTTA TGTGGCTGGA TAAATTGGGC CTGTCAGCCC GAATGGGAAT AGAAGTGGTG 1380  
 ATGAGGCAAG TATTCCTTGG AGCAGGAAAC TACCATTTAG TGGATGAAAA CTTGATCCT 1440  
 TTACCTGATT ATTGGCTATC TCTTCTGTTC AAGAAATTGG TGGGCACCAA GGTGTTAATG 1500  
 GCAAGCGTGC AAGGTTCAAA GAGAAGGAAG CTTGAGTAT ACCTTCATTG CACAAACACT 1560  
 GACATCCAA GGTATAAAGA AGGAGATTTA ACTCTGTATG CCATAAACCT CCATAACGTC 1620  
 ACCAAGTACT TGCGGTTACC CTATCCTTTT TCTAACAAGC AAGTGGATAA ATACCTTCTA 1680  
 AGACCTTTGG GACCTCATGG ATTACTTTCC AAATCTGTCC AACTCAATGG TCTAACTCTA 1740  
 AAGATGGTGG ATGATCAAAC CTTGCCACCT TTAATGGAAA AACCTCTCCG GCCAGGAAGT 1800  
 TCACTGGGCT TGCCAGCTTT CTCATATAGT TTTTTTGTGA TAAGAAATGC CAAAGTTGCT 1860  
 GCTTGCATCT GAAAATAAAA TATACTAGTC CTGACACTG 1899

## (2) INFORMATION FOR SEQ ID NO:14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 592  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: singl  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14

Met Glu Gly Ala Val Gly Gly Val Arg Arg Arg Asn Gly Ala Glu  
 5 10 15  
 Glu Arg Arg Lys Gly Arg Trp Gly Ser Ala Gly Gly Ser Ala Arg  
 20 25 30  
 Ala Leu Asp Ser Pro Leu Arg Gly Ser Trp Arg Gly Glu Gln Pro  
 35 40 45  
 Gly Glu Pro Lys Met Leu Leu Arg Ser Lys Pro Ala Leu Pro Pro  
 50 55 60  
 Pro Leu Met Leu Leu Leu Gly Pro Leu Gly Pro Leu Ser Pro  
 65 70 75  
 Gly Ala Leu Pro Arg Pro Ala Gln Ala Gln Asp Val Val Asp Leu  
 80 85 90  
 Asp Phe Phe Thr Gln Glu Pro Leu His Leu Val Ser Pro Ser Phe  
 95 100 105  
 Leu Ser Val Thr Ile Asp Ala Asn Leu Ala Thr Asp Pro Arg Phe  
 110 115 120  
 Leu Ile Leu Leu Gly Ser Pro Lys Leu Arg Thr Leu Ala Arg Gly  
 125 130 135  
 Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly Thr Lys Thr Asp Phe  
 140 145 150  
 Leu Ile Phe Asp Pro Lys Lys Glu Ser Thr Phe Glu Glu Arg Ser  
 155 160 165  
 Tyr Trp Gln Ser Gln Val Asn Gln Asp Ile Cys Lys Tyr Gly Ser  
 170 175 180  
 Ile Pro Pro Asp Val Glu Glu Lys Leu Arg Leu Glu Trp Pro Tyr  
 185 190 195  
 Gln Glu Gln Leu Leu Arg Glu His Tyr Gln Lys Lys Phe Lys  
 200 205 210  
 Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Val Leu Tyr Thr Phe  
 215 220 225  
 Ala Asn Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu Asn Ala Leu

230	235	240
Leu Arg Thr Ala Asp Leu Gln Trp Asn Ser Ser Asn Ala Gln Leu		
245	250	255
Leu Leu Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile Ser Trp Glu		
260	265	270
Leu Gly Asn Glu Pro Asn Ser Phe Leu Lys Lys Ala Asp Ile Phe		
275	280	285
Ile Asn Gly Ser Gln Leu Gly Glu Asp Tyr Ile Gln Leu His Lys		
290	295	300
Leu Leu Arg Lys Ser Thr Phe Lys Asn Ala Lys Leu Tyr Gly Pro		
305	310	315
Asp Val Gly Gln Pro Arg Arg Lys Thr Ala Lys Met Leu Lys Ser		
320	325	330
Phe Leu Lys Ala Gly Gly Glu Val Ile Asp Ser Val Thr Trp His		
335	340	345
His Tyr Tyr Leu Asn Gly Arg Thr Ala Thr Arg Glu Asp Phe Leu		
350	355	360
Asn Pro Asp Val Leu Asp Ile Phe Ile Ser Ser Val Gln Lys Val		
365	370	375
Phe Gln Val Val Glu Ser Thr Arg Pro Gly Lys Lys Val Trp Leu		
380	385	390
Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala Pro Leu Leu Ser		
395	400	405
Asp Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys Leu Gly Leu		
410	415	420
Ser Ala Arg Met Gly Ile Glu Val Val Met Arg Gln Val Phe Phe		
425	430	435
Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe Asp Pro Leu		
440	445	450
Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu Val Gly Thr		
455	460	465
Lys Val Leu Met Ala Ser Val Gln Gly Ser Lys Arg Arg Lys Leu		
470	475	480
Arg Val Tyr Leu His Cys Thr Asn Thr Asp Asn Pro Arg Tyr Lys		
485	490	495
Glu Gly Asp Leu Thr Leu Tyr Ala Ile Asn Leu His Asn Val Thr		
500	505	510
Lys Tyr Leu Arg Leu Pro Tyr Pro Phe Ser Asn Lys Gln Val Asp		
515	520	525
Lys Tyr Leu Leu Arg Pro Leu Gly Pro His Gly Leu Leu Ser Lys		
530	535	540
Ser Val Gln Leu Asn Gly Leu Thr Leu Lys Met Val Asp Asp Gln		
545	550	555
Thr Leu Pro Pro Leu Met Glu Lys Pro Leu Arg Pro Gly Ser Ser		
560	565	570
Leu Gly Leu Pro Ala Phe Ser Tyr Ser Phe Phe Val Ile Arg Asn		
575	580	585
Ala Lys Val Ala Ala Cys Ile		
590	592	

## (2) INFORMATION FOR SEQ ID NO:15:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1899  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15

														GGG	3
AAA	GCG	AGC	AAG	GAA	GTA	GGA	GAG	AGC	CGG	GCA	GGC	GGG	GCG	GGG	48
TTG	GAT	TGG	GAG	CAG	TGG	GAG	GGA	TGC	AGA	AGA	GGA	GTG	GGA	GGG	93
ATG	GAG	GGC	GCA	GTG	GGA	GGG	GTG	AGG	AGG	CGT	AAC	GGG	GCG	GAG	138
Met	Glu	Gly	Ala	Val	Gly	Gly	Val	Arg	Arg	Arg	Asn	Gly	Ala	Glu	
				5					10					15	
GAA	AGG	AGA	AAA	GGG	CGC	TGG	GGC	TCG	GCG	GGA	GGA	AGT	GCT	AGA	183
Glu	Arg	Arg	Lys	Gly	Arg	Trp	Gly	Ser	Ala	Gly	Gly	Ser	Ala	Arg	
				20					25					30	
GCT	CTC	GAC	TCT	CCG	CTG	CGC	GGC	AGC	TGG	CGG	GGG	GAG	CAG	CCA	228
Ala	Leu	Asp	Ser	Pro	Leu	Arg	Gly	Ser	Trp	Arg	Gly	Glu	Gln	Pro	
				35					40					45	
GGT	GAG	CCC	AAG	ATG	CTG	CTG	CGC	TCG	AAG	CCT	GCG	CTG	CCG	CCG	273
Gly	Glu	Pro	Lys	Met	Leu	Leu	Arg	Ser	Lys	Pro	Ala	Leu	Pro	Pro	
				50					55					60	
CCG	CTG	ATG	CTG	CTG	CTC	CTG	GGG	CCG	CTG	GGT	CCC	CTC	TCC	CCT	318
Pro	Leu	Met	Leu	Leu	Leu	Leu	Gly	Pro	Leu	Gly	Pro	Leu	Ser	Pro	
				65					70					75	
GGC	GCC	CTG	CCC	CGA	CCT	GCG	CAA	GCA	CAG	GAC	GTc	GTG	GAC	CTG	363
Gly	Ala	Leu	Pro	Arg	Pro	Ala	Gln	Ala	Gln	Asp	Val	Val	Asp	Leu	
				80					85					90	
GAC	TTc	TTC	ACC	CAG	GAG	CCG	CTG	CAC	CTG	GTG	AGC	CCC	TCG	TTC	408
Asp	Phe	Phe	Thr	Gln	Glu	Pro	Leu	His	Leu	Val	Ser	Pro	Ser	Phe	
				95					100					105	
CTG	TCC	GTC	ACC	ATT	GAC	GCC	AAC	CTG	GCC	ACG	GAC	CCG	CGG	TTC	453
Leu	Ser	Val	Thr	Ile	Asp	Ala	Asn	Leu	Ala	Thr	Asp	Pro	Arg	Phe	
				110					115					120	
CTC	ATC	CTC	CTG	GGT	TCT	CCA	AAG	CTT	CGT	ACC	TTG	GCC	AGA	GGC	498
Leu	Ile	Leu	Leu	Gly	Ser	Pro	Lys	Leu	Arg	Thr	Leu	Ala	Arg	Gly	
				125					130					135	
TTG	TCT	CCT	GCG	TAC	CTG	AGG	TTT	GGT	GGC	ACC	AAG	ACA	GAC	TTC	543
Leu	Ser	Pro	Ala	Tyr	Leu	Arg	Phe	Gly	Gly	Thr	Lys	Thr	Asp	Phe	
				140					145					150	
CTA	ATT	TTC	GAT	CCC	AAG	AAG	GAA	TCA	ACC	TTT	GAA	GAG	AGA	AGT	588
Leu	Ile	Phe	Asp	Pro	Lys	Lys	Glu	Ser	Thr	Phe	Glu	Glu	Arg	Ser	
				155					160					165	
TAC	TGG	CAA	TCT	CAA	GTC	AAC	CAG	GAT	ATT	TGC	AAA	TAT	GGA	TCC	633
Tyr	Trp	Gln	Ser	Gln	Val	Asn	Gln	Asp	Ile	Cys	Lys	Tyr	Gly	Ser	
				170					175					180	
ATC	CCT	CCT	GAT	GTG	GAG	GAG	AAG	TTA	CGG	TTG	GAA	TGG	CCC	TAC	678
Ile	Pro	Pro	Asp	Val	Glu	Glu	Lys	Leu	Arg	Leu	Glu	Trp	Pro	Tyr	
				185					190					195	
CAG	GAG	CAA	TTG	CTA	CTC	CGA	GAA	CAC	TAC	CAG	AAA	AAG	TTC	AAG	723

Gln Glu Gln Leu Leu Leu Arg Glu His Tyr Gln Lys Lys Phe Lys	
200	210
AAC AGC ACC TAC TCA AGA AGC TCT GTA GAT GTG CTA TAC ACT TTT	768
Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Val Leu Tyr Thr Phe	
215	225
GCA AAC TGC TCA GGA CTG GAC TTG ATC TTT GGC CTA AAT GCG TTA	813
Ala Asn Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu Asn Ala Leu	
230	240
TTA AGA ACA GCA GAT TTG CAG TGG AAC AGT TCT AAT GCT CAG TTG	858
Leu Arg Thr Ala Asp Leu Gln Trp Asn Ser Ser Asn Ala Gln Leu	
245	255
CTC CTG GAC TAC TGC TCT TCC AAG GGG TAT AAC ATT TCT TGG GAA	903
Leu Leu Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile Ser Trp Glu	
260	270
CTA GGC AAT GAA CCT AAC AGT TTC CTT AAG AAG GCT GAT ATT TTC	948
Leu Gly Asn Glu Pro Asn Ser Phe Leu Lys Lys Ala Asp Ile Phe	
275	285
ATC AAT GGG TCG CAG TTA GGA GAA GAT TAT ATT CAA TTG CAT AAA	993
Ile Asn Gly Ser Gln Leu Gly Glu Asp Tyr Ile Gln Leu His Lys	
290	300
CTT CTA AGA AAG TCC ACC TTC AAA AAT GCA AAA CTC TAT GGT CCT	1038
Leu Leu Arg Lys Ser Thr Phe Lys Asn Ala Lys Leu Tyr Gly Pro	
305	315
GAT GTT GGT CAG CCT CGA AGA AAG ACG GCT AAG ATG CTG AAG AGC	1083
Asp Val Gly Gln Pro Arg Arg Lys Thr Ala Lys Met Leu Lys Ser	
320	330
TTC CTG AAG GCT GGT GGA GAA GTG ATT GAT TCA GTT ACA TGG CAT	1128
Phe Leu Lys Ala Gly Gly Glu Val Ile Asp Ser Val Thr Trp His	
335	345
CAC TAC TAT TTG AAT GGA CGG ACT GCT ACC AGG GAA GAT TTT CTA	1173
His Tyr Tyr Leu Asn Gly Arg Thr Ala Thr Arg Glu Asp Phe Leu	
350	360
AAC CCT GAT GTA TTG GAC ATT TTT ATT TCA TCT GTG CAA AAA GTT	1218
Asn Pro Asp Val Leu Asp Ile Phe Ile Ser Ser Val Gln Lys Val	
365	375
TTC CAG GTG GTT GAG AGC ACC AGG CCT GGC AAG AAG GTC TGG TTA	1263
Phe Gln Val Val Glu Ser Thr Arg Pro Gly Lys Lys Val Trp Leu	
380	390
GGA GAA ACA AGC TCT GCA TAT GGA GGC GGA GCG CCC TTG CTA TCC	1308
Gly Glu Thr Ser Ser Ala Tyr Gly Gly Ala Pro Leu Leu Ser	
395	405
GAC ACC TTT GCA GCT GGC TTT ATG TGG CTG GAT AAA TTG GGC CTG	1353
Asp Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys Leu Gly Leu	

410	415	420	
TCA GCC CGA ATG GGA ATA gAA GTG GTG ATG AGG CAA GTA TTC TTT			1398
Ser Ala Arg Met Gly Ile Glu Val Val Met Arg Gln Val Phe Phe			
425	430	435	
GGA GCA GGA AAC TAC CAT TTA GTG GAT GAA AAC TTC GAT CCT TTA			1443
Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe Asp Pro Leu			
440	445	450	
CCT GAT TAT TGG CTA TCT CTT CTG TTC AAG AAA TTG GTG GGC ACC			1488
Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu Val Gly Thr			
455	460	465	
AAG GTG TTA ATG GCA AGC GTG CAA GGT TCA AAG AGA AGG AAG CTT			1533
Lys Val Leu Met Ala Ser Val Gln Gly Ser Lys Arg Arg Lys Leu			
470	475	480	
CGA GTA TAC CTT CAT TGC ACA AAC ACT GAC AAT CCA AGG TAT AAA			1578
Arg Val Tyr Leu His Cys Thr Asn Thr Asp Asn Pro Arg Tyr Lys			
485	490	495	
GAA GGA GAT TTA ACT CTG TAT GCC ATA AAC CTC CAT AAC GTC ACC			1623
Glu Gly Asp Leu Thr Leu Tyr Ala Ile Asn Leu His Asn Val Thr			
500	505	510	
AAG TAC TTG CGG TTA CCC TAT CCT TTT TCT AAC AAG CAA GTG GAT			1668
Lys Tyr Leu Arg Leu Pro Tyr Pro Phe Ser Asn Lys Gln Val Asp			
515	520	525	
AAA TAC CTT CTA AGA CCT TTG GGA CCT CAT GGA TTA CTT TCC AAA			1713
Lys Tyr Leu Leu Arg Pro Leu Gly Pro His Gly Leu Leu Ser Lys			
530	535	540	
TCT GTC CAA CTC AAT GGT CTA ACT CTA AAG ATG GTG GAT GAT CAA			1758
Ser Val Gln Leu Asn Gly Leu Thr Leu Lys Met Val Asp Asp Gln			
545	550	555	
ACC TTG CCA CCT TTA ATG GAA AAA CCT CTC CGG CCA GGA AGT TCA			1803
Thr Leu Pro Pro Leu Met Glu Lys Pro Leu Arg Pro Gly Ser Ser			
560	565	570	
CTG GGC TTG CCA GCT TTC TCA TAT AGT TTT TTT GTG ATA AGA AAT			1848
Leu Gly Leu Pro Ala Phe Ser Tyr Ser Phe Phe Val Ile Arg Asn			
575	580	585	
GCC AAA GTT GCT GCT TGC ATC TGA AAA TAA AAT ATA CTA GTC CTG			1893
Ala Lys Val Ala Ala Cys Ile			
590	592		
ACA CTG			1899

## (2) INFORMATION FOR SEQ ID NO:16:

## (i) SEQUENCE CHARACTERISTICS:

(A)	LENGTH:	594
(B)	TYPE:	nucleic acid
(C)	STRANDEDNESS:	double
(D)	TOPOLOGY:	linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16

ATTACTATAG GGCACGCGTG GTCGACGGCC CGGGCTGGTA TTGTCTTAAT GAGAAGTTGA 60  
TAAAGAATTT TGGGTGGTTG ATCTCTTCC AGCTGCAGTT TAGCGTATGC TGAGGCCAGA 120  
TTTTTTCAGG CAAAAGTAAA ATACCTGAGA AACTGCCTGG CCAGAGGACA ATCAGATTTT 180  
GGCTGGCTCA AGTGACAAGC AAGTGTTTAT AAGCTAGATG GGAGAGGAAG GGATGAATAC 240  
TCCATTGGAG GCTTTACTCG AGGGTCAGAG GGATACCCGG CGCCATCAGA ATGGGATCTG 300  
GGAGTCGGAA ACGCTGGGTT CCCACGAGAG CGCGCAGAAC ACGTGCCTCA GGAAGCCTGG 360  
TCCGGGATGC CCAGCGCTGC TCCCCGGGCG CTCCTCCCGG GGGCTCCTC CCCAGGCCTC 420  
CCGGGCGCTT GGATCCCGGC CATCTCCGCA CCCTTCAAGT GGGTGTGGGT GATTTCGTAA 480  
GTGAACGTGA CCGCCACCGG GGGGAAGCG AGCAAGGAAG TAGGAGAGAG CCGGGCAGGC 540  
GGGGCGGGGT TGGATTGGGA GCAGTGGGAG GGATGCAGAA GAGGAGTGGG AGGG 594

## (2) INFORMATION FOR SEQ ID NO:17:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17  
CCCCAGGAGC AGCAGCATCA G 21

## (2) INFORMATION FOR SEQ ID NO:18:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18  
AGGCTTCGAG CGCAGCAGCA T 21

## (2) INFORMATION FOR SEQ ID NO:19:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19  
GTAATACGAC TCACTATAGG GC 22

## (2) INFORMATION FOR SEQ ID NO:20:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20  
ACTATAGGGC ACGCGTGGT 19

## (2) INFORMATION FOR SEQ ID NO:21:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21  
CTTGGGCTCA CCTGGCTGCT C 21



- (2) INFORMATION FOR SEQ ID NO:22:  
(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 23  
    (B) TYPE: nucleic acid  
    (C) STRANDEDNESS: single  
    (D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22  
AGCTCTGTAG ATGTGCTATA CAC 23
- (2) INFORMATION FOR SEQ ID NO:23:  
(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 22  
    (B) TYPE: nucleic acid  
    (C) STRANDEDNESS: single  
    (D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23  
GCATCTTAGC CGTCTTCTT CG 22
- (2) INFORMATION FOR SEQ ID NO:24:  
(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 23  
    (B) TYPE: nucleic acid  
    (C) STRANDEDNESS: single  
    (D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24  
GAGCAGCCAG GTGAGCCAA GAT 23
- (2) INFORMATION FOR SEQ ID NO:25:  
(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 23  
    (B) TYPE: nucleic acid  
    (C) STRANDEDNESS: single  
    (D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25  
TTCGATCCCA AGAAGGAATC AAC 23
- (2) INFORMATION FOR SEQ ID NO:26:  
(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 23  
    (B) TYPE: nucleic acid  
    (C) STRANDEDNESS: single  
    (D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26  
AGCTCTGTAG ATGTGCTATA CAC 23
- (2) INFORMATION FOR SEQ ID NO:27:  
(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 24  
    (B) TYPE: nucleic acid  
    (C) STRANDEDNESS: single  
    (D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27  
TCAGATGCAA GCAGCAACTT TGGC 24
- (2) INFORMATION FOR SEQ ID NO:28:  
(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28  
GCATCTTAGC CGTCTTTCTT CG 22

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29  
GTAGTGATGC CATGTAAGT AATC 24

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30  
AGGCACCCTA GAGATGTTCC AG 22

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31  
GAAGATTTCT GTTCCATGA CGTG 24

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32  
CCACACTGAA TGTAATACTG AAGTG 25

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33  
CGAAGCTCTG GAACTCGGCA AG 22

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single

(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34  
GCCAGCTGCA AAGGTGTTGG AC 22

(2) INFORMATION FOR SEQ ID NO:35:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 23  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35  
AACACCTGCC TCATCACGAC TTC 23

(2) INFORMATION FOR SEQ ID NO:36:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 22  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36  
GCCAGGCTGG CGTCGATGGT GA 22

(2) INFORMATION FOR SEQ ID NO:37:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 22  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37  
GTCGATGGTG ATGGACAGGA AC 22

(2) INFORMATION FOR SEQ ID NO:38:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 22  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38  
GTAATACGAC TCACTATAGG GC 22

(2) INFORMATION FOR SEQ ID NO:39:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 19  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39  
ACTATAGGGC ACGCGTGGT 19

(2) INFORMATION FOR SEQ ID NO:40:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40  
CCATCCTAAT ACGACTCACT ATAGGGC 27

## (2) INFORMATION FOR SEQ ID NO:41:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:41

ACTCACTATA GGGCTCGAGC GGC 23

## (2) INFORMATION FOR SEQ ID NO:42:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 44848  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:42

GGATCTTGGC	TCAGTGAAT	CTCTGCCTCC	CATGCAATTC	TTATGCATCA	50
GCCTCCTGAG	TAGCTTGGAT	TATAGGTCTG	CGCCACCACT	CCTGGCTACA	100
CCATGTTGCC	CAGGCTGGTC	TTGAACCTCT	GGGCTCTAGT	GATCCACCCG	150
CCTTGGCCTC	CCAAAGTGCT	GGGATTACAG	GTGTGAGCCA	TCACACCCGG	200
CCCCCGTTT	CCATATTAGT	AACTCACATG	TAGACCACAA	GGATGCACTA	250
TTAGAAAAAC	TTGCAATGGT	CCACTTTTCA	AATCACCCAA	ACATGTTAAA	300
GAAATTGGTA	TGACTGGGCA	TGGCACAGTG	GCTCATGCCT	GCAATCCTAG	350
CATTTTGTGA	GGCTGAGACG	GGCAGATCAC	GAGGTCAGGA	GATTGAGACC	400
ATCCTGACAG	ACATGGGTGAA	ATCCCATCTC	TACTAAAAAT	ACAAAACAAT	450
TAGCCGGGGG	TGATGGCAGG	CCCTGTAGT	CCCAGCTACT	CGGGAGGCTG	500
AGGCAGGAGA	ATGGCGTGAA	TCCAGGAGGC	AGAGCTTGCA	GTGAGCCGAG	550
ATGGTGCCAC	TGCACTCCAG	CCTGGGCGAC	AGAGCGAGAC	TCCGTCTCAA	600
AAAAAAAAAA	AAAGAAAGAA	ATTGGTATGA	CTGTTGACTC	ACAACAGGAG	650
TCAGGGGGCAT	GGGGTGGGGT	GTAAGATTAA	TGTCATGACA	AATGTGGAAA	700
AGAAACTTCT	GTTTTTCCAA	CTCCACGTCT	GCTACCATAT	TATTACACTC	750
TTCTGGTAGT	GTGGTGTTTA	TGTGTGAATT	TTTTTTCATA	TGTATACAGT	800
AATTGTAGGA	TATGAACCTG	ATTCTAGTTG	CAAAACTCAC	TATGAGCTTA	850
GCTTTTAAGT	TGCTTAAGAA	TAGGTAGATC	TATGCAAATA	ATGATAATTA	900
TTATTATTAT	TTTAAGAGAG	GGTCTCACTT	TGTCACCCAG	GCTGGAGTGC	950
AGTGGTGTGA	TTAAGGGTCA	CTGCAACCTC	CACCTCCAG	GCTCAAATAA	1000
ACCTCCACC	TCAGCCTCCC	CAGTAGCTGG	AACCACAGGC	ACGGGCCACC	1050
ACGCCTGGCT	AATTTTTTGT	ATTTTTTGTA	GAGATGGGGT	TTTCATCATGT	1100
TGCCCAGGCT	GTCTTGAAAT	TCCTCGGCTC	AAGCAATCCT	CCCACCTTGG	1150
CCTCCCAAAA	TGCTGGCATC	ACAGGCATGA	TGGCATCACT	GGCATCACAT	1200
ACCATGCCTG	GCCTGATTTA	TGCAAATTAG	ATATGCATTT	CAAAATAATC	1250
TATTTTTATT	TGTTGCCTTA	TTGGTGGTAC	AATCTCAAGT	GGAAAAATCT	1300
AAGGGTTTTG	GTGTTATTG	CTTACTCAAC	CAATATTTAT	TAGACTCTTA	1350
CTAAGCACCA	ACATGATCAC	ATGCCTGAGC	TATGGCTAGC	ATAGCGTGTG	1400
AGACAAACTT	AATCTCTGTT	TTGGTGGAGC	ATATAATCTA	GTAGATGAAG	1450
CCAAATGTTGA	GCAACATCAC	AATACTAACA	AATTGAGGAT	GCTACGAGAG	1500
TGTCTAACAA	ATTGAGGATG	CTACGAGAGT	GTCTAACAAA	TTGAGGATGC	1550
TATGAGAGTG	TGTCATGGAG	AGCTGCCTGG	AGATTGAGAG	AAAGCTTCCT	1600
TGAGGGAAGT	TACATTTTCA	CTGAAACACA	CTGCCATCTG	CTCGAGGTTT	1650
TGTAAGTGCA	TTACATCCCC	GATTCTGACA	CTTCACATCC	CGATTCTGAC	1700
ACTTCACCCA	GTTACTGTCT	CAGAGCTTGG	GTCCGCATGT	GTAAACAAG	1750
GACAGTATGC	ACTTGGCAGG	GTTGTGAGAA	GGGAAGAGAA	CACAAGTAAA	1800
GCACCTGTAT	CAGGCATACA	GTAGGCACTA	AGCGTGCGAT	GCTTGCTATG	1850
ATTATACATC	AGTGTAAGCA	TCAAGGAAAA	GCTGAAGAAA	AGTCTGACCA	1900
ACAGCGAAAG	ATAAATGCGC	AGAGGAGAAA	TTTGGCAAAG	GCTCCAAATT	1950
CAGGGGCAGT	CCGTACTCTA	CACTTTGTAT	GGGGGCTTCA	GGTCCTGAGT	2000
TCCAGACATT	GGAGCAACTA	ACCTTTTAAG	ATTGCTAAAT	ATTGCTTTAA	2050
TGAGAAGTTG	ATAAAGAAAT	TTGGGTGGTT	GATCTCTTTC	CAGCTGCAGT	2100
TTAGCGTATG	CTGAGGCCAG	ATTTTTTCAA	GCAAAAGTAA	AATACCTGAG	2150
AAACTGCCTG	GCCAGAGGAC	AATCAGATTT	TGGCTGGCTC	AAGTGACAA	2200
CAAGTGTTTA	TAGCTAGAT	GGGAGAGGAA	GGGATGAATA	TCCATTGGA	2250
GGTTTTACTC	GAGGGTCAGA	GGGATACCCG	GCGCCATCAG	AATGGGATCT	2300
GGGAGTCGGA	AACGCTGGGT	TCCCACGAGA	GCGCGCAGAA	CACGTGCGTC	2350
AGGAAGCCTG	GTCCGGGATG	CCCAGCGCTG	CTCCCGGGGC	GCTCCTCCCC	2400
GGGCGCTCCT	CCCCAGGCCCT	CCCGGGCGCT	TGGATCCCGG	CCATCTCCGC	2450
ACCTTTCAAG	TGGGTGTGGG	TGATTTCGTA	AGTGAACGTG	ACCGCCACCG	2500
AGGGGAAAGC	GAGCAAGGAA	GTAGGAGAGA	GCCGGGCAGG	CGGGGCGGGG	2550
TTGGATTGGG	AGCAGTGGGA	GGGATGCAGA	AGAGGAGTGG	GAGGGATGGA	2600
GGGCGCAGTG	GGAGGGGTGA	GGAGGGGTAA	CGGGGCGGAG	GAAAGGAGAA	2650
AAAGGCGCTG	GGGCTCGGCG	GGAGGAAGTG	CTAGAGCTCT	CGACTCTCCG	2700
CTGCGCGGCA	GCTGGCGGGG	GGAGCAGCCA	GGTGAGCCCA	AGATGCTGCT	2750

GCGCTCGAAG	CCTGCGCTGC	CGCCGCCGCT	GATGCTGCTG	CTCCTGGGGC	2800
CGCTGGGTCC	CCTCTCCCTT	GGCGCCCTGC	CCCACCTGTC	GCAAGCACAG	2850
GACGTGCTGG	ACCTGGACTT	CTTCACCCAG	GAGCCGCTGC	ACCTGGTGAG	2900
CCCCTCGTTC	CTGTCCGTCA	CCATTGACGC	CAACCTGGCC	ACGGACCCGC	2950
GGTTCTCTAT	CCTCTGGGG	TAAGCGCCAG	CCTCTGGTTC	CTGTCCCTTT	3000
TCCTGTCTCT	CTGACACCTA	TGTCTGCCCC	GCCAGCGGCT	CTCCTTCTTT	3050
TGCGCGGAAA	CAACTTCACA	CCGGAACCTC	CCCGCCTGTC	TCTCCCCACC	3100
CCACTTCCCG	CCTCTCATTC	TCCCTCTCCC	TCCTTTACTC	TCAGACCCCA	3150
AACCGCTTTT	TGGGGGGTAT	CATTTAAAAA	ATAGATTTAG	GGGTACAAAG	3200
TGCAGTCTCT	TTCCATGGGT	ATATTGCATT	GTGGTGGCAT	CTGGGCTCTT	3250
AGTGTAACCT	TCACCCGAAT	GTGTACATT	GTATCTAATA	GGTAATTTCT	3300
CATCCCTCAT	CCCTCTCCCA	CCCTCCCACC	TTTTGGAGTC	TCCAGTGTCT	3350
ACTATTCCAC	TAAGTCCATG	TGTACACATT	GTTTAGCGCC	CACTCTAAAT	3400
GAGCCTTTTT	GTTTCATTCA	TTCTGTAAGT	GTGAATAGG	CACCACCTAA	3450
GGTCAGGTAT	AAGTGGAAAT	TTGAAAAAGA	AACTGCCAC	TTGCCCCAGT	3500
ACTTCCCTAG	CCAAGAGGAG	GGAAACCAGG	CAGGTGCACC	TGAAGGCCTG	3550
TGAGTGTCTG	ATTTGCTGTG	CAGTGTAGGA	CAAGTAAGAT	TGTGCATATC	3600
CTTCTGTATT	TAAGCATGTG	TTAGGAAGAT	TTCTCTTTCT	TTTCTTTTCT	3650
TTTTCTTTTT	TCTTTTCTTT	TTTTTTTTTA	GGCAGATGAA	AAGGGCGTCA	3700
CAGAACAGGA	ATAAAAAATCT	AAATATTCAA	TAAATGAGAC	CTAGGAGACT	3750
ACTGCACTGA	CTTACAAAGT	CCTAATAAAA	AGATGTCTCT	CCAAAATGGG	3800
GCTGCAAAAT	GTGGTGCTGC	CTTATCAGCT	CTAAGTTTTT	TCCTTACCTG	3850
AGAAAGAAAG	AACCTGATGC	AGGTTACAGG	CTCCTGCCCC	ATGAATGCAG	3900
GCTGACTCCA	AGATGGGGAG	CTACAGGGAC	AATCCCAGGT	CTTCTAGGCC	3950
TCTTATTTAG	GGCCTGGGAG	CCTCCAGAGA	TGGCCACATC	TTGACCAAGC	4000
CAGATAGAGG	GAAAGATCAC	CATTATCTCA	CCTCTGTGTC	AAATACCTAG	4050
ATGCTGTCTT	CCTGTAGCCC	ACACTATAGT	TGCCAGCGCT	AATTTAATGG	4100
GTAGTGTACT	GGTTAAGAGA	TGGACAGACC	ATCCTGGCTT	GACTCTCAGC	4150
TCTGGCAAGG	ATGAGTGACT	TGGTTTTTCC	ATATCTCTTG	GCCACACCAA	4200
CCTTGATTTC	TTCAAGCTGA	GAATGGAATT	TCTCAAGCTT	GCCTCAAGGA	4250
TTATTGCCCG	AGGATTGTAT	GATATGGTAA	GAGCTTCTCA	GTGTTTGACC	4300
CATAGTAAGT	GTTTGACGTT	TCAAACGAAT	TGTTTCTTTT	TAGGACATGG	4350
TGAGCATTGG	GTAGCCATTC	ACCGGTTTTT	TGTTTCTTTT	GATCATAGTT	4400
AACCTCTCCT	TTTCTTCTG	GCACTACAAT	TTTCTGGTGG	GGAAGAATCC	4450
TTACTTTCTG	CCCTTCCCTT	TAAGGATAGG	AAGCTGATAC	TAGGCAGCAA	4500
CTAGTTGGGG	GATAGGAAGA	TTGTTCCAGA	GAAATGCTGA	ACCATAGGGC	4550
TCCAGATCAC	AGGACCCAG	TCTTAGCTTG	CTGGGGTGTG	GGGTGGGGGG	4600
GGGCGGTTC	TGAACATGGG	TATGAAGTAG	ATGTCCATTT	ACTGAAATGT	4650
GAGGACCTGA	GGCCTCTTCT	ATTGCTGTAG	CCAGCATATT	CCCCAACCTC	4700
TCCCCAAGAA	AGGACAGATG	GGGGTTCCCC	CCTGGAGTAA	CAGGTCCAAA	4750
AGAAAAAACA	TACAGTGGGA	CTTCCAGGAT	CTGGGCCTGA	TCACCCAGCA	4800
GTCAAGCTCC	CCGCAATTGA	CTAACACCCC	CCTAACACGT	AGAAATTCCA	4850
ATCTGCAATT	TAGTGAGGAT	GATACCTTTA	TTCTTCTTAA	ATACATCTCT	4900
TCATTTCCCA	GAGCACCTTT	TTTTCCCTCT	CTCTGCACCT	TTTTGTAAAA	4950
GACTGGAGTA	TAATGAAATA	CCAAGAGAGC	ATAACATGTG	ATACATAAAA	5000
CTTTTTTTCT	GGTTTACAAA	ACAGTTCATT	CTTGTCCTTA	CGTGCTTCTC	5050
TCCAAGGCTG	GCTGCTGTCT	GTTCACGCCC	GCTTCGCTTG	GAGAGGCCAT	5100
CTGCCATACC	TGCTCCCCAC	ACGCATCGAC	AAGCACACCC	AGAGTGTATT	5150
CTGCTAAGAC	CTAAAAGAGG	GAGGAACCCC	CTCTCTCAT	CTAAGACCTA	5200
GGTTCTAAAT	TAGAGTGTGA	GGGTCCATCT	CCCCAGGAGG	GGCAGAGGCC	5250
CCAAACAGCC	CAGCCATCTC	AGAAGACAAC	ACTAAGCTTT	GTAGGGGTCC	5300
ACAGTAGAGG	AGAGTAAGAC	GCCTGTTGTT	TAATTTATTA	CAGTTCCTCA	5350
AAAGTGAAAG	TGTGTGGGCG	GGATGGCAAG	AGCTGAGCAG	ACGAAAGCTG	5400
AAGGAATTAAG	GAAAGAGAGG	AGGACACAAA	CAGCTGACAC	TTCTCAGTT	5450
CTTGCTATTT	GGCTGGCCCT	GTCTAAGCA	CCTTCTAGGT	ATTAATCCAT	5500
TTAGTCTTGG	CTACAACACT	GTGAGTAACT	AGTTTTGTCA	CCCCCATTTT	5550
AAAAATGAAG	AAAGTGAGGC	TCAGGGAGGT	TAAGTAACTT	GGCCACAGTT	5600
TGAAACTAGA	CTCTGATCAC	ATGAGATAAT	AGTGCCCAT	AAAGGGGAAA	5650
GCAGATTATA	TTTTTTAAAG	GAAAGAGAGT	AGGATATGGT	AGAAAAAGAT	5700
TGTTTGGAAA	GGAAATTGAGA	GATTGATATA	ATGAAAAGAA	GCATTACAT	5750
GAGAGTAACA	GTATCAGGGC	CCAAACCTTC	ATCTAAGGTA	CTTCAAAGAG	5800
GCCTAAGCAA	ACTTAGTCAC	TGGCGTGGTT	CTAGTCTCCA	TGATGGCAAA	5850
TACATTGTGT	ACAGCCCAAC	TCCACACAAA	ACTTAAATAC	CAATGATAGA	5900
GCAATCTAAA	ATTTGAAAGA	AAAAATCTTT	CAATTTGTCT	TCTTCCAGAA	5950
GGGACTTAAT	CAAGAAACCA	ATCAAAATAC	TTCTTAAGCC	TAAGTGTGTG	6000
CAGAACTCCA	AAGAGAGCCC	AGCCCTAAAT	CAACACTGTC	CAATGGAAAT	6050
ATAATATAAT	GTGGGCCCTCA	TATGCAAGGT	CATATGTAAT	TTTTAAATTTT	6100
CTAGTAGCCA	TATTAATAAG	GTAAAAAGAA	ACAAGTGAAA	TTAATTTTAA	6150
TAATTTTATT	TAGTTCATAA	GATCCAAAAA	GTTTTCTCAG	CATGTAATCA	6200
ATATAAAAAA	ATTAATGAGG	TATTTATTAT	TCCTTTTCTC	AAACCAAGTC	6250
TATTCTATAA	TCTGGCGTGT	ATTATTTTACA	GCACCTTCTA	GACTATATTT	6300
CTTCTTTTCT	TTTTTTTTCT	CGAGACAATT	TTGCTCTTGT	CACCCAAGCT	6350
AGAGTACAAT	GGCGTTACCT	CGGCTCACTG	CAACCTCCGC	CTCCCGGGTT	6400
CAAGTTATTC	TCTGCGCTCA	GTCTCCCAAG	TAGCTGGGAC	TAGAGGCATG	6450
CACCACCAGC	CCTGGCTAAT	TGTGTATTTT	TAGTAGAGAC	AGGGTTTCAC	6500
CATGTTGGCC	AGGCTAACTT	CAAACTCCTG	AGCTCAGGTG	ATATGCCCAC	6550
CTCGGCCTCC	CAAAGTGTG	GGATTACAGG	CGTGAGCCAC	TGCACCCGGC	6600
CTCAGATTAA	CTATATTTCA	AGCGTTCACT	AGCCACATGT	AGCTAGTGTCT	6650
ATGGTAGTGG	ACAGTACAGA	TCTGCATTTT	AATTAAGACA	CGTATCAAG	6700
CATAGTTTCA	TAAATGCACG	TAAAAAAGG	TATAGTCTCT	AGTCTCTCT	6750

AGAAATCCTA	AATACTGCAG	AGCAAAAGTG	GTACGAACAG	CAATCTCAGT	6800
GATAATGCAA	CCATGCTTGC	TTTTCAATTG	AATTTGCTTA	TTTTCTTCA	6850
GCAAAAGTTCA	TCCATTTTTG	CCAATTCAAT	AAATATTTAC	TGATAAAAAAC	6900
TTTCAATATT	AGATTCTTGC	ATCTTCATAG	ACAGAGTTGC	TTTTACATT	6950
TAGAAAAATTA	CTTATCAATG	TTAAACACAC	GTTTTGATAA	CCAGTGTTGG	7000
AAAGAGGTGC	AGACTCCCCA	TGTGCCTATT	GATGGCAGAA	ATATTCACAG	7050
CCAAAGGGAA	ACAAAGGGCT	GGGGACAATC	ACACACCTCA	TGTCTCCTAA	7100
CTCCTGGGAA	GTGCTGTCCC	TCTGATTGAG	CTCTTATTAT	TGCCTTCCCC	7150
ACTAACCCCTG	TCCACTGTGC	CCTGGAGCCC	TTTGCAGGGT	TACCTGCTCT	7200
GTCCTCTCTCA	CAGAATATCT	CCTCTACCTC	CTTGTCCAAG	CTACAATTG	7250
GCTATTCTCT	GATGACACTG	TCTTCCCTGT	AGCCCTTTTG	AGTAATGGCT	7300
GCATATTCTC	CCATAGTCCA	GTTCTTTTCC	TGTTCTCCAG	TCTGGCTTCT	7350
GGATGACAGC	CCACTAGTTT	GAACCTCCATA	CTGCTATAGT	TCAAGTCCCT	7400
TTTGACTTGT	TACCTTGGGC	AAATTACCTC	CTTTTGTTC	GGTTCCTTGT	7450
TTGTAAATG	ACGATAATAA	TGCCATTGCG	TTCAGTGGGT	TATTTTGAAA	7500
TTGAGTGAAA	GAAGGCGGGT	AGCTTCCCTA	CACGCTCAGT	GTAGACTAGC	7550
CTGATGTGCA	TTACGGGTGA	TGCCATGACT	CAGTGTGTTT	TCCTCATCTC	7600
CACATCTGGC	TCTCATCCAG	TGCTCCTGCT	TACGGCACTC	TGTCCCCCTC	7650
TTACTTACTC	CCCCTTATTA	ACTGAAGACT	GGCACTGATC	TCACAGTTTC	7700
CTCTCCACTT	CCTAGTCTCA	CCATCATCCT	AGATGACTTC	AAGTCACCTA	7750
GATAAACTGT	CTCAGTTTCT	TCACTCACAT	TTTTTTATAA	CAGATAATGT	7800
TACACTCAAG	TTGTAAACAGA	ACCAGCTTAT	CCAGCTCATG	AAATGTATGC	7850
ATTTCACTCT	TAACCTGTAT	TCAGTGACAT	CCTGTGGGTA	TCTGGAAATC	7900
AGCCATGGTG	AGAATATTTA	CCATGGAAAT	TGGCAAATAC	TAAAAAGCAG	7950
AGCACCTTTT	TTTCTGAGAG	CCAGACCATA	GCTCTTCTAC	TCCATAGCAC	8000
CCATCATAAC	AAATTTTAAA	TACCTCCACT	GAACAGCTTC	TTCTCTCTC	8050
TACTTCTTCC	ATATCTGATT	TGAGCTTCTT	AATTTATCAT	GTGAACCACT	8100
CTTGTAATAA	TAACCCCAAA	TCCCTGTTC	ATTGTTCTTC	CTGCTAAAAA	8150
ACTAAACCTG	GTTTAGTCCA	ACCATATTTT	CTCTCTTGG	AATCTACAGG	8200
GTGGCCCAAA	AACCTGGAAG	TGGAAAAATA	TTACTTATTA	ATTTAATGT	8250
ATATTAAATA	GCCATTTTAA	TGCTTCAATT	CCAGTCTCAG	TGGCCACCCT	8300
GTATAGCTGG	GCTATTGAGC	TCTTGCGGGA	GGAGGGAGTG	GACAGTCTCC	8350
CAGCCACACA	GACTGATGTT	GCACCAACAA	TTTTTTAGCT	TCCAGACTTC	8400
CCTGGCCCTT	AGTGTTACCC	TTAACTCTCC	ATTTCTCTGC	CTTTCACATT	8450
CTCTACTTTT	TAAAAATCTC	TGACTCCACC	TTCACTTATC	CATTCTTAGC	8500
ACATGACCAT	ACTTCTGCTT	CCCAAAGAAA	ATGAGCAATT	ACTTCTTTT	8550
CCTTTTCTCT	CTGTGATCAA	ATCTGCAGAC	ATGTCATGCC	TAAGTCCAGC	8600
TTTCTCTCTT	TCTCTGATCT	CAGTCTGCTT	CTTCCATTTC	TGCCCTGAAT	8650
CCCGTCCCTT	CCCCAACCCC	CAAGGACTTC	GCTCTATCAG	TCACCTCTTC	8700
CCTCTCTCTG	ATCTTCAACT	CCTCCCATT	TACTGGCTTC	TTCTCAAGC	8750
CTTTCCCAAA	GCTTTTCCCA	TCTCAATTAC	CTCCTCGCAC	ATGCCCTCTG	8800
AGAAACCACC	CCGTTTCTTC	CCTCCCCTCG	GCAGCCTGTT	CTTCTGTTC	8850
TGCCCTCATG	ATGGCACCAT	CATTGTGTCA	CTAAAAATCAA	TCTCTCCGAC	8900
ATCATCAATG	GCTTCTCTTT	GTGGGAAAC	CTAATAAACA	CTTTATCTTA	8950
TTTGGTCTTT	GTTATGGGTT	GAATGAGGTT	ACCCCGAAAT	CCATATTAGA	9000
AGTCCTAACC	CCCAGTACCT	CAGAATGTGA	CTTTATTTGG	GAATAGGGTC	9050
ATTGCAGACG	TTATTAGTTA	GGATGAGGTC	ATACTGGAAT	GTGATGGGCT	9100
GCTTATCTAA	TATGACTGAT	GTCTTATATA	CAAGGAGAAA	TTTGAGAGACA	9150
GACACGCACA	TAGGGAGAAAT	ACCATGTGAT	GACAGGAGTT	ATGGAGTTGG	9200
ATGCAAAAAG	CTATGGGAAC	TTAGGAGAAA	GACCTGGAAC	AAATCCTTTC	9250
CTGCGCCTAG	AGAGGGAGTA	TGGCCCTGCC	ACTACCTTGA	ATTCAACGTT	9300
TCGGCTTTTC	AAAACGTGTA	GACAATACAT	TTCTGTTGTT	CAAACCAATT	9350
AGTTTGCAGT	ACTCTGCCAG	TGCAGCCCTA	ACAACTAAT	ACAGTCTCTT	9400
GGAGGCATTT	GGCAAGGTG	ACAATGGAAG	CACCTTCTTA	CCCCTTTAGG	9450
TCTGTGCGCT	TTCTTGTGGG	GGGGTGTGTT	CTAACAATTC	CTCTCCATCT	9500
CTCTCTCTCT	AGTTTGTCTT	AAACATTGGT	GTTCTTCAGA	CTTCTGACCT	9550
AGGCCTTCTT	TTCACTTCAC	ATATFCCCT	GGGTGGTCTC	ACCCACTTCC	9600
AGAAATTACT	TAAATTACTG	CTCATGCACT	ACTGTGCTGG	AAACTGTTTA	9650
ACAACCTGGCT	CTCTGGGAAG	AGGGGAGACT	GGTTGATGGT	TTTTGCTGAT	9700
TTCTGTGGTG	TAAATACTCC	CTCCATGGCC	AATTCCAAAC	TGCCAACAGT	9750
TTAACAACCTG	GCTCACAAT	TTTCTCCAAA	TTTAACATTT	GGCTTTCACA	9800
GGCCAACAAC	GTGGTACAGC	CAACTCCAGC	ACACCTCTGC	TTTTGTGTCA	9850
GAGAGAAGTA	ACTTATTTTT	GTACAAAAGG	TAAAAATAAA	ACACCTGCAG	9900
GCCCCCTTTT	TTTCTTAAAC	AAACTGCTCT	AGAAATAGAA	TAGCTGAAGC	9950
TTCTTTTATG	CATTCACTG	TTATTCCAT	GTCAGTGTGG	TGGTGGGATT	10000
ATTTTCTCTT	TATTTTCTT	GTATATGGTT	GAAATACTGT	ACCTTTGATC	10050
AGTTTATAGT	TTATGGCATG	TTTTGCACCC	ATATTAAATC	TAGTTTTTGT	10100
CAGAGGGCGT	CAATATTATT	TTCTCAAAAC	AAGAAATAT	TTCAATTGCAA	10150
AGGAGACAAA	CAAAAAGGTC	CTTAATACCA	AAACTTTGAA	ATGTGATTTT	10200
TTGTACTTGG	CAGTGTCCAA	GTGGTAAACC	CAAACAGTAT	TGGGTTTTTCA	10250
TTTTGTTCAG	GAAAGTCTTT	GTCTGGCAGC	GACTTACCCT	TACATCAGGC	10300
GGGCCTTGCT	CATTCAATTCA	CTTAAGTATT	TATTAACAC	CAGCGGTGTG	10350
CCAAGTACTT	ATCTAGGTAT	CGGGTAGATT	CTGATAAGTC	AGTCAGGTCC	10400
CTGCTCTCAG	GGAGCTTGCA	GCAGAGATGG	GGGCTGCAAT	AGAGAGTAAG	10450
CCAAGGAAAT	GAAAAAGGAA	GTTGATTTCA	GAGAGTGATG	AATGCTATGA	10500
AGAAAAATGAA	GGCAGCGCAG	TGTGATGGAG	AGTGACCCAA	GGTGGTACAG	10550
TTTGTACCTC	TAAAGACCA	ACTGTGACCC	AGGTCACTCA	CAGATGCCCG	10600
TCATGTGATG	CCACAGCAAC	TTTTCCAGGT	GCTCGTTTCC	TCCCACTTCC	10650
CAGTCTCTTG	CCCAGCGCGG	ACTGCTTACA	AATACAGCTA	GAGGAATCTA	10700
AATGAGGTTT	CTCTATCATC	AAACCAATC	AAAATGCCAA	GGAACAGAAT	10750

CAGTGCCTGG	CTGAAGGCAG	TGGAACAGGG	CCAGCCTGGA	GTGGTTCTCT	10800
CTGAGGAAGT	TCCTCATCTT	GGTTTTAGGG	CCATACCTTG	TGACCTGTGA	10850
GCTAGGGGTT	GCCAGTCCCT	GACATTTCTA	CTGAGGACTC	GCCTGTCTAT	10900
ATTCCCAGCC	TGTATGTGTC	TCCTGAGTTC	CAGACACACA	GGCGAAGCG	10950
CCTGATGGAT	GGAAGTATGT	TTTTTGGTGT	TCCATTGGTA	TCTCAAATTC	11000
TACAAAACCT	AGTCCCTCTG	CTCCTCCCTG	TTCTTCCCTA	TCTTCAGTCT	11050
ATCACCTGTT	CCTCATCCAG	CAAAATGATAT	TACCATCTTC	CAAGGAGCTT	11100
CCCAGGAGTA	ATCCTTGACT	CCTCCTCAAC	ATCCAATTAA	TAATCAAATC	11150
TAGGCCAGGT	ACAATAGCTC	ACGCCTATAA	TCCCAGCACT	TTGGGAGGCT	11200
GAGGCAGGTG	GATCATTGTA	GGCCAGGAGT	TCAAGACCAG	CCTGGCCAAC	11250
AAGGTGAAAC	CTGTCTCATT	TAAAAAAAGT	TATTTTAAAA	ACTCAAATCT	11300
ATTATTCTTA	CCTCTAAGTG	TGTCTTGAAT	TTATCCATCT	CTCTCCATCT	11350
CTGAGCTGTT	ACCTTACCTC	AGTCCATCAC	GTTTTGTCTA	CGTTAACATG	11400
ACCAGAGTCT	TGTTCTTAGT	CTGGTGAGGT	CACTCCAGCT	GCTTCAGATC	11450
CTTCCATGGC	TCACCGTTGC	CCTCATATAA	AGTTGGCACT	CCTGGACATG	11500
TGGCTTACGG	GGCCCTCCGT	GATGTGGCCC	TATTTGCTTC	TCCATTCTGT	11550
TCTCTCCAG	CCTCTCTGCC	CCCATCTCTA	GGCACCAACC	ACACCCTTCT	11600
GCTCGTCAAT	GGTCCAGCTT	TCTCTTCTAT	CTCTGGTCTT	TGGACAGACT	11650
TTTCCCTTCA	CCTGGAATGC	TTTCTTCAAT	CCTACCCAC	TCTCTTTAAT	11700
CTAGATAAGG	TTTATTCTTT	TTGAATGTCT	AGCAGTGAAA	CCATTTCCCC	11750
TGAAAAACCT	CCTCTAACCA	ACCCCTTACC	CTCAGCCCAA	GGTCTAGATT	11800
AGGAGTCCCT	CTGAATGTTT	CCATAGCATT	TTTAAAGAAT	TGCCTATTTA	11850
CTTGTTTCGTA	TCTATCACTA	AACTACAAAT	TGTATGAGAA	CAGCCACTAT	11900
CTCTGCCTGG	TTCAACCATC	ATCTCCAGCA	ACTAGCATAA	TGCCTGGCAG	11950
AGTCAGCCTG	CAACAAATAT	TTGTTGAATA	AATTAACAGA	TGGCTTTATC	12000
TCCTTAAAGTA	AATCTTGCTT	TTTTACCTTA	TTAAAACAGA	CGCACAGGCC	12050
AGGTGTGGTG	GCCCATGCC	GTAATCCAG	CACCTTGGCA	GGCTGAGGTG	12100
GGCGGATCAC	CTGAGGTGAG	GAGTTCAAGA	CCAGCCTGGC	CAACATGGTG	12150
AAACCCCATC	TCTAATAAAA	ATACAAAAAT	TAGCTGGGCA	TGGTGGTGGG	12200
TGCGTATAGT	CCAGCTTACT	AGGGAGGCTG	AGGCAAGAGA	ATCGCTTGAA	12250
CCCAGGAGGC	AGAGGTGGCA	GTGAGCCGAG	ATCATGCCAC	TGTACTCCAG	12300
CCTGGATGAC	AGAGACCCTG	TCTCAAAACA	CACACACACA	CACACACACA	12350
CACACACACA	CACACACACA	CACACACACC	AAGTTGTATA	ATTAAAAATA	12400
TAACGTGCTT	GTTATGGAAC	ACTTGTAATA	TACAGGAAAG	TAATGAAAAA	12450
GTCTACCATC	TAGCTCACCA	CATAATGACC	ATTGCTATCA	TCCTGGCATA	12500
ATTCTCTCCT	GTATATAAAT	ATATATTCTT	TTATTGTTAA	AATTACACTA	12550
TGAGTACTAT	TTATTTATTT	TACTGTGGCA	AAATGCGCAA	AACATAAAAT	12600
CTTGCCATTT	TAAGGTATGC	AGTTTGGTGC	ATTCACCACA	CTCACATTGT	12650
TGTGCAATA	TCACCACTAT	CTATCTCAGA	ACTTCTTCGT	CTTCCCAAAC	12700
TGAAACTCTG	TACCCATTAA	ACAATAGTGC	ATCCTCTGTT	TTCCCTCCC	12750
TACAATTTAT	TTTTATTTGG	GTTTGTACCA	AACTGAAAAA	AGCTGCTTCT	12800
TCCTTACTTA	GTTACAGATA	GCATTTCCAT	TTATTTAGCC	GTGGTTTTGA	12850
GGATGCCATG	ACAGATGCCA	TCCTTCCTAG	AGCTCTTTGG	GGCTGTCAGG	12900
TATTTTCAGC	AGGGTGAATT	CGGGTTGATA	ACATTTTAAA	ATCTCACTTT	12950
ATTCTGAGGT	TCTTAGTGTC	AGAGCCCAAC	GTATTTTGTG	GGACTCCCAA	13000
GTTACAAACA	AAAATATGGT	GAGGAGGAAT	CACTGAAGTT	TTAACACAAG	13050
AGACTTACAT	TTTGTTCAT	TTCTATCTTT	TAGTTTATTT	CCTAAGCATA	13100
AAGAAATACT	TTGAAAATTT	TACATAGCAT	TATACATATT	TAATTAAGCA	13150
TGAGCACATC	TTAAAACTTT	AAATTTTAGA	TCAGATCTTT	AATTCCTAGG	13200
ATATTAAAG	GTACTGGCAA	TTTGGCCAGG	TGTGGTGGTT	CACGCCTATA	13250
ATCCCAACAC	TTTGGGAGGG	TGAAGTGGGC	GAATTGCTAG	AGCCCAAGGAG	13300
GTGGAGGCTG	CAATGGCCTG	AGATCACGCC	ATCGTACTCC	AGCCTGGATG	13350
ATGAGAAATGA	AATCCTGTCT	CAAAAAAATA	AAAAAATAAA	AAAAGAAGAA	13400
GAAGAAGTAT	TGGCAATCAT	TGCTCCAGGA	ATAATTTTCT	GACTTGAAAT	13450
AAACCTACAT	GTAGACAAAC	TAATTAGGCC	ATTCCAAGAG	TTGCTAGCAT	13500
TGGTTTAATA	TGTTTTTCAG	GCATTCCAGG	AAGCAGTGTG	GCCAGCATTG	13550
CATGTTTGAT	ACTTCAGAAA	TGTATGACAG	GTGTTTCTCT	TACCCAGGTC	13600
TTCTGTTTTT	TTAGTTTTGC	TCATGTAAAT	ATTTATGAAC	ATCCTCATCT	13650
TTTTGAGGGA	AGGGATTATA	GATCATTTCT	ATTCCATTTT	CTAGCATTTG	13700
GTACCATTTCT	AAGCACATGA	TAGGCACCCA	TTTGGAGCAT	TTTTGGCTTG	13750
ACAGAAATATG	CATTTAGAAT	TGTTCAAATT	AGAGGTGTCA	GTGATGGGAA	13800
TTAGAATACT	ATATAATTCT	AAGTCATTTG	ACTTAAATAC	AAAAGAATGA	13850
TTTTCTTTGG	TGGGGAATGG	TGAAGGGAGG	CAGGAGTTAA	GAAGAGGAGA	13900
AGAGATCCTA	AGTCATTTAT	AACTTCTCT	GGAAAGACAG	GTGTGTGAAG	13950
ACTTTTTTAA	AAGTCATTCA	CCAAATTGTG	TGTGTGTGTG	TGTGTGTGTT	14000
TTAAATAGAC	TTTATTTTTT	AGAGCAGTTT	TAGGTTTACA	GCAAAATTGA	14050
ATGCAAGGAC	AGAGATTTCC	CATAAACCCC	CTGCCCACAC	ACATGCATAG	14100
CCTCCCTCAT	TATCAACATC	CCCACCAGAG	AGGTGTTTGT	TCTAGTTGAT	14150
GAACCTACAC	TGACACATCA	TTATCACCCA	AAGTCCATAG	TTACACGGCAG	14200
GGTTCACTGT	CGGTGTACAT	TCTATGGGTT	TGAGCAAATG	TATAATGACA	14250
TGTATCCACC	ATTATAGTAA	CATACAGAGT	ATTTTTCAGT	CCCTGCAATG	14300
CCCTGTTTCT	CCACCTATTCT	ATCCCTCCCT	CTCTGCATTT	CCACCCCAAG	14350
CCCCTGGTAA	CCGCTGATCT	TTTTACTGTC	CCATAGTTTC	GGACGATCTA	14400
TTTTTTCAGAC	AGACACAGAG	CTGTCTTTCC	CTTAGTTTCT	ATTCTATCAT	14450
TTCTTTCTCC	CATCCCATCA	TAAAAGGCTA	TGAGTTTTTT	TTAAGTGTG	14500
AACACCATCC	TACTTGTCAA	GTTAAACAT	AAGCTCCTGG	CTGGGTACAG	14550
TGGCTCATGC	CTGTAATCTC	AGCATTTTGG	GAGGCTGTGG	CAGAAGCATC	14600
ACTTGAAGCC	AGAAGTTTGA	GACCAGCCTG	GGCAACATAG	CAAGACCCCA	14650
TCCCTCCACA	CACAAACACA	CACACACACA	CACACACACA	CACACACACA	14700
CACACACACA	CACAAAAACA	AGCTCTTGCC	AGATTTAGAG	CTACAAATG	14750

CCCTCAGGTT	CCTAGAAGAT	CAGTCCTTCA	ATTAGATTCA	GATTGAGATG	14800
CTTCCTCTTT	TAAACAATGA	TTCCCTTTCT	ATCATGCCCA	ATAAGAAAAC	14850
AAATAAAAAA	TAAACAATAC	TGCCGTGAAT	CTCAGCTACC	CAGGAGGCAG	14900
AAGCAGAACT	GCTTCAACCC	GGCAAGCAGA	AGTTGCAGTG	AAGTGAGATC	14950
GCGCCACTGC	ACTCCAGCCT	GGGAAACAGA	GCAAGATTCT	GTCTCAAAAA	15000
CAAAACAATG	TGATTTCTCT	CTCTAAGTCC	TGCACAGGGA	AATGTTAAGA	15050
AATAGGTCCA	CCAGGAAAGA	AGGAAGTAAG	AATGTTTGAC	TAGATTGTCT	15100
TGGAAAAAAT	AGTTATACTT	TCTTGCTTGT	CTTCCTAACA	GTTCTCCAAA	15150
GCTTCGTACC	TTGGCCAGAG	GCTTGTCTCC	TGCGTACCTG	AGGTTTGGTG	15200
GCACCAAGAC	AGACTTCCTA	ATTTTCGATC	CCAAGAAGGA	ATCAACCTTT	15250
GAAGAGAGAA	GTTACTGGCA	ATCTCAAGTC	AACCAGGGTG	AAAATTTTTA	15300
AAGATTCACT	CTATATTTTA	ATTAACGTCA	GTCCGTCATG	AGAATGCTTT	15350
GAGAAAACCT	TTATTTCTCA	CACCTAACAA	TTAATGAGAT	TAACCTCCTC	15400
TCCCTCATAT	TGACCTGTGG	AGGAATCTGA	ACAAGAGGAG	GAGGCAGTGG	15450
GCAGGTTTCC	TTATCATGAT	GTTTGTCTAT	TTCAGTGTGA	GGCCTCACAA	15500
AAAAAATAAA	AAAAAATAAA	GGCGTCTGG	ATATAACTGA	GAGCTCATTG	15550
TACAGTAAAT	ATTAATAAAA	CAGTGATTGT	AGCTGAAGGA	TAGAAGTGT	15600
TGGAGGGAGC	AAGTGGGTAG	AATCGCGTCA	AACTAAAGAG	CATTTCTAGC	15650
CAAAGACACA	ATGATAGATT	GAAGGATATT	TATTCTAAAT	ATAGAATATG	15700
GGTGAACGAG	ATCTGTGGAC	TTCTGGGCTC	CAACGTTAGA	TTCTGATTTT	15750
AGCAAGCTTG	TCAGGGGATT	CTGATATTGA	AAGGCTGTGG	CCTTCACCTG	15800
AGAAACCTGC	CCTAGGGGGC	CATGAAAATT	TGTCCTGTCT	TTCAGAAGTG	15850
CTATCAGACA	TCAAATGGAA	GTTAAATCGT	ATCTTAACAA	TTACTAGGAT	15900
GGGCGCAGTG	ACTCACACCT	GTAATCCCAA	CACTTTGGGA	GGCTGAGGCA	15950
GGAGGATCAC	TTGAGCCGAG	GAGTTCGGGA	CCAGCCTGGG	CAACATAGAG	16000
AGACGTTGTC	TCTATTTTTT	AATAATTTAA	AGAGAAAAAA	ATACTGAAAA	16050
TATTTGATAC	ACCACGTAAT	TATAATAATG	TGTATATAAT	GTATATATTC	16100
ATTATGAGGA	ATATTTGATT	ATTTTCATATA	TTATATCTTT	TCCTTCTGTT	16150
TATTTTATCC	AGTTATGAAG	TATTTAGAAC	AATTCATCAG	TAATTGGGGG	16200
TAAATTGACA	GAATAGTAAT	CAGAGAAAAT	AGAAAAAGAG	AGATGGGTAA	16250
TCTTTGAATA	CCAGGTTGGA	GTTGTTTATG	GGTTTGTGTT	TTGTTTGGG	16300
GGCGTTTTTT	TAGACAGAGT	CCCACTCTGT	TGCCCCAGCT	GGAGTGCAGT	16350
GGCACAGCA	TGGCCCACTG	CATCCTTGAC	CTCTTGGGCT	CAAGCAATCT	16400
TCCCACCTTA	GCCTCCTGAG	TAGCTGGGAC	CACAGGTGCA	TGTCACCACA	16450
CCCAGCTAAT	TTTTTTATTT	TTTGTAGAGA	CAGTCTTTCT	ATGTTATCCA	16500
GGCTGATCTC	AAACTCCTGC	ACTCAAGTGA	TCCCCCTGCC	TTGGCGTCCC	16550
AAAGTATTGG	GATTATAGGC	ATAGCCACCA	CACCCAACCT	AGTTTCTATT	16600
TAGACTTGGC	CCTTTCCCAAC	CAGTCAATTT	TGTCCAAAAG	ATCTCATAAA	16650
TGTAGACAGG	AAACTGTCCT	TTGCTCATCA	GTTTTCTTCA	TCCTGTGTCT	16700
AGGGGGATGG	TCGGTGGGGG	AAACTGGGGT	TATGCAAGTT	CCTCTGAAAC	16750
ATCCTCTGTG	AGCCCCAGGA	TGGATGAGGC	ACCAGCCGCC	AGCGAGTCAG	16800
TGTGCAGCTT	TCCAGAAAGG	AAGTCATCAG	CCAGTCAGCC	GGCCTTGCCA	16850
GCCAGCACCC	GGCAACCCTG	CTGTCTTGTC	ATAAAGAAAT	GGTCTGCCTG	16900
ACAGGATGGT	GTGGATTTTT	CTTTTTTCTT	TTTTTTTTTT	TTGAGACAGG	16950
GTCTGGCTCT	CTGCCCCAGG	CTGGAGTGCA	ATGGCGGGAT	CTTGCTCAC	17000
TGCAGCTCT	GCCTCCCAAG	CTCAAGGCAT	CCTCCACCT	CGGTCTCCCG	17050
AGTAGCTGGG	ACCACAGGCA	CACACCACCA	CGCCCAACTA	AGTTTTCGTA	17100
TTTTTAGTAG	AGGCAGGGTT	TTACTATGTT	GTCCAGGCTA	GTCTCAAACT	17150
CCTGAGCTCA	AGCTATCCAT	CTGCCTTGCC	CTCCCAAGGA	GCTGGAATTA	17200
CAAGCGTGAG	CCACTGTGCC	TGACCAGGGT	GGATTTTTTC	AAGTGCACAT	17250
GTTGTGGTCC	CAGAAGCTCT	GATGGTACCA	AATTCCAAGC	GAAAAAAGT	17300
CAATGGTTCC	CACCCATCCT	ACCTCCCATG	ATGGCAAGAG	GAAATCACCA	17350
CAGTCGAGAT	ACAGTCCATG	TAAAACAAAT	TGCTATGGAT	TTTGAAAGTG	17400
AACCTTAAGA	GAATGCACT	ATGTTTTCTT	CATTAGAGTT	CTCTGGTAAT	17450
TTCCAGCTTT	TTTTTTTTTT	TTTTTTAGAC	AGTGTCTCGC	TTTGTGCCCC	17500
AGTGTACCCC	AGGCTGGAGT	GCAGTGACGT	GATCTCGGCT	CAGTGCACCC	17550
TCCGCCTCGT	GGGTGAAGT	GATTCTCTCG	CCTCAGCCTC	CTGAGTAGCT	17600
GTATTTTAGT	AGAGACGAGG	TTTCACCATT	TGGCCAGGCT	GGTCTCGAAC	17650
TCCTGACCTC	AAGTGATTCG	CCCATCTCAG	CCTCCCAAAG	TGCTGGGATT	17700
ACAGGTGTGA	GCCACTGCAC	CGGCCAGTA	ATTTCAAGCT	TCTGAGGAGC	17750
CCTTTGAATT	GTTAAATAAC	TTGTAGCTAT	GTCCAACATA	TCCATGTTCA	17800
GTGTATGTTT	GATATTTCTT	AGGAAACCTG	CCCTTGGTTG	TTTCTTTTGT	17850
GGTAATTCAT	GAGCCGGCAA	ATTTGACATG	TGTTACAGAA	TATACCTTTT	17900
CTCTGCTCTC	CTACCTCATA	ACCAGAACTT	AATTATCCTG	CTTTAGTCAC	17950
ATAAATAGCT	AACTAAATAA	ATATATGAGA	TTTCAGTCTG	CTCACTGTGA	18000
AAATAGACCT	TCTAAATGAT	CTCTTCCACT	TGCAGATATT	TGCAAAATAG	18050
GATCCATCCC	TCCTGATGTG	GAGGAGAAGT	TACGGTTGGA	ATGGCCCTAC	18100
CAGGAGCAAT	TGCTACTCCG	AGAACACTAC	CAGAAAAAGT	TCAAGAACAG	18150
CACCTACTCA	AGTAAGAAAT	GAAAGGCACC	CTAGAGATGT	TCCAGCCCCA	18200
AAGATATTTG	AATAGGTTGG	ACTCGGGCAC	CAATCTAGCA	AGTCCTACGG	18250
AAGTTGTATA	AAGCTGAAAA	TACTGAAGCA	TTTCCCAAAT	GGGAAATCCT	18300
AAACTCAAAA	CTTGCTTTTT	GGTTTTTTTG	TTTGTTTTGT	TTTTCTTCAT	18350
CTGACATTGC	TTAGTAGTCA	CAGAATGAAA	GATAAATCAA	TCATTTCATGA	18400
TCTAACAAATG	ACCTTCAGTG	CTCTAAAAAA	CTACGGAGTC	AAGGAAAAACA	18450
TGAATATATT	CCTCATGTAA	AATTAATAA	CAGACATATA	AAGGGCAAAA	18500
CATGAACATC	ATTCATACCT	TGAGGTCCGT	CCCCCTCCCA	GAAATAACCC	18550
CCAGTATGCC	TTGGTTTAGA	GCATTAAGCA	GGAGGGCCCT	GAGTCACTCC	18600
AGACAGTCTT	GACCACCAAG	CAGCATCTCT	TTTTTGTGTT	CTCTGTGGCT	18650
TTTGCARACA	CAGGGCTAGC	TCAGCTACCC	ATTAGTATGT	TTTCAGTCAC	18700
TAAACAGCTC	TTCCAGTCTT	CAATTAAGGA	TGACATTGTC	ACATGGGGCT	18750



TTAAAGCAAG	TGAAACAAGG	AACCCCCTTT	TTTTTTTTTT	TTGAGATGGA	18800
ATCTCACTCT	TGTCGCCCCAG	CCTGGAGTGC	AATGGCGCAA	TCTTGGCTCA	18850
CTGCAACCTC	CACCTCCAG	GTTCAAGAGA	TTCTCCTGCC	TTAGCCTCCT	18900
ATTCAATTATG	AGGAATATTT	GATTATTCAG	TTCTGTAGG	GTAAAGATAT	18950
TACCCCGGAT	CATATTATTG	ATTATTGAGT	AGCTGAGATT	ACAGGTGCCT	19000
GCCACCACGA	CCGGCTAATT	TTTTGTATTT	TTTAGTAGAG	ACAGGGTTTC	19050
ACCATGTTGG	CCAGGCTCCA	GGCTCGTCTC	GAACCTCTGA	CCTCAGGTGA	19100
TCCACCACCC	TCAGCCTCCC	AAAGTTCTGG	GATTACAGGC	GTGAGCCACC	19150
ACTCCTGGCC	ACAATCCTTT	TTTAACATATG	AAATATATTT	TTATCTGAAG	19200
TTTGATGTTT	ATACCCAACT	GAGGGATGAT	GTTCCCATAT	CTCAGTTAAA	19250
GAAATAACCT	GCTCAGATAC	TTCAAGCTCT	TCTTTTGACT	TTTGAAAAA	19300
AATGATCTTG	AAGTTACTAT	ACTTTGTTTG	GGTTAGTTAA	CATTATTTAA	19350
AGTATATTAT	TTTAATTAAT	TATCTTTGTA	AGATTTTACT	GATACTACC	19400
TGGAGTTCAA	TGTATCAGAT	GGATTTCAA	TTTATGTACA	TTTTTTATGT	19450
ATATGGTACA	GAAAAAATG	TGATCCATAA	GAAATCAGAA	AATAGCGCAT	19500
ATGCTAATAG	CTAAGTTGT	CCTCTAAAAA	ACTTATTTT	GCATTTTAA	19550
GAGGGGGATA	TACTCTGACA	CTTTAATAAG	TGTAATTAAT	TATTGACTGG	19600
AATTTGGCAT	GAGGCAGGGC	CATTTTCAGAT	CCCATTAAAG	GAATGACACA	19650
TACCAGAGAA	CCACAGAAGT	AAGGCCACAT	TTGTAATAAA	TCATTATAGC	19700
TCTGCTAGGA	GAAGACCCAG	TTGTATTAGG	TAATTAATGG	ATTTGCTCTT	19750
AAAACACATG	TCCCGGAAGA	TATAGGTGAG	TCTTGGGGGG	CCGCATTAAA	19800
CATTATACCA	ATGATCTTA	CATTTCTAAG	AAAGTTTAC	TACTTTACAG	19850
GATCTTTCTG	TTACCAAAAT	GGAAGGTTTC	CAACTCCAGG	ACTTGGCTTT	19900
CATAGTTCTT	ACACGAGGGG	AAATGCCTTC	CTTTGCTAAC	TATGCAACCA	19950
GGTTAGTTAG	TGTAAGTCCA	GCCACCCTGT	TGGCAATGCT	AAAAGGTACA	20000
ACAAACACAG	AATTTTATTT	GCATTTGTAA	ACATTTGATT	TCTGGCTCGA	20050
AATTTTCAGT	TTTCATGGGC	ACGTCATGGA	AACAGAAATC	TTCTGTGTTT	20100
AGTTTGGGCA	CCTACTCATT	GTAGTGACAA	ATATTTTCAGA	AGCCAATAGG	20150
GGATTCCACA	AATTGTTCTG	AACCTGTGGC	TGAGACTGGT	AATGGCTGAG	20200
TGACATGGGG	ACATACCACA	AAAGAAGAGG	TAGCAAAAGG	CTGCTGAGAT	20250
AAGGACATGT	TCATTGCTTA	GCTAGTGGCC	TGCACCCTTA	AAACACATGT	20300
CCCAGGCTGG	GTGCTGTGGC	TCACGCCCTGT	AATCCCAGCA	CTTTGGGAGG	20350
CTGAGGGGGG	TGGATTACCT	GAGGTCAGGA	GTTCGAGACC	AACCTGGCCA	20400
ACATAGTGAA	ACCTCATTT	TACTAAAAAT	ACAAAAATTA	GCCAGGCATG	20450
GTGGCGGGCG	CCTGTAGTCC	CAGCTACTCA	GGAGGCAGGC	AGGAGAATTA	20500
CTTGAATCTG	GGAGGCGAGG	GTGTGTGGTGA	GCCGAGATTG	CGCCACCGCA	20550
CGGTAGCCTG	GGGACAAAG	TGAGACTCTG	TCTCAAAAAA	ACAAAAACAA	20600
AAAAACAACA	AACAAAAAAC	AACAACAACA	AAAAAACGGG	TATCCCAGAA	20650
GATACAGGTA	AGTTTTCTAA	CACAGGTCTT	CTTGTATGGT	GCGTTCCT	20700
TAAGTAGAAG	ATGACAAAAA	CATTTGTCT	GAGAATATAG	ACTCACATT	20750
TAAACCTGTT	TGAGCAGGAA	AAGGAAGCAA	TGTTACAGAT	GTAATTCTGG	20800
GTGTGACTGC	AGAAAGGATG	ACTCCCTTAT	TAAAGTAGTC	ATCCTGAGTG	20850
AGCTAACCTT	TTGTACTTCC	TCCTCTCCTC	CTGTCCCT	CATCACCCCA	20900
TTCTTCCGTT	GCCTACACCC	AGGCCACAT	TGGATGCTGA	CATAGACTTA	20950
CATGGTACAG	TCCAAGGGAA	AGATCTGCCA	TTTTTTTCAA	TGTGTCATCT	21000
TGGTTATCTT	CATTCCAAGG	ATCTCTCCAC	TCTTTATACA	GTAAGAGATG	21050
AGAGTCTGGA	AAGGATTGGG	AATAAGATAA	TGAATTGTAA	GTTTTAAATT	21100
GTTCTTCTGA	TTTTGGGGAA	GGAGTAGGCT	AGGTGGTCTT	TCTGTTTTTT	21150
TTTTGTTTTT	TTTTTTAAAG	TAGATGTGGC	CAGACGTGGT	GGCTCACGCC	21200
TGTAATCCCA	GCACCTTGAG	AGGCTGAGGC	AGGTGGATCA	CTTGATGTCA	21250
GGAGTTCAAG	ACCAGCTTGG	CCAACACAGT	GAAACCCCTG	CTTTACTAAA	21300
AATACAAAAA	CTAGCCGGGC	TTGGTGGCGT	CCACCTGTAG	TCCAGCTAC	21350
TGCAGAGGTG	GAGGCAGGAG	AATCACTTGA	ACCCGGGAGG	TGGAGGTTGC	21400
AGTGAGCCAA	GATCATGCCA	TTGTACTCCA	GCCTGGGCGA	CAGAACAATA	21450
CTCTGTCTCA	AAAAAAAAGA	GAAAAAAGAA	GAAAAAAGA	ATGGATTGTA	21500
ACTCAGTCGT	CAATAGCCTC	TATTCAGGA	GATGTTACAG	TTGATTATGT	21550
TATAGGGGGT	GTATAATAGA	ATTTCGAGCT	ATGTAATTC	CAAGTGCATT	21600
TGGAAGAATG	AAGAAATGGA	GGAAGGGTAA	AGTATGAGTG	CAAGCATTTCC	21650
AGGTTTTTTG	AAAATGCTAT	AATCTTTGTT	CAGGGCTAGT	ACAAAGTGCT	21700
ATTTAGCTGT	AAGGGTTTTT	TGTGATTTAC	AGACAGTTTT	CACATGTGTC	21750
ATTTCAACCT	TGGTTTTATG	GCGAAGGCAT	GTGATGGTGC	TTGTCCCAGG	21800
ACTTTAGATC	CATATCTGAG	GTTCCTGTCTG	GGCAAAGATA	TTACCCCTGA	21850
TCATATTATA	GTCTATAAGT	GGGAGAGTTG	TGCCCTGGAGC	TCAAGTCTTA	21900
TGATTTCTGA	TCCAGGGCAC	TTCTTACAAC	ATGATTTTGC	AATATAAAAG	21950
CCTATAATGT	GTGACTAAAG	CAGGTCACCTC	ACCCCTTGTA	ACAGACTCTA	22000
GTAATGGTAC	TGCCACCAAA	CGGCTGCGTG	ATATTTGGGA	AAGACTTACC	22050
TTATTTGAAT	CTCAGTTTCC	TCCTAGAAAA	ATGAGGGTGG	AGGTTAAGCA	22100
TAGGCTGATG	ATCCTAAAGC	CTCCATACTG	CCCTAAACTG	TGGCTCTAAG	22150
ATCCAGTAGA	ATGCTGGGTC	ACAGGACTCT	AGGGAGCTTT	TCAAACCCAA	22200
ATGCTCTGTC	TTCTTGTATG	GTAGGCAGCA	GTTTATGGAA	GTGGCGGACA	22250
CAGCAAAATAT	CAAAATACCT	AAAGCAGCTT	GCAAGAGTTG	TTTCTGCCTA	22300
GTGGTCTTTA	TAGTTAAAT	TAAATAGTTA	ATTTTTTTTT	TTTTTGAGAC	22350
AGAGTCTTGC	TCTGTTACCC	AGGCTGCAGT	GCAGTGGCAC	AATCTCGGCT	22400
CACTGCAACC	TCCACCTCCC	GGGTTTGAGC	AATCTGTCT	CAGCCTCCCA	22450
AGTAGCTGGG	ACTACAGTTG	CATGCCACTG	CACCCAGCTA	ATTTTTGTAT	22500
TTTTAGTAGA	GACGGGGTTT	CACCATATTG	GGCAGGCTGG	TCTCGAACTC	22550
TTGACCTCAG	GTGATCCACC	TGCCCTCAGCC	TCCCAAAGTG	CTGGGATTAC	22600
AGCATGAGC	CACCTGACCC	AGCTTAAATA	GCTAATATTT	AATATTATTC	22650
TATAGTTTAT	CAAGTAATTC	AGGCCAAAGA	CTTAGAAACA	AAACAAAAAG	22700
CCACTTTTTAA	GGAGAAAGGG	TGTAAAGTTG	CCAGATAGAT	AGAGATCTTT	22750

CTTTTTTAAC	TACAAGAGTT	CAGGAATGAA	TTACTCTTTA	ACAAACGACT	22800
ATAGATATAC	ATGAAAATTG	GAAGGACTTA	TTATGCATAT	GATAATCAAT	22850
TTAAAGACAA	CACCTAAAAA	TATATTGTTG	CCACTCTCAA	AAAGTGGTAA	22900
TAGAACAGCT	AATGGTTTAA	AAAGCAGAGT	ACAGAAGTTC	CCAAACTTAT	22950
GGCACCTTAA	TATCGCAGAA	AACCTTTTAA	AGCATGCCTA	GGCCACAAAA	23000
AATACCTGTA	TTTTGATTAT	TAAATTGTAA	GGTCTACACA	ACCTAATAGT	23050
AATAGGTCCA	ATAGTAATGC	TGTCCAATAG	ATGTTGATGT	TTTTTTCCTT	23100
GCAAACCTTAA	AAGATCCTAC	AGTGCCTCTG	TAAATAGCAC	TGCTTGGTTA	23150
GAGTTGAATT	TCAGATAAAT	AATTTTTCCT	ATGTTAATTA	TTTTTCTTTT	23200
CTTTACTTTT	TTTTTTGTTT	TTTGTGTTTT	TTGTTTTTTT	TTTTGAGACA	23250
GGGTCTCATT	CTGTTGCCCA	GGCTGCTGTG	CAATGGCATG	ATCATGGCTC	23300
ACTGCAGCCT	TGACCTCCCT	GGGCTCAGGT	GATCCTCCCA	CCTCAGCCTC	23350
CCAAGTAGCT	AGCTGGGACT	ACAGGTGCTT	ACCATCATGC	CCGGCTAATT	23400
TTTGTGTTTT	TTGTAGAGAT	GTGGTTTTGC	CATGTTGCCC	AGGCTGGTCT	23450
TGAACCTCTG	GGCTCAAGTG	ATCCGCCCGC	CTCGGCCTCC	CAAAGTGCTA	23500
GGATGACAGG	CATGAGCCAC	TGCACCTGGC	CCCTGGGCGA	AGTATTTCTT	23550
AATGGTTTACA	TAGGACATAC	ACTAAACATT	ATTTATTGTC	TATATGAAGT	23600
TCAAGTTTAA	CTAGGTGCCC	TGCACCTTTA	GTGCTAAAT	CCTGTAGCTG	23650
TACCCATGCA	TTCACTGGTG	CTCCCCAGCT	TGCTTGCAC	AGAGTTTGGA	23700
AACCATAGTC	CTATAACTCT	AGGCCAATTT	TTTAATGTAA	AATTTGATTC	23750
ATTTTAAATT	AATAAATAAT	AACAGGAATT	TTTTTAAAAA	TTGTTTTAAA	23800
TATAATTAAA	ATTATCAAAA	TATTTTTTAA	CTGAACCTGT	GACTAGAGAT	23850
ATTTAGATTA	TGAAGAGTGG	GGTTTATGCT	AACTAATGAC	AGTCTGGCTA	23900
TGCATGTGGA	GCACCTAGCT	ATAAATTGTG	GCTTCCCCAA	TTCTCCTGAT	23950
GTCACTTGAA	CAAAACCTAA	GTGTGAGACC	AGAGCTTCTG	GTATCTTCCA	24000
TGGGATTTC	TTCAACAGCT	GGAGCAAATG	AAGTCAGATT	GATTTTTTTT	24050
AATTTGTCCA	ATTTTGTGTG	CTCAAAAACA	TAATTATAAT	CATTATTATG	24100
AAC TAGAATT	TCTTCAGTTT	AACAACAGAA	ATAGTTATTC	ATTATGAAAA	24150
GCGAATCTGG	AGGCCTTCAT	TGTGGTGCCA	ATCTAACCAT	TAAATTGTGA	24200
CGTTTTTCTT	TTAGGAAGCT	CTGTAGATGT	GCTATACACT	TTTGCAAACT	24250
GCTCAGGACT	GGACTTGATC	TTTGGCCTAA	ATGCGTTATT	AAGAACAGCA	24300
GATTTGCAGT	GGAAACAGTT	TAATGCTCAG	TTGCTCCTGG	ACTACTGCTC	24350
TTCCAAGGGG	TATAACATTT	CTTGGGAACT	AGGCAATGGT	GAGTACCCCA	24400
GGGAACAATT	CATTAATAAG	GAGATTCCCC	ACTAGCATTA	TTTCTTTTCT	24450
TTTCTTTTTT	TTTTCTTTTT	TTTTTTTTTT	GAGACAGAGT	CTCGCACTGC	24500
TGCCCAGGCT	GGAGTGCAGT	GGCGCCACCT	CGGCTCACTT	GAAGCTCTGC	24550
CTCCAAAAC	GCCATTCTCC	TGCCCTCAGC	TCCCGAGTAG	CTGGGACTAC	24600
AGGCACCCGC	CACCGCGCCC	GGCTAATTTT	TTTTTTTTTT	TTTTTTTTTT	24650
TTTTTTTGCA	TTTTTAGTAG	AGACGGGGTT	TCACCGTGTG	AGCCAGGATG	24700
GTCTTGATCT	CCTGACCTCG	TGATCTGCCC	TCCTCGGCCT	CCCAAGATGC	24750
TGGGATTACA	GGCGTGAGCC	ACCAGGCCCG	GCTAGCATTA	TTTCTTATGA	24800
CACCTTTTTT	TTTTTTTTTG	GACGGAGTCT	CGCTCTGTG	CCCAGGCTGG	24850
AGTGCAGTGG	CGCCATCTCG	GCTCACTGCA	AGCTCCACCT	CCCAGGTTCA	24900
CGCCATTCTC	CTGCCCTCAG	CTCCCGAGTA	GCTGGGACTA	CACGCACCCG	24950
CCACCACGCC	CGGCTAATTT	TTTTGTATTT	TTAGTAGAGA	CGGGGTTTCA	25000
CCGTGTTAGC	CAGGATGGTC	TCTATATCCT	GACCCCATGA	TCTGCCCGCC	25050
TCGGCCTCCC	AAAGTGGTGG	GATTACAGGC	GTGAGCCACT	GCGCCCGGCC	25100
AACACTCTTT	TTATTATTAG	CAAAATAACT	TCTGCCTGGG	CACATTCTTG	25150
CAAGTGCTCA	ACAATGCAAC	TTTTGGAAAG	GCAATGGGCA	GAAACTCCTG	25200
CTGTATTTAT	TCCAGAACCT	ATTATTGCTA	ATCCCAGTTT	ATGTTACATT	25250
TGAAGTGAGA	ACCAGTTGGA	GCCAGCAACG	TTCCCAGCTC	CAAAGTTCCC	25300
TTGAGATTTT	CAGAATCACT	TAACCTTATT	ATGCTTGGCA	ACCTGGACTC	25350
AGCAAACTG	GGAAAGTCAG	AGTTTGTTTT	ATTCATCCCT	TCCTTTCTCA	25400
GTTTCTCAAA	TGTGTCAGTT	AATCTCAGTA	ACCCCATTCG	AACCTTCATT	25450
ACCTGCCCAA	GCGGCTTAGA	ACTTGCCAGT	ATAGAATCCT	ACGTGGGTCA	25500
AGCTCCTGAC	TGTCTCCTTC	TTCACTCTTT	TTTTGCAAAG	AACCTGTAAA	25550
TTTTAACTAT	AAGTATTACAT	GATTCGCCAC	ATTTATTCAA	AACATAGAGT	25600
GCCTTTTTCA	CATATCAGCC	AATGGAAATA	AGGATTAAAT	GGGAAATGAA	25650
ATGTAGTAAT	AGGATAAGCA	CAAGTCTTCT	TCCTGCTCAA	ACTTTTTTTT	25700
TTTTTTTTTT	CAGACAAGAT	CTTGCTCTGT	TACCCAGGCT	GGAGTGCAGT	25750
GGCGTGTTCA	TAGCTCAATG	TAACCTCCAA	CTCCTGGGCT	CATGCAATCT	25800
CTCACACCTC	AGCCCCCTGA	TTAGCTAGGA	CTACACTATG	CCTAGCCAAT	25850
TTTTTTTCTT	TTGTCTGGTT	GTGTTGCCCA	GGCTGTCTCG	ATCTCCTGGC	25900
CTCAAGTAAT	CCTCCTGCCT	CGGCCTTCTA	AAGTGCTGGG	ATTATAGGCA	25950
TGAGCCACTG	TGCCCGGTCT	CAAACTTTT	TTTCCAAAGT	AAATGAAGTT	26000
ATTAGATATG	GAATATAGTC	TAGTTCCACG	ATATCCATAT	CCATTGGTTT	26050
ATTACCTCTA	TTATTAACTT	CAAATTGTTT	AATAGACCTT	CATATCTCAG	26100
TTATACAGTT	AAAATTTTTG	TTTTGTTTTT	CTGGAGTATC	TTATTTATAA	26150
CTATGAGTTT	TACTTTACTT	ATTTATTTTA	TTTTTTTGAGA	CAGACGCTTG	26200
CTCTGTCACT	CAGGCTGGAG	TGCGGTTGCG	TGATCATGGC	TCACTATGGC	26250
CTCGACCTTC	TGGGCTCAAG	TGATCCTCTC	CCTCAGCCTC	CCAAGCTGAG	26300
ACTACAGGCA	TGCACCACCA	CATCTAGCTA	ATTTTTTTTT	TTCCCATGCG	26350
AACAAGGCTT	TACTATGTTA	CCCAGAGTGG	TCTCAAACCT	CTGGCCTCAG	26400
GGGATCCTCC	TGTCTCAGCC	TACCAAAATG	CTGGGATTAC	AGGCATGAGC	26450
CATAGCGCCA	GACCTGGTTT	TACTTTTCTT	GACTTTGAAT	TACAAGTTTT	26500
TGTAATTTGG	AAAATGTTTT	GTTGCTTTTA	AATACTGCTG	TATGTTTCTT	26550
TTTAAATACA	ACATTTCTCG	ATATATATTT	TGAGAATTGC	TGCTTTTCAG	26600
AACCTAACAG	TTTCTTTTAA	AAGGCTGATA	TTTTCACTCA	TGGGTGCGAG	26650
TTAGGAGAA	ATTTTATTCA	ATTGCATAAA	CTTCTAAGAA	AGTCCACCTT	26700
CAAAAATGCA	AAACTCTATG	GTCTGTATGT	TGGTCAGCCT	CGAAGAAAGA	26750

CGGCTAAGAT	GCTGAAGAGG	TAGGAACTAG	AGGATGCAGA	ATCACTTTAC	26800
TTTTCTTCTT	TTTCCTTTTG	AGACAGAGTC	TCACCTCTGT	AGCCAGACTG	26850
GAGTGCAGTG	GTACAATCAT	GGCTCACTGC	AACTTCGACC	TCCCAGGCTC	26900
AAGCAATCCT	CCCATCTCAG	TCCCACAAAT	AGCTGGGACT	ACAGGTGCAC	26950
ATCACCACAC	CTGGCTACTT	TAAAAAAATT	TTTTTGTAGA	GATGGGGTCT	27000
CCCTGTGTTG	CCCAGGCTGG	TCTCTTGAAT	TCCTGTGCTC	AAGCCATCCT	27050
TCCACCTCAG	CCTCCACAG	TGCCAGGATT	ACAGGCATGA	GCCACCACAC	27100
CCAGCCACCA	CTTTCTTAA	AAAAAATAAA	AGATTCTCTC	TGGTAGACAA	27150
TCCTCAATAG	TCCACATGTT	ATTAACAAT	CTGCTGCCTG	AATACATGAT	27200
TTACCAAAAA	AAGGAAATTT	TGACGGGTTT	AGAATATCAA	GGGATCTGAG	27250
GCAAAATGTCA	CCTATGATAA	AATTTGCTAT	CAAAATTAGG	AAGTTTGTGT	27300
TTACCTGATC	CTAAAGCAGT	AACCAGCCCA	TTTCTAGGGA	ATAAACTCT	27350
CATGCGTATA	TTGTGCATAT	ATATGTATTA	TATGACTGAG	TGATAATAAA	27400
ATTTTTTTTC	TAGCTTCTTG	AAGGCTGGTG	GAGAAGTGAT	TGATTCACTT	27450
ACATGGCATC	AGTAAGTATG	TCTCCTATTC	TTAATACTAG	GAAAGTAAGG	27500
CTAGCTTTAT	TCTTTACCTA	GTATTCAAAA	AGTTAGTTCA	TTAACTGCC	27550
AATTGACTGC	AGTTCAAATA	AGAAACAAAT	AGTGTCTCAA	GTAGCACTGT	27600
ACTCCAATTT	TAATATTAAT	AAAAAAATTT	TTAAGTTATT	TTAAATAATG	27650
TAGTGGTTTC	TATAAAGATC	ACTTTATACA	GAAGAACAGT	GCCAATTAAC	27700
CCATGGAACA	TATAAGTAGC	TAAAACCAAT	TGCTTGCCAA	AGAACCAGTA	27750
ACCCAGGAGT	ACATGTCCTT	GCCACTGTGT	TTTTTCAAGA	CAGAGTAAC	27800
GATTTCTAGT	TACTTGCTAT	GAATGGACTC	CTCCTCATAA	CTCCCTTCCA	27850
TCTTGGTCTT	TCCCTAGTAG	AACCTCTACC	TTTTTTTAGT	AACAGGTGAG	27900
TGGGAGAGGT	AAGAAGGAGA	ATAAGGTCAG	CAATTAACCT	AAAAGCAGAA	27950
AGTAAATTTT	GTTATTTTTT	TTCTGAATAT	TTTCTGTGTA	ATTAGCTAC	28000
TATTTGAATG	GACGGACTGC	TACCAGGGAA	GATTTTCTAA	ACCCTGATGT	28050
ATTGGACATT	TTTATTTTAT	CTGTGCAAAA	AGTTTTCCAG	GTAATAGTCT	28100
TTTTAACTTT	TTTAATGTAA	AACCAGAATC	CTTATTTTAT	AGTCTAGCTA	28150
GTTCTAAATT	CTATAGGTAT	GTATATTTAC	ATGTTTTTCT	AATTTTAGAG	28200
AACAAGCACT	ATGACTTATC	CACTGTTAGT	TTTCCCTTAA	GCATTGGGTC	28250
TTACCCCATG	TACGTGATTA	GAAATTTGAA	ATATTTCCAA	TAGCCTTTAG	28300
TAGAATTAAC	TCACATAGAT	GATAAGAATG	GGTTGGTTCA	CTTCATGTTT	28350
CTTCCACAGC	CTACTATTTT	AATAAAAGAA	AGTTTCCCAA	GACCTAAATG	28400
ACTATGAACA	TATTTTATTA	CTATATAGGA	GGGGTGGGTC	TAGGAATACA	28450
AAGTTTTGAA	TGCTGTTAAT	CTTCAACACC	ACAGTTGAAA	CCACAGGTCA	28500
GCTTTTTTGC	AATTACCATT	GATACTTTTC	TGTTCTATAG	GTGGTTGAGA	28550
GCACCAGGCC	TGGCAAGAAG	GTCTGGTTAG	GAGAAACAAG	CTCTGCATAT	28600
GGAGGCGGAG	CGCCCTTGCT	ATCCGACACC	TTTGCAGCTG	GCTTTATGTG	28650
AGTGAAGCAG	CGCTGGGCTT	AGGGGTGAGA	GTGCAGCTCT	TCTCCATCCT	28700
TCTATTCTGC	TGAAATAGCT	CCCCAGCCAA	AAAGCAGATC	AAAGACCGTT	28750
TCAGTGGCTG	AGCCCCAAAA	TTTATGCCAG	ATTTTGCAAG	AAAATGATTT	28800
ACTAAAGCTT	GAGGGACATC	TTTAACAAGT	GTTCACAAAT	AATCACTATA	28850
AGGATGAATT	GTTTTCAGAA	TTTTGGCTTT	TAATTATGGC	CCATAAATAT	28900
GTCAGTAGT	CCTTACTCTA	AAGAAGTACA	CTGTAAGAGA	ATGCATATAG	28950
CCGGATATGG	TAGTTCCTCT	TAATCCCAAT	ACTTTGGGAG	GCCAAAGGTG	29000
GAGGATTGCT	TGAGCCCAAG	AGTTTGAGGC	TGCAGTGAGT	TATGATGGTG	29050
CCACTGCACT	CTAGACTGGG	CAACAGAGTG	AGACTGTCTT	TTTTTTTCCC	29100
CTCTGTCACT	CAGACTGGAG	GGCAGTGCCA	CGATCTCACC	TCACTGCAAC	29150
CTCTGCCTCC	CGATTCTGAG	CGATTCTCCT	GCCTCAGGCT	CCTGAGTAGC	29200
TGGGACTACA	GGAGTATCAC	CGCACTGGGC	TAATTTTTGT	ATTTTTAGTA	29250
GAGACGGGGT	TTTGACATGT	TGCCCAGGCT	GGTCTGAAAC	CCATGAGCTC	29300
AAGTGATCTG	CCTACCTCAG	CCTTCCAAAA	TGCTGGGATT	ACGGACATGA	29350
GCTACCACGC	CCGGCCACAC	CCTGTCTCTT	AAAAAAAAAA	AAAATGCAAG	29400
TTAGAGCATA	TTACAGCTTT	GTCTCTCAGG	AGGATACTTA	GTGTATGTAG	29450
CTATAAATCA	TAGATTCCCA	AGAAGTTTAG	AGCCTAAAGT	ATGAGGTCCC	29500
ACCAAGAGGG	CTATCATTAA	ATTTAAAGAT	TTGTTAAATC	ATCTCATTGT	29550
CCAACACCAC	AAACTTGATT	GCTTTAAAAA	ACTGGTTTAG	TTACATTTAG	29600
TAACCTCTATT	AGTCTCTTTA	ATCTATACTG	CTATATCCTC	ACATTGAGAT	29650
TTTTTTTCTT	TTCTCTTCCA	TCCTCATTTT	TTTTTCTCTC	ATCCTCATTC	29700
TTATAAGCCT	AGAATACATC	ACAAATCCTT	TATGCCCATG	GAAGCAAGAG	29750
GAATAAAGAA	TGGAGATGTT	TGTTTTGCCA	TTAACTAAAG	ATCTGGGGTG	29800
TCGGGGAGAA	GGGGGATAGA	GAAGGAGAAG	TGGGAAGAGG	TGTCCATAAT	29850
AGCTTAGGTG	CAATTCTGCT	TATTTTACAT	TTTACCCCGG	CTGACTGCCA	29900
CTTTTCTCTC	AGCCCTCACA	CATTGTTTGT	GCAGGGACCT	CATAGGACCA	29950
GGAATTGTCT	ATAGAGGTGG	GAATTTGTCT	CACCTGAAA	GGGATACCTC	30000
TAGCATGGTA	ATAGTCTTCT	AGGATTTGTT	ATCATATGGA	AAGATGTAAA	30050
GGGAGGGATT	CTGCTGCTGC	TGCTGCTGCT	GCATGCAGTT	GCCATTTTCA	30100
TTAAATGACT	TATTTATTAAT	TGATGACACT	TTTCTGGCTT	CCTGTTAATT	30150
CCTCCCTCAA	AGATCAATAA	ACCAGAACCA	GGCATGGTGG	CATGCACCTG	30200
TGGTCTCTGA	AGCAACCAAC	AGGTTTCACT	TGCTGCTGCT	CTAGATAGAG	30250
CCAATTATCA	AGACAGGGGA	ATTGCAAGAG	AGAAAGAGTA	ATTTATGCAG	30300
AGCCAGCTGT	GCAGGAGACC	AGAGTTTAT	TATTAATCAA	ATCAGTCTCC	30350
CCGAACATTC	GAGGATCAGA	GCTTTTAAAG	ATAATTTGGC	CGGTAGGGGC	30400
TTAGGAAGTG	GAGAGTGCTG	GTTGGTCAGG	TTGGAGATGG	AATCACAGGG	30450
AGTGGAAGTG	AGGTTTCTCT	GCTGTCTTCT	GTTCCTGGAT	GGGATGGCAG	30500
AACTGGTTGG	CGCAGATTAC	CGGTCTGGGT	GGTCTCAAT	GATCCACCCA	30550
GTTCAAGGTC	TGCAAGATAT	CTCAAGCACT	GATCTTAGGT	TTTACAACAG	30600
TGATGTTATC	CCCAGGAACA	ATTTGGGGAG	GTTCAGACTC	TTGGAGCCAG	30650
AGGCTGCATT	ATCCCTAAAC	CGTAATCTCT	AATGTTGTAG	CTAATTTGTT	30700
AGTCTGCAAA	AGGTAGACTT	GTCCCAAGGC	AGAAGGGGGG	TCTTTTCAGA	30750

AAAGGGCTAT	TATCATT TTTT	GTTTCAGAGT	CAAACCATGA	ACTGAAT TTC	30800
TTCCCAAAGT	TAGTTCAGCC	TACACCCAGG	AATGAAGAAG	GACAGCTTAA	30850
AGGTAGAAAG	CAAGATGGAG	TCAATGAGGT	CTGATCTCTT	TCACTGTCAT	30900
AATTTCCCTCA	GTTATAATTT	TTGCAAAGGC	GGTTTCAGTC	CCAGCTACTT	30950
GGGAGGCTGA	GACAGGAGGA	TTAATGGAGC	CCAGGAGTTT	GAGGTTGCAG	31000
AGAGCTATGA	TCACGCCACT	GCACCTCAGC	CTGGGTGACA	GAGTGAGACC	31050
CTGTCTCTAA	ATAAATAAAT	AAGTAAATAA	ATAAATACAT	AAATAAAATC	31100
AAGATGGTGT	GCAATTAGAA	TTGAGCGATT	TTGTTTCCAA	ACCTCAGAA	31150
AGCTTGGTCT	TGCTCTGTCC	CAGGTGGCTG	GATAAAT TGG	GCCTGTCAGC	31200
CCGAATGGGA	ATAGAAGTGG	TGATGAGGCA	AGTATTCTTT	GGAGCAGGAA	31250
ACTACCATT T	AGTGGATGAA	AACTTCGATC	CTTTACCTGT	AAGTGACCAT	31300
TATTTTCCTA	ATTCTAGTGG	AGTAGATTAA	AGTCAACTCA	GGACCTCTGG	31350
TGTTAACCTC	CTATGAACAG	TCAGTCTCT	CAGTAACTAG	CCAAATCATG	31400
AGATGATGAA	TTAGAAGGAG	CCTTAGATAG	CATCCAATCT	AACATTT TTT	31450
TGTGTGTTTG	AAGAGAAGAA	ATCAAGAGCT	AGGAATAACT	TTT TAAAGGT	31500
AAGCCATT TG	CAGTATAGTG	TGGATTTTGT	TTAAAAGGGG	ATAATTTGAA	31550
ATTTTATGAC	TCATTTATACA	AGACAAAATA	AGTTGGATTT	TCAAATGTTT	31600
TACAAAGTAA	ATCAAAGTTA	TAATTGCCTA	CAGTACGCAA	AGCTTCAAAA	31650
CATTTTTTAT	GTTATGAAAT	TGTAATTTAT	TTAACCTTAA	AATGAGCCAG	31700
TACCATGTGT	TTGCTTAAAA	ATCTCATGCT	AAGAATTTAC	TATGTTGTTA	31750
ATAATCTTCA	AGATATTTAT	GAATAAAGTC	TTATTTCTAA	TCCTTCCTCC	31800
AACTGTATCT	GGTGCTAAAT	CAGGAAATGT	TTCTTCCCAA	AAAGCCTCGT	31850
GGAAGATCTG	TATGTCTAAA	TATATGTCAG	GGATAATACA	GATGTAGCCC	31900
TGCGAAGCAT	GACCTTGATT	TTTATAGTCT	AAAATGTCAT	TTGCAGATAT	31950
CTATTTTCTA	AGAATAATTC	CTAAAAGAAT	TATTTGAATG	TTGTAGGAAA	32000
GCTAAGAAAT	TTTGCAAGAA	GCGTACGTGA	AAATATAAGC	TAGGCTTTTG	32050
TGGTTTGTGG	ATAGACTTCC	CAACAAAATT	GCTTTTATC	TATAGTGATC	32100
CAAGCTTGTG	GAACATATTA	GTCATCTTTT	TTTAGAAAAT	TCTTAGAAAA	32150
GTGATCTTGC	AAAAATGGAA	TTTATCTTTC	CCCAAGTATA	TTCTGTCATG	32200
TATAGAGTTA	AACTAAGCAT	AGTAATT TCA	CCAGACAAAC	ATTCAAAAATC	32250
TACTCTGTAC	CTTTTATCT	CATCCAAATT	TTCCCAGGGC	CCAGACATAA	32300
ACCTTTGCCT	TACGAAC TCT	TTGTATATGC	ACTAAATATG	CTTCTCCTTC	32350
AAGGTTCTCA	GTCAGCTAGA	AAAATGTGCA	AGAGTAAATG	GTACCCTTCT	32400
CAC TTGTAGA	TCCAAGAGAA	TTAGACTTAA	ACTCACTCTA	CATGTCTGTG	32450
ACTTTATTTT	ATTTGCAATG	CAGTCCCTGTG	AGGTGGCAAG	GCAGGTATCT	32500
TGGATCCATT	TTTTAGATAA	GGAAGTTCAA	ATTGAGAAGA	GGTTGCATGA	32550
TTTACAGGAA	GCCATACTGT	AGTCCTATGT	TACTCTTAAA	AATCCCATTTC	32600
AAATCCTGCT	TCTGAGGCCT	GCATACTTTC	TACCCTACCA	GTCAATTGACC	32650
CATGCTTATG	TCTCCTTTGA	AAACATTGAT	TCCACTCTTG	TCTCCAGTGA	32700
AAAAGTGGA	TTTAAGCAGA	GAAACAAAAG	CCATTTGTCT	TGTTAAGTCT	32750
ACTTTCCCTC	TACTTTCAAG	AAGGAAAGTT	GGGGTATGTG	TTGAATGGTG	32800
ATTTATTTAT	TTATTTATTA	TTTTAAAAAT	TGATACAAGG	TCCTACTGTA	32850
TTGTGCAGGC	TGGTCTCAAA	CTCCTGGGCT	CAAGTGATCA	TCCCACCTCA	32900
GCCTCCCACT	GTTGGGATTA	CAGCATGAAC	CATTGTGCCC	ACCACCGATC	32950
CGCAGTTTTT	TAGA AAAAAC	TTTTACTATA	GAAAATTTTA	ATCATATACA	33000
AAATACAGAG	GAAAGTATAT	GAACCCACTT	TAGGAGACTA	GAATATGCCA	33050
CCCCAAAATA	TGCCACTTTG	GCATAAGGAT	TATTTTCGAGC	TAAAGGCAAC	33100
TGGGAAGGAA	CACATAGAAG	AAAAGTTCTC	TGTCCTTCTC	CATTTGCCTA	33150
AAAGCAGGAC	ATGAATCTTA	AAAGTCCCCC	TCCTTCCCTT	TCTACCAGGA	33200
AAAACAAGAG	TTAATCACTG	AAGATAACTT	CAGACCCTTA	TCAGTGTAGA	33250
GATGGCACTA	GAAGAATCTA	TATTACATAC	TCATTTATTT	TCCTTCCAC	33300
AACTTGCCAC	CCCAGAGACT	AAAAATCCTT	TTCTTTTGTC	ATGTCTCTTG	33350
TCCAAAATTT	TGCTCTATAA	GCTGGAGTTC	TAAGCCACCT	CTTTGAGAAT	33400
TACTTGT TCC	CTGTTATTTT	CTGTTAACAT	ACATGTATTA	ATATACATGT	33450
TAACAAGCTT	CTGTTTGT TT	TTCTCCTGTT	TTCTGTCTTG	TTACAGAGGT	33500
CCATCCCAAC	TAAGA ACTAA	AGAGTAGGAG	GAAAATATAA	TTCTCTCTG	33550
CATACTTTGA	TCTTGT TTA	TCCGTAACCC	TTCCC ACTTT	TCACCTCCTA	33600
CCTATTAGAT	TACTTTGAAG	CAAAATTT CAG	ATATATTACT	TTATCTATAA	33650
ATATTTT CAGT	ATGTGCTAGT	TGTGGTGGCT	CACACCTGTA	ATCCCAACAC	33700
TTTGGGAAGC	TGAGGCAGGA	GGATCACTTG	AGCCAGGAG	TTCAAGACCA	33750
GCTACGGCAA	CAAAAAATCA	AAAAC TTATC	TGGGCATGGT	GGCACATGCC	33800
TGTGGTCCCA	GCTACATGAG	AGGCTGAGGC	AGGAGGATCG	CTTTAGCCCA	33850
GGAGTTGAG	GCTGCAGTAA	GCTGCATTCA	CACCACTGCA	CTCCAGCCTG	33900
GGTGACAGAG	TAAAGCCATG	TCTCAAAAAA	ATACATATTT	TAGTATGTAT	33950
CCTTTTGTGA	AAAAACACAAT	ACTTTTATCA	TACTTTAAAT	AATAACAATA	34000
ATTCTTTAGT	ATCACCAAAAT	ATTTTGT CAG	TGCTCTACAT	TTTCTTTATT	34050
GTCTAAAATA	TTGTTGATAG	TTATTCAAAT	CAGAATCCAA	ACAAGGTCCA	34100
TATATTACAT	TTGGTTGACA	AGTCTCTTAA	GTTTGTTCAT	CTTTAAGTTC	34150
TTCTTCCCTC	TCTTTCACTC	CTTGTAATTT	ATTAATGTGA	AAAAACAGGT	34200
AATTTGTTCT	ATAGTATTTT	CTACATTATA	GAGTTTGCTA	CATTTATTCC	34250
CTATGATATC	ATTTAGCATG	TTCTCTGTG	CCCTGTGTTT	CCTGTAAACT	34300
GGTAGTTATA	CTAGAAGCT	TGAGTTTATT	CAGGTTTTTA	ATTGTATTTT	34350
TTTTGCAAGA	ATTCTTTTAT	ATCTGCTTCT	GGAAGCACAG	AATGTCTGGT	34400
TGTGTCTGGT	TTTGATCTTG	ACAGCTACTG	ATGACCATTG	CCTAATCCAT	34450
TACTTTTATTG	GGGTGGGGGG	AATAAGGTTT	TAAAATAAAT	TTTTTTTAAA	34500
GATTTT TTTTA	ACTGTTATTT	TGAGACAGTG	TCTCATTTTCG	TTTCCCAGGC	34550
TGGAGTGACG	TGGCACAATC	ACGGCTCACT	GCAGCCTTGA	CCTCCTGGGA	34600
TCAGGTGATC	TCTCACCTC	AGCCTCCTGG	GTACCTGGAA	CTACAGGTGC	34650
ACACCACCAC	ACCTGGCTAA	TTTTTTGTAT	TTTGTGTACA	GAAGGGGTTT	34700
CATCATGTTT	CCACAGACTGG	TCTTGAATTT	CTGGGTTCAA	GTGATCTACC	34750

CAC TTCAGCT	TCCCAAATC	CTGGGATTAC	ACTTTGGCCA	CCGTGCCTGG	34800
CCTAAATGAA	ATTATTGTG	TCTAAACAGA	CAGAAGTTT	ACTTTAAAA	34850
TTTGTCTTTG	TGTGTACATG	TGTTTGTGTA	TGTGTGTGTG	TCTAAAAGTT	34900
TGGCTTTGAG	CTTTGCTTTG	AATTCCTTGA	TGAACAATA	CCAAGAATAC	34950
TTAAACTCTG	ATCATTCTTG	ACAGATATCC	CCTACAGGCT	ATGGCCTTTT	35000
GAATTGTGTC	CTCCAGTGAT	AAAAAGCAGC	AAGCACGATA	CTGCTCTCAG	35050
ATTCATGGTG	GTCACATGTG	AGGTGAAAA	AAAAAAAAG	ATGAATCCTA	35100
TTTAAATGCC	CCCAGGATA	CAGTGATACT	CTTTGTAGGA	TAATATTTTG	35150
CTTGCCACTG	GTTTCATTAA	ATAAGGCAT	AAGTAAAGAT	CTATTTTGT	35200
CTCTTTCTCC	CCAACCACCA	CAACTAGGAT	TATTGGCTAT	CTCTTCTGTT	35250
CAAGAAATTG	GTGGGCACCA	AGGTGTTAAT	GGCAAGCGTG	CAAGGTTCAA	35300
AGAGAAGGAA	GCTTCGAGTA	TACCTTCATT	GCACAAACAC	TGACAAGTAA	35350
GTATGAAACA	CACCCCTTAC	CAATCATCAA	GTTTATAGTG	GTAAAGCCTG	35400
AAC TTTACTC	AAACACCCTG	TTGCATGTGT	CTATACATTG	CATAAGTATA	35450
GGCAGTTGCA	ATTTAGTAAA	GTTTTATACA	ACGATTTTAT	TTTATTTTAT	35500
TTTTAGAAGA	AAAATGCTAC	TTTTGTTGTT	GTGTTTTTTT	GAGACGGGGC	35550
CTCGCTCGTC	ACCCAGGGTG	GAGTGCAGTG	GTGCAATCTC	AGCTCACTGC	35600
AACCTCCGCC	TCCCGGGTTC	AAGTGATTCT	TGAAGAGGAG	AACAATAATA	35650
ACAACAATAT	TATTTTCAAA	AGTTGTGACC	GCAGTTTCTG	GAGTTGAGAA	35700
GACATCGAGA	TTTTTGTAGC	CTCATACTCT	TGCTTTAGGT	AGCAAAAAAT	35750
GTTCTTAAAT	CTCAGGAATA	TTCTCTAGAT	AGGTTTCAAT	CTATCATTCC	35800
TGATAAGATG	ATGCTGAAAT	ACTAATTCTA	GCCAAAAAAG	ACCAGCTACC	35850
ATTTCCGATT	GTTGGGGACT	GGGAACCTCG	GATAGTGAGG	ACCCAGTAG	35900
GAAGTAGCGA	GGGGAATGGT	TTGAATGGAT	AAATTCATAA	AAAATGTCAG	35950
TAGATTTAAT	TTTCTTATAC	ATTTAGTCT	TTTTATAAAG	CTAGGAAAA	36000
CCCTTGTTTT	TATGGTTTAT	AATTTGAATT	CACATGAACC	CACAAAATTT	36050
GCCTTTTACC	TTCTATGTGC	TGAAAAATGA	TAGTCTGGCT	GGCCTCTTAA	36100
CAACCCAGCT	GGCAGAGCTG	TGAGGATCTC	AGTGTGCTCT	AGCCAGACA	36150
TTGGTAGCAT	GAACGGCAAC	ATTTTAAATT	GTGTTTTCAA	AATAGGAGCA	36200
CACTAGCGGT	CTAAAAACGAT	CATAAAAGAA	GGATACTAAG	AGGGCCACT	36250
GTCAATTATG	ATCTTAATAC	TTAGGATGCA	TTATGGATTG	TCATTATGGA	36300
TACTAATACT	TAGGATCACA	TTTGTAATTG	AGTTTTTAAT	TGCTTAAATT	36350
AGATACATAT	TTCTATTAAG	TTAACCTCTT	TGCTTTTAGT	CCAAGGTATA	36400
AAGAAGGAGA	TTTAACTCTG	TATGCCATAA	ACCTCCATAA	TGTCACCAAG	36450
TACTTGCGGT	TACCTATACC	TTTTTCTAAC	AAGCAAGTGG	ATAAATACCT	36500
TCTAAGACCT	TTGGGACCTC	ATGGATTACT	TTCCAAGTAA	GTAATTTTCC	36550
TTGTTCAATC	CAAACCTTTC	ATAAATTAT	TGGTGTTTAT	CAGAATAGAG	36600
AGTTTGGACA	GGGAGCAAAA	GACAAAGTCA	ACTATATCAA	GTTCTAATAA	36650
TTCTTAATAT	TCAGGAAATT	TATGTATGAA	TACTTACTAA	TATGAGTATA	36700
ACTCATCCTA	AGAGTCTTAA	GCAAAAGGAT	GTGAACACAA	ACTAGCAGTT	36750
ATCTTAGAGA	ATAAGTTTGC	ATTTCAAAAT	AAC TTGACAT	ATCAAGATCC	36800
ACTCAACGCA	TTTAAATTAT	TTACTCTAAA	AAGACATAAT	TCTTGGTAAC	36850
ACATTCACTA	AAGCAAAATA	TACCTTTATA	TAATTGCTAT	CAAAGGTATG	36900
TGGGTTGGTA	TAAATATACA	TACCATGTGA	GATCAGTGTG	ATTCTTTTAC	36950
AGCATTAATT	TTTATTGGTT	AGAGTAAGAA	AAAGAATAGC	TAGAGTATAT	37000
TTCTTAAGTA	GATTCTCATA	CAC TTTGGTT	TCAAAAACCA	ATTATTGACT	37050
ACATCTTATA	AAAGCCTGTA	TTCAATGGAG	TGCCAAAAAA	TGACTATGAG	37100
TCTTAAAGAG	TTAGGCATAT	AAATATTTTA	AGGTTTCTGT	TCAATGTATG	37150
TTGGAAGGAG	TTCC TTTCTC	ATGACTATTC	TCATATTGGA	GCATAAAAA	37200
AGTTTACAGG	CTTGGCGCAG	TGGCTCATGC	CTGTAATCCC	AATACTTTGG	37250
GAAGCTGAAG	CAGGCAGATC	ACTTCAGCCC	AGGAGTTTGA	GACCAGCCTG	37300
GGCAATATGG	CAAACTCTC	TCTACAAAAT	ATACCAAAAT	TAGCCAGGCG	37350
TGGTGGTGCA	TGCTGTAGT	CCCAGCTACT	TGGGAAGCTG	AGGTGGGAGG	37400
ATTGCTTGAG	CCCAGGGGGG	TCATGGCTGC	AGTGAGCTGT	GATGGTGCCT	37450
CTGTCAACCA	GCCTGGGTGA	CAGAGTGAGA	CCCTGTCTCA	AAAAAATAAA	37500
TAAATAAAAA	TTAAGAGTTT	ACAAAATTCT	CACCATCTCC	TCCCATCTTT	37550
GCAAAATGCCA	CATAAGTGAT	GTGTTCCAGG	ACTATTAGCC	TCGGAACCTG	37600
AGGCAGTACA	GTAAGCAGCG	TTTCTCCAAA	GTCTGTCC	CCACAGACAA	37650
ACATTATTTA	CACTGGGTAC	TGCTCTTTTA	TTTTTTCCCC	TCTATGCTTT	37700
ATTTTACTAT	AACTATAATC	ATATAACATG	TAATAGGAAA	AAGGCAGGGT	37750
CGGGGGAGAG	ATCCAGAAAT	CTTCCCAAGA	GCCTTTCCAA	CATAGCCTCT	37800
GTAGACATTT	TTTCTTTCTT	CTTTTTTTTT	TTTTTTTTTT	TTCTGAGACA	37850
GAGTCTCACT	CTGTTGTCCA	GGCTAGAGTG	CAGTGGCGTG	ATCTAGGCTC	37900
ACTGCAACCT	CCGCTCCTGT	GGTTCAAGCA	ATTCTCCAC	CTCAGCCTCC	37950
CTAGTAGCTG	GGATTAGAGG	CATGCATCAC	CACGCCTGGC	TAATTTTGT	38000
ATTTTATAGTA	GAGATGAGGT	TTCAACCATGT	GGGCCAGGCT	GGTCTTGAAC	38050
TCCTGACCTC	AAAGTGATCA	CCTGCCTTAG	CCTCCCAAG	TGCTAGGATT	38100
ACACGAGTGA	GCCACCGTGC	CCTGCCCTTA	TTACATTCTG	ATCACACATT	38150
TCATGTTTTA	TAATTGGAAA	ACTGGTGAAA	TTATAGACAA	TGTTTTGTTC	38200
CCCTAAATTC	TCTTTGATGA	GTATATATTA	CTTACACTCT	TCTGTCTTTA	38250
AAATTTTGCA	AAATAGTATC	CTAGATAAGT	TTATGAGTGC	ACAGTCTGTA	38300
CGCTTACTCA	TATTAATGAC	CTCGGAGAGT	TAAACAACAG	TCACCTTTAA	38350
AAATTTATTAC	TATCATTTATC	ATTATTTTGT	AGGCGGGGGT	CTCATCTGT	38400
CTCCCAGGCT	GGAGAGTAGT	GGTGGCGTCA	CAGCTCACTG	CAGCCACCGC	38450
TACCTGGGCT	CAAGTGATCC	TTCTCTCTCA	GCCTTCTGAG	TAGCTGAGAC	38500
CACAGGCTTA	TGCTACCCAA	CCTGGCTAAT	TTTTTAACTT	TTTGTAGAGA	38550
CGATGTCTCA	TTATGTTGCC	CAGGCTGGTC	TCAAACCTCT	AAGCTCAAGT	38600
GATCTTCTCT	AGCCTCCCAA	AGTGCTGGGA	TTACAGGCAT	GAAATCTG	38650
ACCCAGGCTT	AAAAATTAT	AGGCTCCTGC	ATAGTAAGAC	TTTAAATAAT	38700
ATTTAAATGA	ACATCTGGTT	TTTTTAAAA	AAAAATAGAG	ACAAGGTCTC	38750

ACTATATTGC	CCAAGCTGGT	CTCGAACTCC	TGGACTCAGC	CAATCCTGCT	38800
GCCTTAGCCG	CCCAAAGTGC	TGGGATTACA	GGCATGACCC	ACCTCATCTG	38850
GGCTGAGTGA	ACATATTTTT	AACATAAAGG	CCGTATTTTA	TATTTATCTC	38900
ATACATTTTTG	CCCAGCATCC	CCATTTCCGC	CGAATCTGTT	GCTTGCTAAT	38950
TCCTTCCAGC	TTCATTTTCAT	CTGAAATTTG	ACAAACATCT	TCTATTTCTT	39000
TGTCGTCATG	TTATTGACTT	CAGAAATATA	AATAAAACAC	TATACCCAAA	39050
TTAAACCCCA	CCCTCATTGC	CCAGCCTGAT	GTGAAAATAA	TCAGCATACA	39100
TTAAGCTTAC	CCTTGATATA	TGTGTAGCAT	CTTTTAGATA	AATATACAGC	39150
TGATTAAACA	ATATAGCCTG	ATGGTATAAT	ATCTTGCCCA	TGTACCTCAT	39200
CTTATCTCCA	GCAGGATTAA	TTCACAGTGA	TCAGATTTAC	CTTTAACTT	39250
TGTAGCAAAA	TATCCTCTCC	AAAAGCATAT	CTAAAACCTT	TGTGTGTACT	39300
CTTGCAAGTT	TCTTAATTTT	ATGCAGAACA	GGCTCTTACC	ACTGTTAGCT	39350
GGAGATATTT	TCAAGACCTA	TTTTTGTTTG	TGGTTTCCTG	ATGATGGTCA	39400
TGGCATTTC	CCCTTCACTC	CATCTAAAAA	TTGAGGTGAT	ACAGGCTTTT	39450
AAACAAACC	AACTCATATA	GACTGAGTAC	AACTGCAATG	CAGGCATGCT	39500
AACCTCTGCT	ACAATCATGG	GCGTGCTATT	GATATGTCTT	AAGTTACAGA	39550
ACACAGGGCT	GAGCGTCTCA	TTAGGTCAAA	ATGTAAACCA	GTTTTCTGCT	39600
TCAGTATGTC	TTAATGAGGA	CAGGGTGTGA	GAGATTTCTT	TAAGGAAAAA	39650
AAATATATAA	TAATGCTACA	TGGAAAAATA	TCTAACATTA	GAGAATTAAG	39700
TAAATAAACT	AATATACTCA	CACCATGGAA	TCTTGTGCAG	ACATTAATAA	39750
TATGTAGTGG	ATGGATGTTT	AATGGTGTGA	GAAAAAGTTA	GGATGTGCTG	39800
GGGTGGGGGG	AAGAATCAAG	TTTTAAGAAA	ATACAGTATA	CCCATACTTA	39850
AGTAAAAAAA	AAAAAAAAGG	TATGTACAGT	CATGTGTGTC	TTAATGATGG	39900
GGATACATTC	CGAGAAATGT	GTCGATAGGT	GATTTTCATCC	TTGTGTGAAC	39950
ATCATAGAGT	GAACCTTACAC	AAACCTAGAT	GGTCTAGCCT	ACTATGTATC	40000
TAGGCTATAT	GACTAGCCTG	TTGCTCCTAG	GCTACAAACC	TGTAAAGCAT	40050
GTTACTGTAG	CGAATATACA	AATACTTAAC	ACAATGGCAA	GCTATCATTG	40100
TGTTAAGTAG	TTGTGTATCT	AAACATATCT	AAAACATAGA	AACTAATGTT	40150
GTTGTGCTAC	AATGTTACAA	TGACTATGAC	ATTGCTAGGC	AATAGGAATT	40200
ATAATTTTAT	CCTTTTATGG	AACCACACTT	ATATATGCGG	TCCATGGTGG	40250
ACCAAAACAT	CCTTATGTGG	CATATGACTG	TATACATGTA	CACAAAAAAT	40300
AGATGAAAGA	ATGAATATAC	ATCAAAATAT	TTAAATGGT	TATAATGACT	40350
TAGGTTACTT	TTATTTATCT	TAGTAATAAT	AATGATGATA	GATAATACTT	40400
TTATAGTGTT	TACTATATAA	AAGACACTGT	TATAAGTGTT	CTACATACTT	40450
TACATGTATT	ACCTAAATGA	TATAAATATA	ACTCTGACAG	TAACATACTT	40500
TATACGTCTT	CTTTTCTTTT	TTTTTTTTTT	CTTTTTTTAG	ACAGAATCTT	40550
GCTCTACCAG	GCTGGAGTGC	AGGGTGCAAT	CTCGGCTCAC	TGCAACCTCC	40600
GCCTCCAGG	TTCAAAACGAT	TCTCATGTCT	CAGCCTCCTG	AGTAGCTGGG	40650
ACTACAGGCA	CACACCACCA	TGCCCGGCTA	ATTTTTGTAT	TTTTGGGTAG	40700
AGATGGAGTT	TTGCCATGTT	GGCCAGGCTG	ATCTTGAAC	CCTGGCCTCA	40750
AGTGATCTGC	CTCCCAAGT	GCTGGGATTA	CAGGTGTGAA	40800	
CACTGTGCT	CGGCCTAATC	TTACAAGTTT	TCAATATTTA	AAGAGTGCTA	40850
ACTTTGTTGA	CAATATAAAA	CATATTTGAG	AAAAAGAGAT	ATAAGCATCT	40900
TATTTAGAAAT	TATGAAAATA	TCAATAGACC	TACAGCCGAC	TAAAGCTTTT	40950
CTTCATAAGC	TCTTGCCAT	ATTGATTGCG	TCCTGTGAAT	ATGCATTAAT	41000
TTGATTTAAA	TAATAAGTAT	GTATAAGAAA	TAACACTTTT	CCTTAATTTT	41050
TAAGAACGTT	CAACAGTTT	TAATTTGAAT	TCCAATAGTG	AAATACATAG	41100
AAAATATAAA	ATTTTCTGTA	GTTAGCCAA	ATTGTTTTTG	TTTCACCACA	41150
GCATTCTACC	AAAATTTCTT	AATAACAGTA	AGAAAATGAA	TGCATACCTC	41200
TGCGAGGGAG	AGGGGAGTTA	GGCAGTTTAT	GGGCATAGTT	ACAAGTGAGA	41250
AATTTTCATTG	GCTACCATT	ACGCTAAAT	CATAAAACT	GCATTCAATT	41300
CTATATATCT	ATTTTCTTTA	CATAAAAAAG	GTTTCAATTA	TTGGCCATTA	41350
ATAAAAAATAG	CCACCATTCC	AGAAGTTGTG	TCATGTTTAT	CCTTTTTATA	41400
CCACCATCAT	ATTGCCTATT	ATATAGATTG	TGTGTGTTCC	ATTTTCTGTA	41450
ATGGGCCAGA	CAGTAAGTAT	TTCTGGCTTT	GGAGTCCATA	TGGTCTCTAT	41500
CATAACTACT	CATCTCTGCC	ATTGTAGCTT	AAAGATTATC	TAGGTCAAAT	41550
GCCTAAGTGA	TATAGTGTG	AAATACAAGT	TATATAATAT	AGGCTGCCAC	41600
AAAAAAAAT	TTATTTGGTC	TAAAAAAGAT	TTTCATGACT	TTGTAGCAGC	41650
ATGGGTGGGG	CATGCACCAC	TTGGTTAACT	CGGTGTATCT	TTCTCCTTTG	41700
CAGATCTGTC	CACTCAATG	GTCTAACTCT	AAAGATGGTG	GATGATCAAA	41750
CCTTGCCACC	TTTAATGGAA	AAACCTCTCC	GGCCAGGAAG	TTCACTGGGC	41800
TTGCCAGCTT	TCTCATATAG	TTTTTTTGTG	ATAAGAAATG	CCAAAGTTGC	41850
TGCTTGCAAT	TGAAAATAAA	ATATACTAGT	CCTGACACTG	AAATTTTCAA	41900
GTATACTAAG	AGTAAAGCAA	CTCAAGTTAT	AGGAAAGGAA	GCAGATACCT	41950
TGCAAAAGCAA	CTAGTGGGTG	CTTGAGAGAC	ACTGGGACAC	TGTCAGTGCT	42000
AGATTTAGCA	CAGTATTTTG	ATCTCGCTAG	GTAGAACACT	GCTAATAATA	42050
ATAGCTAATA	ATACCTTGTT	CCAAACTCTG	CTTAGCATTT	TGCATGTTTT	42100
ACTTTTATCT	AAAGTTTTGT	TTTGTTTTAT	TATTTATTTA	TTTATTTATT	42150
TTGAGACAGA	ATCTCTCTCT	GTCACCCAGG	CTGGAGTGCC	ATGGTGCAT	42200
CTTGGCTCAC	TGCAACTTTA	AGCAATTCTC	CTGCCTCAGC	TTCTGAGTA	42250
GCTGGGATTA	TAGGCGTGTG	CCACCACGCC	CAGCTACTTT	CTATATTTTT	42300
TGTAGAGATG	GAGTTTCGCC	ATATTGGCCA	AGCTGGTCTC	GAACCTCTGT	42350
CCTCGAAGTC	CTGCTCTCAA	GTGATCCACC	CGCCTCAGCC	TCTCAAAGTG	42400
CTGGGATTAC	AGGTGTGAGC	CACCACACCC	AGCAGTGTTT	TATTTTGTAG	42450
ACAGGGTATC	ATTCTGTTGC	CCAGGCTTGA	GTGCAAGTGT	GCAATCATAG	42500
ATCACTGCAG	CCTTTTAACT	CCTGGGCTCA	AGTCATCCTC	CTGCTTAGCC	42550
TCCCAAGTAG	CTAGGACCAC	AGACACATGC	CATCACACTT	GGCTATTTTT	42600
AAAAAATTTT	TTGTAGAGAT	GGGGTCTCGC	TATGTTACCC	AACTGGTCTC	42650
TGAACTCCTG	GACTCAATTC	ATCCTCCAC	CTTGGCCTTC	CAGGTGCTGG	42700
GATTTCTTTG	GGAGTACAGC	ATGGTACAGC	AGGAGATCAT	TTGATGTTAC	42750

CTCTGTGCAG	TGTTGCTAGT	CAGCGAAAGA	CTATAATACC	TGTGGGGACA	42800
GCGATTAGCC	ACCACAACCA	GTCTTTATTT	AAAGTTATTA	AAAATGGCTG	42850
GGCGCAGTGG	CTCACACCTG	TAATCCTAGC	ACTTTGGGAG	GCCGAGGCAG	42900
ATGGATCACC	TGACGTGAGG	AATTTGAGAC	CAGCCTGGCC	AACATGGTGA	42950
AACCCCATCT	CTACTAAAAA	ATACAAAAAT	TAGCTGGGTG	TGGTCCTGTA	43000
GTCCCAAGTA	CTTGGGAGGC	TGGGGCAGGA	GAATTACTTG	AACCCAGGAG	43050
GCAGAGGTTG	CAGTGAGCCG	AGATTGTGCC	ACTGCACTCC	AGCCTGGGTG	43100
ACAGAGAGAG	ATTCCATCTC	AAAAAAACAA	GTTATTAAAA	ATGTATATGA	43150
ATGCTCCTAA	TATGGTCAGG	AAGCAAGGAA	GCGAAGGATA	TATTATGAGT	43200
TTTAAGAAGG	TGCTTAGCTG	TATATTATC	TTTCAAAATG	TATTAGAAGA	43250
TTTTAGAAAT	CTTTCCTTCA	TGTGCCATCT	CTACAGGCAC	CCATCAGAAA	43300
AAGCATACTG	CCGTTACCGT	GAAACTGGTT	GTAAGAGAGA	AACTATCTAT	43350
TTGCACCTTA	AAAGACAGCT	AGATTTTGCT	GATTTTCTTC	TTTCGGTTTT	43400
CTTTGTGAGC	AATAATATGT	GAGAGGACAG	ATTGTTAGAT	ATGATAGTAT	43450
AAAAAATGGT	TAATGACAAT	TCAGAGGCGA	GGAGATTCTG	TAAACTTAAA	43500
ATTACTATAA	ATGAAATTGA	TTTGTCAAGA	GGATAAATTT	TAGAAAACAC	43550
CCAATACCTT	ATAACTGTCT	GTTAATGCTT	GCTTTTTCTC	TACCTTTCTT	43600
CCTTGTTTCA	GTTGGGAAGC	TTTTGGCTGC	AAGTAACAGA	AACCTCTAAT	43650
TCAAATGGCT	TAAGCAATAA	GGAAATGTAT	ATTCCCACAT	AACTAGACGT	43700
TCAAACAGGC	CAGGCTCCAG	CACCTTCAGT	CGTCACCAGG	GATCTGGGTT	43750
CTTCCAGCTG	CTCTGCTCTG	CCATCTTTAG	CGCTGGCTTC	ATTCTCAGAC	43800
TCTGGTAGCA	TGATGGCTGT	AGCTGTTTCA	TGGGCCCTTT	CAAACCTCAT	43850
AGCAACCAGA	GGAAGAAAAT	GAGCCATTTT	TTGAGTCTCC	TTCATAGACT	43900
TGAATAACTC	TTTTTCAGAG	CTTCTCACAG	CAAACCTCTC	CTCATGTCTC	43950
CTCATGTCTT	ATTGTTCAGA	AATGGGTAAT	GTGGCCATTT	CACCAAGTCA	44000
TGCCAACAAC	AACGAGGTTC	CTATAATTGT	CTCTGAGTAA	CCCTTTGGAA	44050
TGGAGAGGGT	GTTGGTCAGT	CTACAACTG	AACACTGCAG	TTCTGCGCTT	44100
TTTACCAGTG	AAAAAATGTA	ATTATTTTCC	CCTCTTAAGG	ATTAATATTC	44150
TTCAAATGTA	TGCCTGTTAT	GGATATAGTA	TCTTTAAAAT	TTTTTATTTT	44200
AATAGCTTTA	GGGGTACACA	CTTTTTGCTT	ACAGGGGTGA	ATTGTGTAGT	44250
GGTGAAGACT	CGGCTTTTAA	TGTACTTGTC	ACCTGAGTGA	TGTACATTGT	44300
ACCCAATAGG	TAATTTTTC	TCCATTACCC	TCCCTCCGCC	CTCTTCCCTT	44350
CTGAGTCTCC	AACATCCCTT	ATACCACTGT	GTATGTTCTT	GTGTACCTAC	44400
AGCTAAGCTT	CCACTTATAA	GTGAGAACAT	GCAGTATTTG	GTTTTCCATT	44450
CCTGAGTTAC	TCCCTTAGG	ATAACAGCCC	CCAGTTCCTG	CCAAGTTGCT	44500
GCAAAATACA	TTATCTTCTT	TTATGGCTGA	GTAATAGTCC	ATGGTACATA	44550
TATACCACAT	TTTCTTTATC	CACCTATCAG	TTGATGGACA	CTTAGGTTAA	44600
TTCCATTCAA	TTTCATTCAA	TTTAAGTATA	TTTGTAAGGA	GCTAAAGCTG	44650
AAAATTTAAAT	TTTAGATCTT	TCAATACTCT	TAAATTTTAT	ATGTAAGTGG	44700
TTTTTATATT	TTACATTTG	AAATAAAGTA	ATTTTATATA	CCTTGATATT	44750
GTATGACTAT	TCTTTTAGTA	ATGTAAAGCC	TACAGACTCC	TACATTTTGA	44800
ACCACTAGTG	TGTTGTTTCA	CCCTTTGTTA	TACTATCAGG	ATCCTCGA	44848

## (2) INFORMATION FOR SEQ ID NO:43:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2396  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:43

TTTCTAGTTG	CTTTTAGCCA	ATGTCGGATC	AGGTTTTTCA	AGCGACAAAG	50
AGATACTGAG	ATCCTGGGCA	GAGGACATCC	TAGCTCGGTC	AGATTTGGGC	100
AGGCTCAAGT	GACCAAGTGC	TTAAGGCAGA	AGGGAGTCGG	GGTAGGCTCT	150
GGCTGAACCC	TCAACCGGGG	CTTTTAACTC	AGGGTCTAGT	CCTGGCGCCA	200
AATGGATGGG	ACCTAGAAAA	GGTGACAGAG	TGCGCAGGAC	ACCAGGAAGC	250
TGGTCCCACC	CCTGCGCGGC	TCCCGGGCGC	TCCCTCCCCA	GGCCTCCGAG	300
GATCTTGGAT	TCTGGCCACC	TCCGCACCC	TTGGATGGGT	GTGGATGATT	350
TCAAAAGTGG	ACGTGACCGC	GGCGGAGGGG	AAAGCCAGCA	CGGAAATGAA	400
AGAGAGCGAG	GAGGGGAGGG	CGGGGAGGGG	AGGGCGCTAG	GGAGGGACTC	450
CCGGGAGGGG	TGGGAGGGAT	GGAGCGCTGT	GGGAGGGTAC	TGAGTCCTGG	500
CGCCAGAGGC	GAAGCAGGAC	CGGTTGCAGG	GGGCTTGAGC	CAGCGCGCCG	550
GCTGCCCCAG	CTCTCCCGGC	AGCGGGCGGT	CCAGCCAGGT	GGGATGCTGA	600
GGCTGCTGCT	GCTGTGGCTC	TGGGGGCCGC	TCGGTGCCCT	GGCCAGGGC	650
GCCCCGCGGG	GGACCGCGCC	GACCGACGAC	GTGGTAGACT	TGGAGTTTTA	700
CACCAAGCGG	CCGCTCCGAA	GCGTGAGTCC	CTCGTTCCCTG	TCCATCACCA	750
TCGACGCCAG	CCTGGCCACC	GACCCGCGCT	TCCTCACCTT	CCTGGGCTCT	800
CCAAGGCTCC	GTGCTCTGGC	TAGAGGCTTA	TCTCCTGCAT	ACTTGAGATT	850
TGGCGGCACA	AAGACTGACT	TCCTTATTTT	TGATCCGGAC	AAGGAACCGA	900
CTTCCGAAGA	AAGAGTTTAC	TGGAAATCTC	AAGTCACCCA	TGATATTTGC	950

```

AGGTCTGAGC CGGTCTCTGC TCGGGTGTG AGGAACTCC AGGTGGAATG 1000
GCCCTTCCAG GAGCTGTTGC TGCTCCGAGA GCAGTACCAA AAGGAGTTCA 1050
AGAACAGCAC CTACTCAAGA AGCTCAGTGG ACATGCTCTA CAGTTTTGCC 1100
AAGTGCTCGG GGTTAGACCT GATCTTTGGT CTAAATGCGT TACTACGAAC 1150
CCCAGACTTA CGGTGGAACA GcTCCAACGC CCAGCTTCTC CTTGACTACT 1200
GCTCTTCCAA GGGTTATAAC ATcTCCTGGG AACTGGGCAA TGAGCCCAAC 1250
AGTTTcTGGA AGAAAGCTCA CATTCTCATC GATGGGTGTC AGTTAGGAGA 1300
AGACTTTGTG GAGTTGCATA AACTTcTACA AAGGTCAGCT TTCCAAAATG 1350
CAAAACTCTA TGGTCTGAC ATCGGTCAGC CTCGAGGGAA GACAGTTAAA 1400
CTGCTGAGGA GTTTCCTGAA GGCTGGCGGA GAAGTGATCG ACTCTCTTAC 1450
ATGGCATCAC TATTACTTGA ATGGACGCAT CGCTACCAA GAAGATTTTC 1500
TGAGCTCTGA TGCGCTGGAC ACTTTTATTC TCTCTGTGCA AAAAATTCTG 1550
AAGGTCACTA AAGAGATCAC ACCTGGCAAG AAGGTCTGGT TGGGAGAGAC 1600
GAGCTCAGCT TACGGTGGCG GTGCACCCTT GCTGTCCAAC ACCTTTGCAG 1650
CTGGCTTTAT GTGGCTGGAT AAATTGGGCC TGTCAGCCCA GATGGGCATA 1700
GAAGTCGTGA TGAGGCAGGT GTTCTTCGGA GCAGGCAACT ACCACTTAGT 1750
GGATGAAAC TTTGAGCCTT TACCTGATTA CTGGCTCTCT CTTCTGTTCA 1800
AGAAACTGGT AGGTCCCAGG GTGTTACTGT CAAGAGTGAA AGGCCAGAC 1850
AGGAGCAAAC TCCGAGTGTA TCTCCACTGC ACTAACGTCT ATCACCACG 1900
ATATCAGGAA GGAGACTTAA CTCTGTATGT CCTGAACCTC CATAATGTCA 1950
CCAAGCACTT GAAGGTACCG CCTCCGTTGT TCAGGAAACC AGTGGATACG 2000
TACCTTCTGA AGCCTTCGGG GCCGGATGGA TTACTTTCCA AATCTGTCCA 2050
ACTGAACGGT CAAATTCTGA AGATGGTGGA TGAGCAGACC CTGCCAGCTT 2100
TGACAGAAAA ACCTCTCCCC GCAGGAAGTG CACTAAGCCT GCCTGCCTTT 2150
TCCTATGGTT TTTTGTGCAT AAGAAATGCC AAAATCGCTG CTGTATATG 2200
AAAATAAAAG GCATACGGTA CCCCTGAGAC AAAAGCCGAG GGGGGTGTTA 2250
TTCATAAAAC AAAACCCTAG TTAGGAGGC CACCTCCTTG CCGAGTTCCA 2300
GAGCTTCGGG AGGGTGGGGT ACACTTCAGT ATTACATTCA GTGTGGTGTT 2350
CTCTCTAAGA AGAATACTGC AGGTGGTGAC AGTTAATAGC ACTGTG 2396

```

## (2) INFORMATION FOR SEQ ID NO:44:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 535  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:44

```

Met Leu Arg Leu Leu Leu Trp Leu Trp Gly Pro Leu Gly Ala
      5                      10                      15
Leu Ala Gln Gly Ala Pro Ala Gly Thr Ala Pro Thr Asp Asp Val
      20                      25                      30
Val Asp Leu Glu Phe Tyr Thr Lys Arg Pro Leu Arg Ser Val Ser
      35                      40                      45
Pro Ser Phe Leu Ser Ile Thr Ile Asp Ala Ser Leu Ala Thr Asp
      50                      55                      60
Pro Arg Phe Leu Thr Phe Leu Gly Ser Pro Arg Leu Arg Ala Leu
      65                      70                      75
Ala Arg Gly Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly Thr Lys
      80                      85                      90
Thr Asp Phe Leu Ile Phe Asp Pro Asp Lys Glu Pro Thr Ser Glu
      95                      100                     105
Glu Arg Ser Tyr Trp Lys Ser Gln Val Asn His Asp Ile Cys Arg
     110                     115                     120
Ser Glu Pro Val Ser Ala Ala Val Leu Arg Lys Leu Gln Val Glu
     125                     130                     135
Trp Pro Phe Gln Glu Leu Leu Leu Leu Arg Glu Gln Tyr Gln Lys
     140                     145                     150

```



Glu Phe Lys Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Met Leu	155	160	165
Tyr Ser Phe Ala Lys Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu	170	175	180
Asn Ala Leu Leu Arg Thr Pro Asp Leu Arg Trp Asn Ser Ser Asn	185	190	195
Ala Gln Leu Leu Leu Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile	200	205	210
Ser Trp Glu Leu Gly Asn Glu Pro Asn Ser Phe Trp Lys Lys Ala	215	220	225
His Ile Leu Ile Asp Gly Leu Gln Leu Gly Glu Asp Phe Val Glu	230	235	240
Leu His Lys Leu Leu Gln Arg Ser Ala Phe Gln Asn Ala Lys Leu	245	250	255
Tyr Gly Pro Asp Ile Gly Gln Pro Arg Gly Lys Thr Val Lys Leu	260	265	270
Leu Arg Ser Phe Leu Lys Ala Gly Gly Glu Val Ile Asp Ser Leu	275	280	285
Thr Trp His His Tyr Tyr Leu Asn Gly Arg Ile Ala Thr Lys Glu	290	295	300
Asp Phe Leu Ser Ser Asp Ala Leu Asp Thr Phe Ile Leu Ser Val	305	310	315
Gln Lys Ile Leu Lys Val Thr Lys Glu Ile Thr Pro Gly Lys Lys	320	325	330
Val Trp Leu Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala Pro	335	340	345
Leu Leu Ser Asn Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys	350	355	360
Leu Gly Leu Ser Ala Gln Met Gly Ile Glu Val Val Met Arg Gln	365	370	375
Val Phe Phe Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe	380	385	390
Glu Pro Leu Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu	395	400	405
Val Gly Pro Arg Val Leu Leu Ser Arg Val Lys Gly Pro Asp Arg	410	415	420
Ser Lys Leu Arg Val Tyr Leu His Cys Thr Asn Val Tyr His Pro	425	430	435
Arg Tyr Gln Glu Gly Asp Leu Thr Leu Tyr Val Leu Asn Leu His	440	445	450
Asn Val Thr Lys His Leu Lys Val Pro Pro Pro Leu Phe Arg Lys	455	460	465
Pro Val Asp Thr Tyr Leu Leu Lys Pro Ser Gly Pro Asp Gly Leu	470	475	480
Leu Ser Lys Ser Val Gln Leu Asn Gly Gln Ile Leu Lys Met Val	485	490	495
Asp Glu Gln Thr Leu Pro Ala Leu Thr Glu Lys Pro Leu Pro Ala	500	505	510
Gly Ser Ala Leu Ser Leu Pro Ala Phe Ser Tyr Gly Phe Phe Val	515	520	525
Ile Arg Asn Ala Lys Ile Ala Ala Cys Ile	530	535	

## (2) INFORMATION FOR SEQ ID NO:45:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2396  
 (E) TYPE: nucleic acid

(C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:45

	TT TCT AGT	8
TGC TTT TAG CCA ATG TCG GAT CAG GTT TTT CAA GCG ACA AAG AGA		53
TAC TGA GAT CCT GGG CAG AGG ACA TCC TAG CTC GGT CAG ATT TGG		98
GCA GGC TCA AGT GAC CAG TGT CTT AAG GCA GAA GGG AGT CGG GGT		143
AGG GTC TGG CTG AAC CCT CAA CCG GGG CTT TTA ACT CAG GGT CTA		188
GTC CTG GCG CCA AAT GGA TGG GAC CTA GAA AAG GTG ACA GAG TGC		233
GCA GGA CAC CAG GAA GCT GGT CCC ACC CCT GCG CGG CTC CCG GGC		278
GCT CCC TCC CCA GGC CTC CGA GGA TCT TGG ATT CTG GCC ACC TCC		323
GCA CCC TTT GGA TGG GTG TGG ATG ATT TCA AAA GTG GAC GTG ACC		368
GCG GCG GAG GGG AAA GCC AGC ACG GAA ATG AAA GAG AGC GAG GAG		413
GGG AGG GCG GGG AGG GGA GGG GCG TAG GGA GGG ACT CCC GGG AGG		458
GGT GGG AGG GAT GGA GCG CTG TGG GAG GGT ACT GAG TCC TGG CGC		503
CAG AGG CGA AGC AGG ACC GGT TGC AGG GGG CTT GAG CCA GCG CGC		548
CGG CTG CCC CAG CTC TCC CGG CAG CGG GCG GTC CAG CCA GGT GGG		593
ATG CTG AGG CTG CTG CTG CTG TGG CTC TGG GGG CCG CTC GGT GCC		638
Met Leu Arg Leu Leu Leu Leu Trp Leu Trp Gly Pro Leu Gly Ala		
5 10 15		
CTG GCC CAG GGC GCC CCC GCG GGG ACC GCG CCG ACC GAC GAC GTG	683	
Leu Ala Gln Gly Ala Pro Ala Gly Thr Ala Pro Thr Asp Asp Val		
20 25 30		
GTA GAC TTG GAG TTT TAC ACC AAG CGG CCG CTC CGA AGC GTG AGT	728	
Val Asp Leu Glu Phe Tyr Thr Lys Arg Pro Leu Arg Ser Val Ser		
35 40 45		
CCC TCG TTC CTG TCC ATC ACC ATC GAC GCC AGC CTG GCC ACC GAC	773	
Pro Ser Phe Leu Ser Ile Thr Ile Asp Ala Ser Leu Ala Thr Asp		
50 55 60		
CCG CGC TTC CTC ACC TTC CTG GGC TCT CCA AGG CTC CGT GCT CTG	818	
Pro Arg Phe Leu Thr Phe Leu Gly Ser Pro Arg Leu Arg Ala Leu		
65 70 75		
GCT AGA GGC TTA TCT CCT GCA TAC TTG AGA TTT GGC GGC ACA AAG	863	
Ala Arg Gly Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly Thr Lys		
80 85 90		
ACT GAC TTC CTT ATT TTT GAT CCG GAC AAG GAA CCG ACT TCC GAA	908	
Thr Asp Phe Leu Ile Phe Asp Pro Asp Lys Glu Pro Thr Ser Glu		
95 100 105		
GAA AGA AGT TAC TGG AAA TCT CAA GTC AAC CAT GAT ATT TGC AGG	953	
Glu Arg Ser Tyr Trp Lys Ser Gln Val Asn His Asp Ile Cys Arg		
110 115 120		
TCT GAG CCG GTC TCT GCT GCG GTG TTG AGG AAA CTC CAG GTG GAA	998	
Ser Glu Pro Val Ser Ala Ala Val Leu Arg Lys Leu Gln Val Glu		
125 130 135		
TGG CCC TTC CAG GAG CTG TTG CTG CTC CGA GAG CAG TAC CAA AAG	1043	
Trp Pro Phe Gln Glu Leu Leu Leu Leu Arg Glu Gln Tyr Gln Lys		
140 145 150		

GAG TTC AAG AAC AGC ACC TAC TCA AGA AGC TCA GTG GAC ATG CTC	1088
Glu Phe Lys Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Met Leu	
155 160 165	
TAC AGT TTT GCC AAG TGC TCG GGG TTA GAC CTG ATC TTT GGT CTA	1133
Tyr Ser Phe Ala Lys Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu	
170 175 180	
AAT GCG TTA CTA CGA ACC CCA GAC TTA CGG TGG AAC AGC TCC AAC	1178
Asn Ala Leu Leu Arg Thr Pro Asp Leu Arg Trp Asn Ser Ser Asn	
185 190 195	
GCC CAG CTT CTC CTT GAC TAC TGC TCT TCC AAG GGT TAT AAC ATC	1223
Ala Gln Leu Leu Leu Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile	
200 205 210	
TCC TGG GAA CTG GGC AAT GAG CCC AAC AGT TTC TGG AAG AAA GCT	1268
Ser Trp Glu Leu Gly Asn Glu Pro Asn Ser Phe Trp Lys Lys Ala	
215 220 225	
CAC ATT CTC ATC GAT GGG TTG CAG TTA GGA GAA GAC TTT GTG GAG	1313
His Ile Leu Ile Asp Gly Leu Gln Leu Gly Glu Asp Phe Val Glu	
230 235 240	
TTG CAT AAA CTT CTA CAA AGG TCA GCT TTC CAA AAT GCA AAA CTC	1358
Leu His Lys Leu Leu Gln Arg Ser Ala Phe Gln Asn Ala Lys Leu	
245 250 255	
TAT GGT CCT GAC ATC GGT CAG CCT CGA GGG AAG ACA GTT AAA CTG	1403
Tyr Gly Pro Asp Ile Gly Gln Pro Arg Gly Lys Thr Val Lys Leu	
260 265 270	
CTG AGG AGT TTC CTG AAG GCT GGC GGA GAA GTG ATC GAC TCT CTT	1448
Leu Arg Ser Phe Leu Lys Ala Gly Gly Glu Val Ile Asp Ser Leu	
275 280 285	
ACA TGG CAT CAC TAT TAC TTG AAT GGA CGC ATC GCT ACC AAA GAA	1493
Thr Trp His His Tyr Tyr Leu Asn Gly Arg Ile Ala Thr Lys Glu	
290 295 300	
GAT TTT CTG AGC TCT GAT GCG CTG GAC ACT TTT ATT CTC TCT GTG	1538
Asp Phe Leu Ser Ser Asp Ala Leu Asp Thr Phe Ile Leu Ser Val	
305 310 315	
CAA AAA ATT CTG AAG GTC ACT AAA GAG ATC ACA CCT GGC AAG AAG	1583
Gln Lys Ile Leu Lys Val Thr Lys Glu Ile Thr Pro Gly Lys Lys	
320 325 330	
GTC TGG TTG GGA GAG ACG AGC TCA GCT TAC GGT GGC GGT GCA CCC	1628
Val Trp Leu Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala Pro	
335 340 345	
TTG CTG TCC AAC ACC TTT GCA GCT GGC TTT ATG TGG CTG GAT AAA	1673
Leu Leu Ser Asn Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys	
350 355 360	

TTG GGC CTG TCA GCC CAG ATG GGC ATA GAA GTC GTG ATG AGG CAG	1718
Leu Gly Leu Ser Ala Gln Met Gly Ile Glu Val Val Met Arg Gln	
365 370 375	
GTG TTC TTC GGA GCA GGC AAC TAC CAC TTA GTG GAT GAA AAC TTT	1763
Val Phe Phe Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe	
380 385 390	
GAG CCT TTA CCT GAT TAC TGG CTC TCT CTT CTG TTC AAG AAA CTG	1808
Glu Pro Leu Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu	
395 400 405	
GTA GGT CCC AGG GTG TTA CTG TCA AGA GTG AAA GGC CCA GAC AGG	1853
Val Gly Pro Arg Val Leu Leu Ser Arg Val Lys Gly Pro Asp Arg	
410 415 420	
AGC AAA CTC CGA GTG TAT CTC CAC TGC ACT AAC GTC TAT CAC CCA	1898
Ser Lys Leu Arg Val Tyr Leu His Cys Thr Asn Val Tyr His Pro	
425 430 435	
CGA TAT CAG GAA GGA GAT CTA ACT CTG TAT GTC CTG AAC CTC CAT	1943
Arg Tyr Gln Glu Gly Asp Leu Thr Leu Tyr Val Leu Asn Leu His	
440 445 450	
AAT GTC ACC AAG CAC TTG AAG GTA CCG CCT CCG TTG TTC AGG AAA	1988
Asn Val Thr Lys His Leu Lys Val Pro Pro Pro Leu Phe Arg Lys	
455 460 465	
CCA GTG GAT ACG TAC CTT CTG AAG CCT TCG GGG CCG GAT GGA TTA	2033
Pro Val Asp Thr Tyr Leu Leu Lys Pro Ser Gly Pro Asp Gly Leu	
470 475 480	
CTT TCC AAA TCT GTC CAA CTG AAC GGT CAA ATT CTG AAG ATG GTG	2078
Leu Ser Lys Ser Val Gln Leu Asn Gly Gln Ile Leu Lys Met Val	
485 490 495	
GAT GAG CAG ACC CTG CCA GCT TTG ACA GAA AAA CCT CTC CCC GCA	2123
Asp Glu Gln Thr Leu Pro Ala Leu Thr Glu Lys Pro Leu Pro Ala	
500 505 510	
GGA AGT GCA CTA AGC CTG CCT GCC TTT TCC TAT GGT TTT TTT GTC	2168
Gly Ser Ala Leu Ser Leu Pro Ala Phe Ser Tyr Gly Phe Phe Val	
515 520 525	
ATA AGA AAT GCC AAA ATC GCT GCT TGT ATA TGA AAA TAA AAG GCA	2213
Ile Arg Asn Ala Lys Ile Ala Ala Cys Ile	
530 535	
TAC GGT ACC CCT GAG ACA AAA GCC GAG GGG GGT GTT ATT CAT AAA	2258
ACA AAA CCC TAG TTT AGG AGG CCA CCT CCT TGC CGA GTT CCA GAG	2303
CTT CGG GAG GGT GGG GTA CAC TTC AGT ATT ACA TTC AGT GTG GTG	2348
TTC TCT CTA AGA AGA ATA CTG CAG GTG GTG ACA GTT AAT AGC ACT	2393
GTG	2396

## (2) INFORMATION FOR SEQ ID NO:46:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 385

(B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46

```
CGGCCGCTGC TGCTGCTGTG GCTCTGGGGG CGGCTCCGTG CCCTGACCCA 50
AGGCACTCCG GCGGGGACCG CGCCGACCAA AGACGTGGTG GACTTGGAGT 100
TTTACACCAA GAGGCTATTC CAAAGCGTGA GTCCCTCGTT CCTGTCCATC 150
ACCATCGACG CCAGTCTGGC CACCGACCCT CGGTCCTCA CCTTCCTGAG 200
CTCTCCACGG CTTCGAGCCC TGTCTAGAGG CTTATCTCCT GCGTACTTGA 250
GATTTGGCGG CACCAAGACT GACTTCCTTA TTTTGTATCC CAACAACGAA 300
CCACCTCTG AAGAAAGAAG TTACTGGCAA TCTCAAGACA ACAATGATAT 350
TTGCGGGTCT GACCGGGTCT CCGCTGACGT GTTGA 385
```

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 541  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47

```
AAATCAGGAC ATATCCTTCA CTTATTTGCC TCTTGGTCAT ATTGGAGGCA 50
TTTGATTCA TTTTAAATAA CCCTCAAAAT AGTGCATGCA AAGTGCTAAG 100
CGTCATTTGC CACATGGTGC CATTAAGTGT CACCACCTGC AGTGGTCTAC 150
TTAGAGAACA CCGCACTGGA TGTTAAACACT GAAGCGCGTG CCCC GCCCTC 200
CCGAGGCTCT GGATCCAGCG TTGAAGCTTG CCCC GCCCTC CCGAGGCTCT 250
GGATCCAGCA CTGGAGCATG CCCC GCCCTC CCGAGGCTCT GGAGCTTGCT 300
AAGGAGTCCG CTCCCTACCG CTGGGGTTTT GCTTTATTCT TATGAATGAC 350
ACCCCTGACC GCTTTCGTCT CAGGGGTACT GTAATGCCTT TTATTTTCAT 400
ATACAAGCTG CGATTTTGGC ATTTCTTATG AAAAAAACC CATAGGAAAA 450
GGCGGGCAGC CTTAGTGAGC TTCCTGCGGG GAGAGGTTTT TCTGTTAGAG 500
CTGGCANGGT CTGCTCATCG ACCATCTTCA GGCCTCGTGC C 541
```